



FNP International Annual Report 2019

Appendices: FNP Analysis of data to 31st December 2018

Prepared by Health Intelligence, PHA, May 2019

Contents

INTRODUCTION	1
CLIENT CASELOAD CHARACTERISTICS	12
1. Referrals and dispositions	13
2. Client engagement and characteristics	16
Summary	16
Age at LMP	17
Gestation at enrolment	23
Enrolled by the end of 16 weeks gestation	25
Enrolled by the end of 28 weeks gestation	27
3. Annual Income	29
4. Ethnicity	32
5. Education, Employment and Training (EET)	34
Intake	34
24 months	36
PROGRAMME ACTIVITY AND FIDELITY	38
6. Dosage	39
Pregnancy Dosage	39
Infancy Dosage	45
Toddlerhood Dosage	51
7. Visit Duration	56
Pregnancy Duration	56
Infancy Duration	63
Toddlerhood Duration	68
8. Visit Content	73
Visit Content Pregnancy Completers	73
Visit Content Infancy completers	80
Visit Content Toddlerhood Completers	87

9. Father Engagement	94
10. Client Parent Engagement	95
11. Client Retention	96
Client Attrition	100
Retention Pregnancy	104
Retention Infancy	106
Retention Toddlerhood	108

SHORT-TERM OUTCOMES **110**

12. Hospital Anxiety and Depression Scale (HADs)	111
13. Breastfeeding	115
14. Tobacco Use	121
15. Alcohol Use	129
16. Illegal Drug Use	134
17. Mastery	136
Mastery at Intake	136
Mastery at 24 Months	138
18. Hospital Admissions due to injury or ingestion	141
Admissions Due to Injury/Ingestion Birth to 6 Months	141
Admissions Due to Injury/Ingestion Birth to 12 Months	143
Admissions 12 to 18 Months	145
Admissions 12 to 24 Months	147
19. ED attendances due to injury or ingestion	149
A&E Birth to 6 Months	149
A&E Birth to 12 months	152
A&E 12 to 18 months	154
A&E 12 to 24 months	156
20. ASQ	158
ASQ Infancy	158
ASQ Toddlerhood	170
ASQ-SE	182
21. Immunisations	188

Immunisations 6 Months	188
Immunisations 12 months	190
Immunisations 18 months	192
Immunisations 24 months	194

List of Tables

Table 1: Referrals and dispositions in total (all clients up to the end of 2018) by site and overall	13
Table 2: Referrals and dispositions in total by year referred (All sites)	14
Table 3: Summary of client characteristics at enrolment for all sites by year enrolled (All sites)	16
Table 4: Age at LMP overall and by site	17
Table 5: Age at LMP – Average, SD and range by year of enrolment (All sites)	17
Table 6: Age at LMP – Average, SD and range by site and year of enrolment	18
Table 7: Aged 19 or under at LMP – overall by year of enrolment (All sites)	20
Table 8: Aged 19 or under at LMP by site and year of enrolment.....	21
Table 9: Gestation at enrolment (weeks) – Average, SD and range for all sites by year of enrolment	23
Table 10: Gestation at enrolment (weeks) – Average, SD and range by site and year of enrolment.....	23
Table 11: Percentage of clients (%) enrolled by end of 16 weeks gestation by year of enrolment (All sites).....	25
Table 12: Percentage of clients (%) enrolled by end of 16 weeks gestation by site and year of enrolment.....	25
Table 13: Percentage of clients (%) enrolled by the end of 28 weeks gestation by year of enrolment (All sites).....	27
Table 14: Percentage of clients (%) enrolled by the end of 28 weeks gestation by site and year of enrolment.....	27
Table 15: Annual Income – All sites	29
Table 16: Ethnicity – All sites	32
Table 17: Education, Employment and Training (EET) Intake – All sites	34
Table 18: Education, Employment and Training (EET) 24 Months – All sites	36
Table 19: Dosage Pregnancy Stage – All sites by year enrolled	39
Table 20: Dosage Pregnancy Stage – BHSCT by year enrolled.....	40
Table 21: Dosage Pregnancy Stage – NHSCT by year enrolled	41
Table 22: Dosage Pregnancy Stage – SEHSCT by year enrolled	42
Table 23: Dosage Pregnancy Stage – SHSCT by year enrolled.....	43
Table 24: Dosage Pregnancy Stage – WHSCT by year enrolled.....	44
Table 25: Dosage Infancy Stage – All sites by year enrolled	45
Table 26: Dosage Infancy Stage – BHSCT by year enrolled	46
Table 27: Dosage Infancy Stage – NHSCT by year enrolled.....	47
Table 28: Dosage Infancy Stage – SEHSCT by year enrolled.....	48
Table 29: Dosage Infancy Stage – SHSCT by year enrolled	49

Table 30: Dosage Infancy Stage – WHSCT by year enrolled	50
Table 31: Dosage Toddlerhood Stage – All sites by year enrolled.....	51
Table 32: Dosage Toddlerhood Stage – BHSCT by year enrolled.....	52
Table 33: Dosage Toddlerhood Stage – NHSCT by year enrolled.....	53
Table 34: Dosage Toddlerhood Stage – SEHSCT by year enrolled	53
Table 35: Dosage Toddlerhood Stage – SHSCT by year enrolled.....	54
Table 36: Dosage Toddlerhood Stage – WHSCT by year enrolled.....	55
Table 37: Pregnancy Duration – All sites by year enrolled	56
Table 38: Pregnancy Duration – By site by year enrolled	57
Table 39: Pregnancy Duration – BHSCT by year enrolled.....	58
Table 40: Pregnancy Duration – NHSCT by year enrolled	59
Table 41: Pregnancy Duration – SEHSCT by year enrolled	60
Table 42: Pregnancy Duration – SHSCT by year enrolled.....	61
Table 43: Pregnancy Duration – WHSCT by year enrolled.....	62
Table 44: Infancy duration – All sites by year enrolled.....	63
Table 45: Infancy duration – By site by year enrolled	64
Table 46: Infancy duration – BHSCT by year enrolled.....	65
Table 47: Infancy duration – NHSCT by year enrolled.....	65
Table 48: Infancy duration – SEHSCT by year enrolled	66
Table 49: Infancy duration – SHSCT by year enrolled.....	66
Table 50: Infancy duration – WHSCT by year enrolled.....	67
Table 51: Toddlerhood Duration – All sites by year of enrolment	68
Table 52: Toddlerhood Duration – By site by year enrolled	69
Table 53: Toddlerhood Duration – BHSCT by year enrolled.....	70
Table 54: Toddlerhood Duration – NHSCT by year enrolled.....	70
Table 55: Toddlerhood Duration – SEHSCT by year enrolled	71
Table 56: Toddlerhood Duration – SHSCT by year enrolled.....	71
Table 57: Toddlerhood Duration – WHSCT by year enrolled.....	72
Table 58: Visit content – Overall active pregnancy completers by site (until end of 2018)	73
Table 59: Visit content – All sites active pregnancy completers by year enrolled (until end of 2018)	73
Table 60: Visit content – BHSCT active pregnancy completers by year enrolled (until end of 2018)	75

Table 61: Visit content – NHSCT active pregnancy completers by year enrolled (until end of 2018)	76
Table 62: Visit content – SEHSCT active pregnancy completers by year enrolled (until end of 2018)	77
Table 63: Visit content – SHSCT active pregnancy completers by year enrolled (until end of 2018)	78
Table 64: Visit content – WHSCT active pregnancy completers by year enrolled (until end of 2018)	79
Table 65: Visit content – active infancy completers by site (until end of 2018)	80
Table 66: Visit content - All sites active infancy completers by year of enrolment (until end of 2018)	80
Table 67: Visit content – BHSCT active infancy completers by year of enrolment (until end of 2018).....	82
Table 68: Visit content – NHSCT active infancy completers by year of enrolment (until end of 2018)	83
Table 69: Visit content – SEHSCT active infancy completers by year of enrolment (until end of 2018)	84
Table 70: Visit content – SHSCT active infancy completers by year of enrolment (until end of 2018).....	85
Table 71: Visit content – WHSCT active infancy completers by year of enrolment (until end of 2018).....	86
Table 72: Visit Content – Toddlerhood completers by site (up to end of December 2018).....	87
Table 73: Visit Content – Toddlerhood (All sites) completers by year of enrolment (up to end of December 2018)	87
Table 74: Visit Content – BHSCT toddlerhood completers by year of enrolment (up to end of December 2018).....	89
Table 75: Visit Content – NHSCT toddlerhood completers by year of enrolment (up to end of December 2018).....	90
Table 76: Visit Content – SEHSCT toddlerhood completers by year of enrolment (up to end of December 2018).....	91
Table 77: Visit Content – SHSCT toddlerhood completers by year of enrolment (up to end of December 2018).....	92
Table 78: Visit Content – WHSCT toddlerhood completers by year of enrolment (up to end of December 2018).....	93
Table 79: Father Engagement – All sites and by stage for active completers by 31.12.2018.....	94
Table 80: Client parent engagement – All sites and by stage for active completers by 31.12.2018	95
Table 81: Client Retention by Year – All sites	96
Table 82: Client Retention by Year – All sites	97
Table 83: Attrition by Stage and Cumulative Attrition for Stage Completers (All sites).....	100
Table 84: Total Attrition by Stage for Stage Completers (All sites)	101
Table 85: Total Attrition by Stage for Stage Completers BHSCT	101
Table 86: Total Attrition by Stage for Stage Completers NHSCT.....	102
Table 87: Total Attrition by Stage for Stage Completers SEHSCT.....	102
Table 88: Total Attrition by Stage for Stage Completers SHSCT	103
Table 89: Total Attrition by Stage for Stage Completers WHSCT	103
Table 90: Client Retention by Site and by Year – Pregnancy Stage.....	104
Table 91: Client Retention by Site and by Year – Infancy Stage	106

Table 92: Client Retention by Site and by Year – Toddlerhood Stage	108
Table 93: HADS* Anxiety – All sites	111
Table 94: HADS Depression – All sites	111
Table 95: Breastfeeding – All sites	115
Table 96: Breastfeeding – BHSCT and all sites.....	116
Table 97: Breastfeeding – NHSCT and all sites	117
Table 98: Breastfeeding – SEHSCT and all sites	118
Table 99: Breastfeeding – SHSCT and all sites.....	119
Table 100: Breastfeeding – WHSCT and all sites.....	120
Table 101: Change in self-reported smoking during pregnancy (intake to 36 weeks) – All sites	121
Table 102: Change in self-reported alcohol use during pregnancy (intake to 36 weeks) – All sites	129
Table 103: Change in self-reported drug use during pregnancy (intake to 36 weeks) – All sites	134
Table 104: Mastery at intake – All sites.....	136
Table 105: Mastery at 24 Months – All sites.....	138
Table 106: Admissions Birth to 6 Months – All sites	141
Table 107: Admissions Birth to 12 Months – All sites	143
Table 108: Admissions 12 to 18 Months – All sites	145
Table 109: Admissions 12 to 24 Months – All sites	147
Table 110: A&E Birth to 6 Months – All sites	149
Table 111: A&E Birth to 12 months – All sites	152
Table 112: A&E 12 to 18 months – All sites	154
Table 113: A&E 12 to 24 months – All sites	156
Table 114: ASQ Infancy – All sites	158
Table 115: ASQ Infancy – BHSCT and all sites.....	160
Table 116 ASQ Infancy- NHSCT and all sites	162
Table 117: ASQ Infancy – SEHSCT and all sites	164
Table 118: ASQ Infancy – SHSCT and all sites.....	166
Table 119: ASQ Infancy – WHSCT and all sites.....	168
Table 120: ASQ Toddlerhood – All sites	170
Table 121: ASQ Toddlerhood – BHSCT and all sites	172
Table 122: ASQ Toddlerhood – NHSCT and all sites.....	174

Table 123: ASQ Toddlerhood – SEHSCT and all sites..... 176

Table 124: ASQ Toddlerhood – SHSCT and all sites 178

Table 125: ASQ Toddlerhood – WHSCT and all sites 180

Table 126: ASQ-SE 6, 12, 18 and 24 months – All sites 182

Table 127: ASQ-SE 6, 12, 18 and 24 months – BHSCT and all sites 183

Table 128: ASQ-SE 6, 12, 18 and 24 months – NHSCT and all sites..... 184

Table 129: ASQ-SE 6, 12, 18 and 24 months – SEHSCT and all sites..... 185

Table 130: ASQ-SE 6, 12, 18 and 24 months – SHSCT and all sites 186

Table 131: ASQ-SE 6, 12, 18 and 24 months – WHSCT and all sites 187

Table 132: Immunisations 6 Months – All sites..... 188

Table 133: Immunisations 12 months – All sites..... 190

Table 134: Immunisations 18 months – All sites..... 192

Table 135: Immunisations 24 months – All sites..... 194

Introduction

The following tables are based on overall data for the five Northern Ireland FNP sites up to the end of 31st December 2018 to inform the 2019 International Annual Report.

The tables below provide a summary of the key data pertaining to client caseload characteristics, programme activity and fidelity and short term outcomes.

Data Overview (Overall, for all sites)

Client Caseload Characteristics	
Referrals	<ul style="list-style-type: none"> Overall, 72.1% (N=918) of definitely eligible clients offered the programme prior to the end of December 2018 (referred by 31.12.2018) were enrolled on the programme (79.6% BHSCT (N=183), 66.7% NHSCT (N=112), 88.1% SEHSCT (N=140), 63.4% SHSCT (N=194) and 70.5% WHSCT (N=289). The fidelity goal for % enrolled who are offered the programme is 75%. In 2018, 62.9% (N= 124) of definitely eligible clients offered the programme were enrolled.
Client Characteristics at Enrolment	<ul style="list-style-type: none"> Overall, of the 1931 clients who were referred to the programme prior to end of December 2018, 913 clients were enrolled (had valid completed visit) on the programme prior to the end of December 2018. 43.6% of clients were enrolled by the end of the 16th week gestation and 99.9% of clients were enrolled by the end of the 28th week gestation. The mean gestational age (week's gestation) at enrolment was 18.5 and this ranged from 7 weeks to 29 weeks gestation. Overall, across all sites 99.8% (N=911) of clients were aged 19 or under at LMP. Two clients (0.2%) were aged 20 at LMP. Overall, across all sites (N=913) the average age at LMP was 17.2 and this ranged from 12.8 to 20.5.
Annual Income	<ul style="list-style-type: none"> Annual income was recorded for 94% (N= 857) clients at intake. 79.8% (N=684) had an annual income of less than £13,000. However, given the level of completeness (i.e. 48.4% (N=415) missing data) in relation to the source of income as household or self only it is not possible to make comparisons against currently available poverty indicators. Relative poverty can be defined as living in a household with an income below 60% of the average income. Annual income in N.I, before housing costs, was £23,504 per year in 2016/2017 and 60% of this is £14,102.

	<p>Approximately 18% of the population in Northern Ireland were living in relative poverty before housing costs in 2016/2017 (Northern Ireland Poverty Bulletin 2016/2017 [NISRA, 2018]). We have presented comparison data for clients with income under £13,000 due to this most closely matching the £14,102 poverty cut off.</p>
Ethnicity	<ul style="list-style-type: none"> • Overall, ethnicity was recorded for 94% (N=857) of the 912 who were scheduled to have their ethnicity recorded by the report end date (31.12.18). • For the 857 clients who had their ethnicity recorded, 46% (N= 394) reported that they were White British, 47.4% (N=406) reported that they were White Irish, 3.7% (N=32) reported that they were White Other, 2.9% (N=25) reported that they were Other ethnic group.
EET	<p>Intake</p> <ul style="list-style-type: none"> • Education and employment status was recorded for 94% (N=857) of clients at intake. At intake 46.6% (N=399) clients were in education, 19.6% (N=168) of clients were in employment and 9.6% (N=82) of clients were in both education and employment. • At intake, 56.6% (N=485) clients were either in education or employment (EET) and 43.4% (N=372) clients were not in education or employment (NEET). <p>24 months</p> <ul style="list-style-type: none"> • Education and employment status was recorded for 84.5% (N=430) of clients at 24 months. At 24 months 30.9% (N=133) clients were in education, 21.2% (N=91) of clients were in employment and 7.9% (N=34) of clients were in both education and employment. • At 24 months, 44.2% (N=190) clients were either in education or employment (EET) and 55.8% (N=240) clients were not in education or employment (NEET).

Programme Activity and Fidelity	
Pregnancy Dosage	<ul style="list-style-type: none"> • 845 active clients completed the pregnancy stage by the end of 31.12.2018. • Overall, clients were scheduled to have an average of 12.1 visits and they completed an average of 9.6 valid visits in pregnancy. • Overall to date for all clients who enrolled since 2010, 79.5% of expected visits (based on gestational age) and 68.4% of expected visits (based on 14 expected visits) were achieved by the end of 2018. • For clients who enrolled in 2018, 79.5% of expected visits (based on gestational age) and 60.3% of expected visits (based on 14 expected visits) were achieved by the end of 2018.
Infancy Dosage	<ul style="list-style-type: none"> • 660 active clients completed the infancy stage by the end of 31.12.2018.

	<ul style="list-style-type: none"> Overall, clients were scheduled to have an average of 28 visits and they completed an average of 20 valid visits in infancy. Overall to date for all clients who enrolled since 2010, 71.3% of expected visits were achieved by the end of 2018. For clients who enrolled in 2017, 63.6% of expected infancy visits were achieved by the end of 2018.
Toddlerhood dosage	<ul style="list-style-type: none"> 488 active clients completed the toddlerhood stage by the end of 31.12.18 Overall, clients were scheduled to have an average of 22 visits and they completed an average of 14 valid visits in toddlerhood. Overall to date for all clients who enrolled since 2010, 63.7% of expected visits were achieved by the end of 2018. For clients who enrolled in 2016, 61.4% of expected toddlerhood visits were achieved by the end of 2018.
Visit Duration Pregnancy	<ul style="list-style-type: none"> Overall, the average visit duration for valid pregnancy visits (N= 8095 visits) for pregnancy completers (N=845) was 1 hour and 10 minutes. This ranged from 15 minutes to 5 hours and 33 minutes. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.
Visit Duration Infancy	<ul style="list-style-type: none"> Overall, the average visit duration for valid infancy visits (N= 13153 visits) for infancy completers (N= 660) was one hour and six minutes. This ranged from 15 minutes to six hours. The fidelity stretch goal for length of visits is that, on average, infancy visits should be 60 minutes or more.
Visit Duration Toddlerhood	<ul style="list-style-type: none"> Overall, the average visit duration for valid toddlerhood visits (N= 6834 visits) for toddlerhood completers (N=488) was one hour and five minutes. This ranged from 15 minutes to eight hours. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.
Visit content pregnancy	<ul style="list-style-type: none"> Overall (N=8095 valid visits), life course development (12%) was within the visit content goal (10-15%), family and friends (15.1%) was just above the visit content goal (10-15%), maternal role (27.7%) was above the visit content goal (23-25%), personal health (36.2%) was within the visit content goal (35-40%) and environmental health (9%) was above the visit content goal (5-7%).
Visit content infancy	<ul style="list-style-type: none"> Overall (N=13153 valid visits), life course development (11.1%) was within the visit content goal (10-15%), family and friends (13.4%) was within the visit content goal (10-15%), maternal role (42.7%) was below the visit content goal (45-50%), personal health (21.8%) was above the visit content goal (14-20%) and environmental health (11%) was above the visit content goal (7-10%).

Visit content toddlerhood	<ul style="list-style-type: none"> Overall (N= 6834 valid visits), life course development (14.6%) was below the visit content goal (18-20%), family and friends (14.3%) was within the visit content goal (10-15%), maternal role (40.1%) was within the visit content goal (40-45%), personal health (19.2%) was above the visit content goal (10-15%) and environmental health (11.9%) was above the visit content goal (7-10%).
Father engagement	<ul style="list-style-type: none"> Overall, father engagement was 16.6% for pregnancy completers, 16.5% for infancy completers and 12.7% for toddlerhood completers.
Parent Engagement	<ul style="list-style-type: none"> Overall, client parent engagement was 18.7% for pregnancy completers, 17.3% for infancy completers and 10.8% for toddlerhood completers.
Client retention	<ul style="list-style-type: none"> 98.3% (845 active clients of 860 clients scheduled to complete the stage) clients completed the pregnancy stage. 89.9% (660 active clients of 734 clients scheduled to complete the stage) clients completed the infancy stage. 83% (488 active clients of 588 clients scheduled to complete the stage) clients completed the toddlerhood stage.
Attrition	<p>Overall, of the 913 clients ever enrolled, attrition by the 31.12.2018 is as follows:</p> <ul style="list-style-type: none"> 15 (1.7%) left during pregnancy (11 were leavers and 4 became inactive); 63 (8.6%) left during infancy (29 were leavers and 34 became inactive); and 41 (7%) left during toddlerhood (17 were leavers and 27 became inactive, one of these clients became inactive, returned to the programme and then left the programme, 2 further clients became inactive and then left in toddlerhood).

Short Term Outcomes	
HADS Pregnancy	<p>Anxiety</p> <ul style="list-style-type: none"> Overall, anxiety (HADS) scores were recorded for 85.3% (N=736) clients in pregnancy. Of these, 79.8% (N=587) were found to have an unlikely presence of anxiety, 12% (N=88) were found to have borderline presence of anxiety, 6% (N=44) were found to have probable presence of anxiety and 2.2% (N=16) were found to have a high likelihood of anxiety in pregnancy. <p>Depression</p> <ul style="list-style-type: none"> Overall, depression (HADS) scores were recorded for 85.1% (N=734) clients in pregnancy. Of these, 90.5% (N=664) were found to have an unlikely presence of depression, 6.4% (N=47) were found to

	<p>have borderline presence of depression, 2.7% (N=20) were found to have probable presence of depression and 0.3% (N=2) clients were found to have a high likelihood of depression in pregnancy.</p>
Breastfeeding	<ul style="list-style-type: none"> • Breastfeeding initiation was recorded for 94.9% (N=785) of clients with infants. Of these 43.9% (N=345) reported initiating breastfeeding. • Breastfeeding status was recorded for 92.7% (N=745) of clients who had reached 6 weeks infancy. Of these, 8.1% (N=60) reported that they were still breastfeeding at 6 weeks. • Breastfeeding status was recorded for 84.4% (N=628) of clients who had reached 6 months infancy. Of these, 3% (N=19) reported that they were still breastfeeding at 6 months. • Exclusive breastfeeding status at either 6 weeks or 6 months was recorded for 94.9% (N=763) clients. For these clients, the average duration that clients exclusively breastfed their infant was 3.9 weeks.
Tobacco Use	<p>Intake</p> <ul style="list-style-type: none"> • Information on smoking during pregnancy was recorded for 93.5% (N= 854) of clients at intake. Of these, 40.9 (N=349) reported that they had smoked at some time during pregnancy at intake. • Information on smoking during the last 48 hours was recorded for 93.5% (N= 854) of clients at intake. Of these, 27.4% (N=234) reported that they had smoked during the last 48 hours at intake. • The mean number of cigarettes smoked was 10.1 for those who smoked during the last 48 hours at intake. • The mean number of cigarettes smoked was 16.1 for those who smoked more than 5 cigarettes per day during the last 48 hours at intake. <p>36 weeks pregnancy</p> <ul style="list-style-type: none"> • Information on smoking during pregnancy was recorded for 85.7% (N= 740) of clients at 36 weeks pregnancy. Of these, 39.6% (N=293) reported that they had smoked at some time during the pregnancy at 36 weeks pregnancy • Information on smoking during the last 48 hours was recorded for 85.7% (N= 740) of clients at 36 weeks pregnancy. Of these, 22.7% (N=168) reported that they had smoked during the last 48 hours at 36 weeks pregnancy. • The mean number of cigarettes smoked was 9.1 for those who smoked during the last 48 hours at 36 weeks pregnancy. • The mean number of cigarettes smoked was 15.7 for those who smoked more than 5 cigarettes per day during the last 48 hours at 36 weeks pregnancy. <p>Intake and 36 weeks</p> <ul style="list-style-type: none"> • 727 clients had information on smoking status recorded at both intake and 36 weeks pregnancy. For these

	<p>clients:</p> <ul style="list-style-type: none"> ○ a - 5% change in smoking in the last 48 hours was achieved between intake and 36 weeks. ○ a - 18% change in the proportion of clients smoking in in the last 48 hours was achieved between intake and 36 weeks. ○ 17.7% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks. ○ 75.3% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks for clients who smoked more than 5 cigarettes per day at intake. ○ 7.3% (N=53) clients smoked more at 36 weeks than at intake. ○ 1.7% (N=12) clients smoked at 36 weeks who denied smoking at intake.
Alcohol Use	<p>Intake</p> <ul style="list-style-type: none"> ● Information on alcohol use was recorded at intake for 93.5% of clients (N=854) at intake. Of these, 1.4% (N=12) reported drinking alcohol in the last 14 days at intake. 0.4% (N=3) clients reported drinking excessively in the last 14 days at intake. <p>36 weeks pregnancy</p> <ul style="list-style-type: none"> ● Information on alcohol use was recorded at 36 weeks pregnancy for 85.7% (N=740) clients. Of these, 0.8% (N=6) reported drinking alcohol in the last 14 days at 36 weeks. 0.3% (N=2) clients reported drinking excessively in the last 14 days at 36 weeks pregnancy. <p>Intake and 36 weeks</p> <ul style="list-style-type: none"> ● Information on alcohol use was recorded at both intake and 36 weeks for 727 clients. For these clients, there was: <ul style="list-style-type: none"> ○ -44.4% change in the proportion of clients drinking alcohol between intake and 36 weeks. ○ -0.6% change in the number of clients drinking alcohol between intake and 36 weeks. ○ -50% change in the proportion of clients drinking alcohol excessively between intake and 36 weeks. ○ -0.1% change in the number of clients drinking alcohol excessively between intake and 36 weeks. ○ 0.4% (N=3) clients started drinking between intake and 36 weeks. ○ 1% (N=7) clients stopped drinking between intake and 36 weeks. ● Please note the small numbers of clients who reported drinking alcohol when interpreting the figures for % change in proportion of clients drinking alcohol.
Illegal drug use	<ul style="list-style-type: none"> ● Information on illegal drug use was recorded for 93.5% (N=854) clients at intake and 85.7% (N=740) clients at 36 weeks pregnancy. 0.1% (N=1) client reported illegal drug use during pregnancy or the last 14 days at intake. 0.1% (N=1) client reported illegal drug use in last 14 days at 36 weeks.

	<ul style="list-style-type: none"> Information on illegal drug use was recorded for 727 clients at both intake and 36 weeks. For these clients, there was a 0.1% change in the number of clients who reported using illegal drugs between intake and 36 weeks.
Mastery	<p>Intake</p> <ul style="list-style-type: none"> Overall, mastery was recorded for 93.5% (N=854) of clients at intake. The mean mastery score was 22 (Range= 9 - 28). At intake 16.4% clients (N=140) had low mastery and 83.6% (N=714) of clients did not have low mastery. <p>24 months</p> <ul style="list-style-type: none"> Overall, mastery was recorded for 76.2% (N=377) of clients at 24 months. The mean mastery score was 22.7 (Range= 13 - 28). At 24 months 12.7% clients (N=48) had low mastery and 87.3% (N=329) of clients did not have low mastery. There was a reduction (-2.2%) in the number of clients who had low mastery from intake to 24 months for clients who had mastery recorded at both time points (N=362). There was also an increase in the number of clients who had higher mastery scores from intake to 24 months as 45.3% (N=164) of clients across all sites had a higher mastery score at 24 months than at intake.
Hospital Admissions	<p>6 months</p> <ul style="list-style-type: none"> Overall, 748 infants reached 6 months infancy and information on hospital admissions was recorded for 84.5% (N=632) of these clients. Two infants (0.3%) were recorded with having 1 hospital admission due to injury/ingestion between birth and 6 months. Overall, the mean hospital admission due to injury or ingestion between birth and 6 months is 0.003. <p>12 months</p> <ul style="list-style-type: none"> Overall, 671 infants reached 12 months infancy and information on hospital admissions was recorded for 81.7% (N=548) of these clients. Three infants (0.5%) had 1 hospital admission due to injury or ingestion between birth and 12 months. No infants recorded 2 or more hospital admission due to injury or ingestion between birth and 12 months. Overall, the mean hospital admission due to injury or ingestion between birth and 12 months is 0.01. <p>18 months</p> <ul style="list-style-type: none"> Overall, 580 infants reached 18 months toddlerhood and information on hospital admissions was recorded for 74.8% (N=434) of these clients.

	<ul style="list-style-type: none"> • Six infants (1.4%) were recorded with having 1 hospital admission due to injury or ingestion between 12 and 18 months. • One infant (0.2%) was recorded with having 2 or more hospital admissions due to injury or ingestion between 12 and 18 months. • Overall, the mean hospital admissions due to injury or ingestion between 12 and 18 months is 0.02. <p>24 months</p> <ul style="list-style-type: none"> • Overall, 510 infants reached 24 months toddlerhood and information on hospital admissions was recorded for 81.8% (N=417) of these clients. • Nine infants (2.2%) were recorded with having 1 hospital admission due to injury or ingestion between 12 and 24 months. • No infants recorded 2 or more hospital admission due to injury or ingestion between 12 and 24 months. • Overall, the mean hospital admissions due to injury or ingestion between 12 and 24 months is 0.02.
A&E Attendances	<p>6 months</p> <ul style="list-style-type: none"> • 748 infants reached 6 months by the end of December 2018 and information on A&E visits was recorded for 84.5% (N=632) of these clients. • Overall, 4.1% (N=26) infants had 1 A&E visit due to injury or ingestion between birth and 6 months. • Overall, 0.2% (N=1) infant had 2 or more A&E visits due to injury or ingestion between birth and 6 months. • Overall, the mean visits to A&E due to injury or ingestion between birth and 6 months is 0.04. <p>12 months</p> <ul style="list-style-type: none"> • 671 infants reached 12 months by the end of December 2018 and information on A&E visits was recorded for 81.7% (N=548) of these clients. • Overall, 4.8% (N=32) infants had 1 A&E visit due to injury or ingestion between birth and 12 months. • Overall, 0.4% (N=3) infant had 2 or more A&E visits due to injury or ingestion between birth and 12 months. • Overall, the mean visits to A&E due to injury or ingestion between birth and 12 months is 0.06. <p>18 months</p> <ul style="list-style-type: none"> • 580 infants reached 18 months by the end of December 2018 and information on A&E visits was recorded for 74.8% (N=434) of these clients. • Overall, 4.6% (N=20) infants had 1 A&E visit due to injury or ingestion between 12 and 18 months. • Overall, 0.9% (N=4) infant had 2 or more A&E visits due to injury or ingestion between 12 and 18 months. • Overall, the mean visits to A&E due to injury or ingestion between 12 and 18 months is 0.07.

	<p>24 months</p> <ul style="list-style-type: none"> • 510 infants reached 24 months by the end of December 2018 and information on A&E visits was recorded for 81.8% (N=417) of these clients. • Overall, 5.8% (N=24) infants had 1 A&E visit due to injury or ingestion between 12 and 24 months. • Overall, 0.7% (N=3) infant had 2 or more A&E visits due to injury or ingestion between 12 and 24 months. • Overall, the mean visits to A&E due to injury or ingestion between 12 and 24 months is 0.07.
ASQ	<p>4 Months</p> <ul style="list-style-type: none"> • Overall, 748 infants reached 4 months infancy by the end of December 2018 and ASQ data was recorded for 83.2% (N=622) of these clients. • Communication: 1.6% (N=10) infants had a score below the seek consultation cut-off (34.60). The overall mean for communication is 56.3. • Gross motor: 4.8% (N=30) infants had a score below the seek consultation cut-off (38.41). The overall mean for gross motor is 54.7. • Fine motor: 2.1% (N=13) infants had a score below the seek consultation cut-off (29.62). The overall mean for fine motor is 54.8. • Problem solving: 1.6% (N=10) infants had a score below the seek consultation cut-off (34.98). The overall mean for problem solving is 56.2. • Personal Social: 1.9% (N=12) infants had a score below the seek consultation cut-off (33.16). The overall mean for personal social is 55.9. <p>10 Months</p> <ul style="list-style-type: none"> • Overall, 671 infants reached 10 months infancy by the end of December 2018 and ASQ data was recorded for 80.9% (N=543) of these clients. • Communication: 1.3% (N=7) infants had a score below the seek consultation cut-off (22.87). The overall mean for communication is 55.7. • Gross motor: 15.8% (N=86) infants had a score below the seek consultation cut-off (30.07). The overall mean for gross motor is 50.1. • Fine motor: 3.5% (N=19) infants had a score below the seek consultation cut-off (37.97). The overall mean for fine motor is 56.0. • Problem solving: 3.7% (N=20) infants had a score below the seek consultation cut-off (32.51). The overall mean for problem solving is 54.7. • Personal Social: 1.7% (N=9) infants had a score below the seek consultation cut-off (27.25). The overall mean for personal social is 53.9.

	<p>14 Months</p> <ul style="list-style-type: none"> • Overall, 580 infants reached 14 months toddlerhood by the end of December 2018 and ASQ data was recorded for 73.4% (N=426) of these clients. • Communication: 2.3% (N=10) infants had a score below the seek consultation cut-off (17.40). The overall mean for communication is 53.5. • Gross motor: 5.3% (N=21) infants had a score below the seek consultation cut-off (25.8). The overall mean for gross motor is 54.5. • Fine motor: 2.5% (N=11) infants had a score below the seek consultation cut-off (23.06). The overall mean for fine motor is 54.4. • Problem solving: 2.8% (N=12) infants had a score below the seek consultation cut-off (22.56). The overall mean for problem solving is 52.6. • Personal Social: 1.6% (N=7) infants had a score below the seek consultation cut-off (23.18). The overall mean for personal social is 55.4. <p>20 Months</p> <ul style="list-style-type: none"> • Overall, 510 infants reached 20 months toddlerhood by the end of December 2018 and ASQ data was recorded for 78.8% (N=402) of these clients. • Communication: 5.2% (N=21) infants had a score below the seek consultation cut-off (20.50). The overall mean for communication is 52.5. • Gross motor: 3.2% (N=13) infants had a score below the seek consultation cut-off (39.89). The overall mean for gross motor is 56.3. • Fine motor: 2.7% (N=11) infants had a score below the seek consultation cut-off (36.05). The overall mean for fine motor is 55.6. • Problem solving: 1.7% (N=7) infants had a score below the seek consultation cut-off (28.84). The overall mean for problem solving is 53.4. • Personal Social: 2.7% (N=11) infants had a score below the seek consultation cut-off (33.36). The overall mean for personal social is 55.6.
ASQ-SE	<ul style="list-style-type: none"> • Overall, 748 infants reached 6 months infancy by the end of December 2018 and ASQ-SE data was recorded for 82.8% (N=619) of these clients. 0.8% (N=5) infants scored above the seek consultation cut-off (45). The overall mean ASQ-SE score is 7.8. • Overall, 671 infants reached 12 months infancy by the end of December 2018 and ASQ-SE data was recorded for 79.7% (N=535) of these clients. 0.6% (N=3) infants scored above the seek consultation cut-off (48). The overall mean ASQ-SE score is 7.6.

	<ul style="list-style-type: none"> • Overall, 580 infants reached 18 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 73.3% (N=425) of these clients. 1.4% (N=6) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 9.8. • Overall, 510 infants reached 24 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 80% (N=408) of these clients. 2.9% (N=12) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 11.6.
Immunisations	<p>6 months</p> <ul style="list-style-type: none"> • Overall, 748 infants reached 6 months infancy and immunisation status was recorded for 84.5% (N=632) of these clients. 94.9% (N= 600) had up-to-date immunisations (written record or mother’s self-report) and 5.1% (N=32) did not have records reported as up-to-date. <p>12 months</p> <ul style="list-style-type: none"> • Overall, 671 infants reached 12 months infancy and immunisation status was recorded for 81.7% (N=548) of these clients. • 93.4% (N=512) had up-to-date immunisations (written record or mother’s self-report) and 6.6% (N=36) did not have records recorded as up-to-date. <p>18 months</p> <ul style="list-style-type: none"> • Overall, 580 infants reached 18 months toddlerhood and immunisation status was recorded for 74.7% (N=433) of these clients. 96.8% (N=419) had up-to-date immunisations (written record or mother’s self-report) and 3.2% (N=14) did not have records reported as up-to-date. <p>24 months</p> <ul style="list-style-type: none"> • Overall, 510 infants reached 24 months toddlerhood and immunisation status was recorded for 81.8% (N=417) of these clients. 98.6% (N=411) had up-to-date immunisations (written record or mother’s self-report) and 1.4% (N=6) did not have records reported as up-to-date.

More detailed tables on the FNP data are provided in the following sections. Data for individual FNP sites are also provided for reference.

Client Caseload Characteristics

Referrals and dispositions

Table 1: Referrals and dispositions in total (all clients up to the end of 2018) by site and overall

	Overall (All sites)	BHSCT	NHSCT	SEHSCT	SHSCT	WHSCT
Total Referrals	1931	405	191	168	517	650
Disposition						
1 Enrolled in programme	918	183	112	140	194	289
2 Refused participation in programme	323	45	56	19	85	118
3 Unable to locate	96	39	6	5	16	30
4a Didn't meet criteria: Employed/Qualifications	2	-	-	-	-	2
4b Didn't meet criteria: > 28 wks. pregnant	63	12	7	-	22	22
4c Didn't meet criteria: multiple problems/mental health	3	-	-	-	1	2
4d Didn't meet criteria: miscarried/other medical crisis	44	8	3	-	11	22
4e Didn't meet criteria: incorrect geographical area	106	49	2	4	45	6
4f Didn't meet criteria: unknown/other	38	5	-	-	7	26
6 Quota full for the month	306	62	5	-	109	130
7 Eligible Awaiting Enrolment	32	2	-	-	26	3
Total Enrolled	918	183	112	140	194	289
Total Referrals	1931	405	191	168	517	650
Total possibly eligible*	1675	331	179	164	431	570
Enrolled as a percentage of possibly eligible (%)	54.8	55.3	62.6	85.4	45.0	50.7
Total definitely eligible**	1273	230	168	159	306	410
Enrolled as a percentage of definitely eligible (%)	72.1	79.6	66.7	88.1	63.4	70.5

*Total possibly eligible calculated as: total referrals minus not eligible (not eligible [all reasons [4a-4f]])

** Total definitely eligible calculated as: total referrals minus not eligible (not eligible [all reasons [4a-4f] and not recruited unable to locate [3], [language issues [5], Quota full for the month [6]])

All sites

- Overall, 72.1% (N=918) of definitely eligible clients offered the programme prior to the end of December 2018 (referred by 31.12.2018) were enrolled on the programme (79.6% BHSCT (N=183), 66.7% NHSCT (N=112), 88.1% SEHSCT (N=140), 63.4% SHSCT (N=194) and 70.5% WHSCT (N=289). The fidelity goal for % enrolled who are offered the programme is 75%.
- In 2018, 62.9% (N= 124) of definitely eligible clients offered the programme were enrolled.

Table 2: Referrals and dispositions in total by year referred (All sites)

	Total 2010	Total 2011	Total 2012	Total 2013	Total 2014	Total 2015	Total 2016	Total 2017	Total 2018	All Sites
Total Referrals (All Sites)	13	136	95	287	246	250	364	265	275	1931
Disposition										
1 Enrolled in programme	9	77	38	152	101	92	191	134	124	918
2 Refused participation in programme		25	13	57	36	33	71	47	41	323
3 Unable to locate	1	9	2	6	17	6	32	7	16	96
4a Didn't meet criteria: Employed/Qualifications									2	2
4b Didn't meet criteria: > 28 wks. pregnant	3	5	4	12	8	7	13	8	3	63
4c Didn't meet criteria: multiple problems/mental health	-	-	-	2	-	-	-	1	-	3
4d Didn't meet criteria: miscarried/other medical crisis	-	4	7	10	8	-	8	5	2	44
4e Didn't meet criteria: incorrect geographical area	-	3	3	16	20	13	19	15	17	106
4f Didn't meet criteria: unknown/other	-	9	2	6	11	3	2	3	2	38
6 Quota full for the month	-	4	26	26	45	96	28	45	36	306
7 Eligible Awaiting Enrolment	-	-	-	-	-	-	-	-	32	32
Total Enrolled	9	77	38	152	101	92	191	134	124	918
Total Referrals	13	136	95	287	246	250	364	265	275	1931
Total possibly eligible *	10	115	79	241	199	227	322	233	249	1675
Enrolled as a percentage of possibly eligible (%)	90	67.0	48.1	63.1	50.8	40.5	59.3	57.5	49.8	54.8
Total definitely eligible **	9	102	51	209	137	125	262	181	197	1273
Enrolled as a percentage of definitely eligible (%)	100	75.5	74.5	72.7	73.7	73.6	72.9	74.0	62.9	72.1

*Total possibly eligible calculated as: total referrals minus not eligible (not eligible [all reasons [4a-4f]])

** Total definitely eligible calculated as: total referrals minus not eligible (not eligible [all reasons [4a-4f] and not recruited unable to locate [3], [language issues [5], Quota full for the month [6]])

All sites

- 62.9% (N=124) of definitely eligible clients referred to the programme in 2018 were enrolled by 31/12/2018.
- 74% (N=134) of definitely eligible clients referred to the programme in 2017 were enrolled by 31/12/2017.
- 72.9% (N=191) of definitely eligible clients referred to the programme in 2016 were enrolled by 31/12/2016.

- 73.6% (N=92) of definitely eligible clients referred to the programme in 2015 were enrolled by 31/12/2015.
- 73.7% (N=101) of definitely eligible clients referred to the programme in 2014 were enrolled by 31/12/2014.
- 72.7% (N=152) of definitely eligible clients referred to the programme in 2013 were enrolled by 31/12/2013.
- 74.5% (N=38) of definitely eligible clients referred to the programme in 2012 were enrolled by 31/12/2012.
- 75.5% (N=77) of definitely eligible clients referred to the programme in 2011 were enrolled by 31/12/2011.
- 100% (N=9) of definitely eligible clients referred to the programme in 2010 were enrolled by 31/12/2010.
- Overall, 72.1% (N=918) of definitely eligible clients referred to the programme by the end of 2018.

Client engagement and characteristics

Summary

Table 3: Summary of client characteristics at enrolment for all sites by year enrolled (All sites)

All Sites	Enrolled N	% of those eligible* offered the programme who enrol	% Enrolled by end of 16 th week gestation	% Enrolled by end of 28 th week gestation	Mean gestational age at enrolment (weeks)
2010	7	100	14.3	100	26.4 (SD=4.8, range=14 - 27)
2011	76	75.5	46.1	100	17.8 (SD=3.9, range=12 - 27)
2012	35	74.5	34.3	100	18.9 (SD=4.3, range=11 - 27)
2013	138	72.7	60.1	100	16.9 (SD=4.0, range=7 - 27)
2014	116	73.7	56.0	100	17.1 (SD=4.8 , range=7 - 28)
2015	80	73.6	47.5	100	18.3 (SD=5.2, range=8 - 28)
2016	202	72.9	42.6	99.5	18.6 (SD=4.7, range=10 - 29)
2017	132	74	30.3	100	19.8 (SD=4.9, range=11 - 27)
2018	127	62.9	29.9	100	19.9 (SD=4.5, range=12 - 27)
Overall	913	72.1	43.6	99.9	18.5 (SD=4.7, range=7 - 29)

*Enrolled as % definitely eligible

All sites

- Overall, of the 1931 clients who were referred to the programme prior to end of December 2018, 913 clients were enrolled (had valid completed visit) on the programme prior to the end of December 2018.
- 43.6% of clients were enrolled by the end of the 16th week gestation and 99.9% of clients were enrolled by the end of the 28th week gestation.
- The mean gestational age (week's gestation) at enrolment was 18.5 and this ranged from 7 weeks to 29 weeks gestation.

Age at LMP

Table 4: Age at LMP overall and by site

Site	Clients	Age (years)		
	N	Average	Standard deviation	Range
BHSCT	183	17.1	1.5	12.8 – 20.5
NHSCT	110	17.0	1.3	13.7 – 20.0
SEHSCT	138	17.2	1.3	14.6 – 19.9
SHSCT	193	17.2	1.3	13.9 – 19.8
WHsCT	289	17.3	1.3	13.8 – 19.9
Overall	913	17.2	1.4	12.8 – 20.5

Table 5: Age at LMP – Average, SD and range by year of enrolment (All sites)

Year	Clients	Average	Standard Deviation	Range
2010	7	17.3	1.3	15.9 – 19.5
2011	76	17.5	1.3	14.6 – 19.9
2012	35	16.9	1.6	13.8 – 19.8
2013	138	17.2	1.4	13.4 – 19.8
2014	116	17.3	1.4	13.3 – 20.5
2015	80	17.1	1.5	12.8 – 20.0
2016	202	17.1	1.3	14.1 – 20.0
2017	132	17.2	1.2	14.5 – 19.9
2018	127	17.1	1.4	13.7 – 19.8
Total	913	17.2	1.4	12.8 – 20.5

Table 6: Age at LMP – Average, SD and range by site and year of enrolment

Site	Year	Clients	Average	Standard Deviation	Range
BHSCT	2013	43	17.0	1.6	13.4 – 19.8
	2014	43	17.1	1.6	13.3 – 20.5
	2015	8	17.4	2.4	12.8 – 20.0
	2016	43	17.2	1.4	14.5 – 19.7
	2017	25	17.0	1.3	14.8 – 19.7
	2018	21	16.8	1.2	14.6 – 19.0
	Total		183	17.1	1.5
NHSCT	2015	19	16.9	1.2	16.0 – 20.0
	2016	41	17.0	1.2	15.6 – 20.0
	2017	28	17.3	1.3	14.5 – 19.9
	2018	22	16.6	1.8	13.7 – 19.7
	Total		110	17.0	1.3
SEHSCT	2014	1	17.0	-	-
	2015	21	17.3	1.3	15.2 – 19.8
	2016	58	17.1	1.3	15.2 – 19.9
	2017	18	17.1	1.1	14.5 – 19.9
	2018	40	17.2	1.4	13.7 – 19.7
	Total		138	17.2	1.3
SHSCT	2012	13	16.4	1.4	14.7 – 19.0
	2013	72	17.3	1.3	13.9 – 19.8
	2014	23	17.5	1.1	15.3 – 19.8
	2015	5	17.2	0.8	16.6 – 18.4
	2016	32	17.0	1.5	14.1 – 19.7
	2017	20	16.9	1.3	15.0 – 18.9
	2018	28	17.2	1.1	15.7 – 19.8
	Total		193	17.2	1.3
WHSCT	2010	7	17.3	1.3	15.9 – 19.5
	2011	76	17.5	1.3	14.6 – 19.9
	2012	22	17.2	1.7	13.8 – 19.8
	2013	23	17.2	1.2	14.1 – 19.8
	2014	49	17.5	1.3	14.2 – 19.8

	2015	27	16.9	1.5	14.1 – 19.6
	2016	28	17.2	1.3	15.0 – 19.8
	2017	41	17.4	1.2	15.2 – 19.7
	2018	16	17.6	1.3	15.9 – 19.7
	Total	289	17.3	1.3	13.8 – 19.9
All site total		913	17.2	1.4	12.8 – 20.5

All sites

- Overall, for all sites, the average age for all clients enrolled to the end of 2018 (N=913) at LMP was 17.2 (SD= 1.4, range 12.8 to 20.5).
- For BHSCT, the average age for all clients enrolled to the end of 2018 (N= 183) at LMP was 17.1 (SD= 1.5, range 12.8 to 20.5).
- For NHSCT, the average age for all clients enrolled to the end of 2018 (N= 110) at LMP was 17.0 (SD= 1.3, range 13.7 to 20.0).
- For SEHSCT, the average age for all clients enrolled to the end of 2018 (N= 138) at LMP was 17.2 (SD= 1.3, range 14.6 to 19.9).
- For SHSCT, the average age for all clients enrolled to the end of 2018 (N= 193) at LMP was 17.2 (SD=1.3, range 13.9 to 19.8).
- For WHSCT, the average age for all clients enrolled to the end of 2018 (N= 289) at LMP was 17.3 (SD= 1.3, range 13.8 to 19.9).

Table 7: Aged 19 or under at LMP – overall by year of enrolment (All sites)

Year	Yes (N)	No (N)	Total (N)	Age 19 or under at LMP (%)
2010	7	-	7	100
2011	76	-	76	100
2012	35	-	35	100
2013	138	-	138	100
2014	115	1	116	99.1
2015	79	1	80	98.8
2016	202	-	202	100
2017	132	-	132	100
2018	127	-	127	100
Total	911	2	913	99.8

*Two clients were aged 20 at LMP.

Table 8: Aged 19 or under at LMP by site and year of enrolment

Site	Year	Yes (N)	No (N)	Total (N)	Age 19 or under at LMP (%)	Age 20 or over at LMP (%)
BHSCT	2013	43	-	43	100.0	0.0
	2014	42	1	43	97.7	2.3
	2015	8	-	8	100.0	0.0
	2016	43	-	43	100.0	0.0
	2017	25	-	25	100.0	0.0
	2018	21	-	21	100.0	0.0
	Total		182	1	183	99.5
NHSCT	2015	18	1	19	94.7	5.3
	2016	41	-	41	100.0	0.0
	2017	28	-	28	100.0	0.0
	2018	22	-	22	100.0	0.0
	Total	109	1	110	99.1	0.9
SEHSCT	2014	1	-	1	100.0	0.0
	2015	21	-	21	100.0	0.0
	2016	58	-	58	100.0	0.0
	2017	18	-	18	100.0	0.0
	2018	40	-	40	100.0	0.0
	Total	138	-	138	100.0	0.0
SHSCT	2012	13	-	13	100.0	0.0
	2013	72	-	72	100.0	0.0
	2014	23	-	23	100.0	0.0
	2015	5	-	5	100.0	0.0
	2016	32	-	32	100.0	0.0
	2017	20	-	20	100.0	0.0
	2018	28	-	28	100.0	0.0
	Total	193	-	193	100.0	0.0
WHSCT	2010	7	-	7	100.0	0.0
	2011	76	-	76	100.0	0.0
	2012	22	-	22	100.0	0.0
	2013	23	-	23	100.0	0.0

	2014	49	-	49	100.0	0.0
	2015	27	-	27	100.0	0.0
	2016	28	-	28	100.0	0.0
	2017	41	-	41	100.0	0.0
	2018	16	-	16	100.0	0.0
	Total	289	-	289	100.0	0.0
All site total		911	2	913	99.8	0.2

*Two clients were aged 20 at LMP.

- Overall, across all sites 99.8% (N=911) of clients were aged 19 or under at LMP. Two clients (0.2%) were aged 20 at LMP.
- In BHSCT 99.5% (N=182) of clients were aged 19 or under at LMP. One client (0.5%) was aged 20 at LMP.
- In NHSCT 99.1% (N=109) clients were aged 19 or under at LMP. One client (0.9%) was aged 20 at LMP.
- In SEHSCT 100% (N=138) clients were aged 19 or under at LMP.
- In SHSCT 100% (N=193) clients were aged 19 or under at LMP.
- In WHSCT 100% (N=289) clients were aged 19 or under at LMP.

Gestation at enrolment

Table 9: Gestation at enrolment (weeks) – Average, SD and range for all sites by year of enrolment

Year Enrolled	Clients	Average	Standard Deviation	Range
2010	7	24.6	4.8	14 – 27
2011	76	17.8	3.9	12 – 27
2012	35	18.9	4.3	11 – 27
2013	138	16.9	4.0	7 – 27
2014	116	17.1	4.8	7 – 28
2015	80	18.3	5.2	8 – 28
2016	202	18.6	4.7	10 – 29
2017	132	19.8	4.9	11 – 27
2018	127	19.9	4.5	12 – 27
All site total	913	18.5	4.7	7 – 29

Table 10: Gestation at enrolment (weeks) – Average, SD and range by site and year of enrolment

Site	Year Enrolled	Clients	Average	Standard Deviation	Range
BHSCT	2013	43	17.1	4.6	8 – 27
	2014	43	15.8	4.9	7 – 27
	2015	8	20.5	5.7	11 – 27
	2016	43	19.9	4.9	12 – 27
	2017	25	22.0	4.4	13 – 27
	2018	21	22.4	3.8	15 – 27
	Total	183	18.9	5.2	7 – 27
NHSCT	2015	19	21.4	5.9	12 – 28
	2016	41	21.4	4.1	11 – 29
	2017	28	24.5	3.6	12 – 27
	2018	22	24.4	2.9	18 – 27
	Total	110	22.8	4.4	11 – 29
SEHSCT	2014	1	12.0	-	-
	2015	21	16.6	4.6	8 – 26

	2016	58	16.8	4.1	10 – 28
	2017	18	16.3	3.7	11 – 25
	2018	40	17.1	3.3	12 – 26
	Total	138	16.7	3.9	8 – 28
SHSCT	2012	13	19.1	4.9	11 – 27
	2013	72	16.8	3.9	7 – 27
	2014	23	18.5	4.3	12 – 28
	2015	5	19.6	5.3	14 – 27
	2016	32	19.7	4.8	12 – 27
	2017	20	20.6	3.5	15 – 26
	2018	28	19.7	3.9	13 – 26
	Total	193	18.5	4.3	7 – 28
WHSCT	2010	7	24.6	4.8	14 – 27
	2011	76	17.8	3.9	12 – 27
	2012	22	18.8	4.1	13 – 26
	2013	23	16.7	3.1	13 – 26
	2014	49	17.8	4.6	11 – 27
	2015	27	16.4	3.6	10 – 24
	2016	28	15.4	3.1	11 – 23
	2017	41	16.5	3.5	11 – 27
	2018	16	17.8	4.3	13 – 27
	Total	289	17.4	4.1	10 – 27
All Site Total		913	18.5	4.7	7 – 29

All sites

- The mean gestational age (weeks gestation) at enrolment was 18.5 and this ranged from 7 – 29 weeks gestation.
- In the BHSCT, the mean gestational age (weeks gestation) at enrolment was 18.9 and this ranged from 7 – 27 weeks gestation.
- In the NHSCT, the mean gestational age (weeks gestation) at enrolment was 22.8 and this ranged from 11 – 29 weeks gestation.
- In the SEHSCT, the mean gestational age (weeks gestation) at enrolment was 16.7 and this ranged from 8 – 28 weeks gestation.
- In the SHSCT, the mean gestational age (weeks gestation) at enrolment was 18.5 and this ranged from 7 – 28 weeks gestation.
- In the WHSCT, the mean gestational age (weeks gestation) at enrolment was 17.4 and this ranged from 10 – 27 weeks gestation.

Enrolled by the end of 16 weeks gestation

Table 11: Percentage of clients (%) enrolled by end of 16 weeks gestation by year of enrolment (All sites)

Year	Yes	No	Total	Enrolled by end of 16th week (%)
2010	1	6	7	14.3
2011	35	41	76	46.1
2012	12	23	35	34.3
2013	83	55	138	60.1
2014	65	51	116	56.0
2015	38	42	80	47.5
2016	86	116	202	42.6
2017	40	92	132	30.3
2018	38	89	127	29.9
Total	398	515	913	43.6

Table 12: Percentage of clients (%) enrolled by end of 16 weeks gestation by site and year of enrolment

Site	Year	Yes	No	Total	Enrolled by end of 16th week (%)
BHSCT	2013	26	17	43	60.5
	2014	31	12	43	72.1
	2015	2	6	8	25.0
	2016	12	31	43	27.9
	2017	3	22	25	12.0
	2018	2	19	21	9.5
	Total	76	107	183	41.5
NHSCT	2015	5	14	19	26.3
	2016	5	36	41	12.2
	2017	1	27	28	3.6
	2018	-	22	22	0.0
Total	11	99	110	10.0	
SEHSCT	2014	1	-	1	100.0

	2015	14	7	21	66.7
	2016	34	24	58	58.6
	2017	10	8	18	55.6
	2018	21	19	40	52.5
	Total	80	58	138	58.0
SHSCT	2012	4	9	13	30.8
	2013	41	31	72	56.9
	2014	7	16	23	30.4
	2015	2	3	5	40.0
	2016	13	19	32	40.6
	2017	3	17	20	15.0
	2018	7	21	28	25.0
	Total	77	116	193	39.9
WHSCT	2010	1	6	7	14.3
	2011	35	41	76	46.1
	2012	8	14	22	36.4
	2013	16	7	23	69.6
	2014	26	23	49	53.1
	2015	15	12	27	55.6
	2016	22	6	28	78.6
	2017	23	18	41	56.1
	2018	8	8	16	50.0
	Total	154	135	289	53.3
All site total		398	515	913	43.6

All sites

- Overall, 43.6% (N=398) of clients were enrolled by the end of 16 weeks gestation. The fidelity goal for recruitment by 16 weeks is 60%.
- 41.5% (N=76) of clients in the BHSCT were enrolled by the end of 16 weeks gestation.
- 10% (N=11) of clients in the NHSCT were enrolled by the end of 16 weeks gestation.
- 58% (N=80) of clients in the SEHSCT were enrolled by the end of 16 weeks gestation.
- 39.9% (N=77) of clients in the SHSCT were enrolled by the end of 16 weeks gestation.
- 53.3% (N=154) of clients in the WHSCT were enrolled by the end of 16 weeks gestation.

Enrolled by the end of 28 weeks gestation

Table 13: Percentage of clients (%) enrolled by the end of 28 weeks gestation by year of enrolment (All sites)

Year	Yes	No	Total	Enrolled by end 28 th week gestation
2010	7	-	7	100
2011	76	-	76	100
2012	35	-	35	100
2013	138	-	138	100
2014	116	-	116	100
2015	80	-	80	100
2016	201	1	202	99.5
2017	132	-	132	100
2018	127	-	127	100
Total	912	1	913	99.9

Table 14: Percentage of clients (%) enrolled by the end of 28 weeks gestation by site and year of enrolment

Site	Year	Yes	No	Grand Total	Enrolled by end 28th week gestation
BHSCT	2013	43	-	43	100
	2014	43	-	43	100
	2015	8	-	8	100
	2016	43	-	43	100
	2017	25	-	25	100
	2018	21	-	21	100
	Total	183	-	183	100
NHSCT	2015	19	-	19	100
	2016	40	1	41	97.6
	2017	28	-	28	100
	2018	22	-	22	100
	Total	109	1	110	99.1
SEHSCT	2014	1	-	1	100

	2015	21	-	21	100
	2016	58	-	58	100
	2017	18	-	18	100
	2018	40	-	40	100
	Total	138	-	138	100
SHSCT	2012	13	-	13	100
	2013	72	-	72	100
	2014	23	-	23	100
	2015	5	-	5	100
	2016	32	-	32	100
	2017	20	-	20	100
	2018	28	-	28	100
	Total	193	-	193	100
WHSCT	2010	7	-	7	100
	2011	76	-	76	100
	2012	22	-	22	100
	2013	23	-	23	100
	2014	49	-	49	100
	2015	27	-	27	100
	2016	28	-	28	100
	2017	41	-	41	100
	2018	16	-	16	100
	Total	289	-	289	100
All Site total		912	1	913	99.9

All sites

- Overall, 99.9% (N=912) of clients were enrolled by the end of 28 weeks gestation. The fidelity goal for recruitment by 28 weeks is 100%.
- 100% (N=183) of clients in the BHSCT were enrolled by the end of 28 weeks gestation.
- 99.1% (N=109) of clients in the NHSCT were enrolled by the end of 28 weeks gestation.
- 100% (N=138) of clients in the SEHSCT were enrolled by the end of 28 weeks gestation.
- 100% (N=193) of clients in the SHSCT were enrolled by the end of 28 weeks gestation.
- 100% (N=289) of clients in the WHSCT were enrolled by the end of 28 weeks gestation.

Annual Income

Table 15: Annual Income – All sites

Annual Income	BHSCT		NHSCT		SEHSCT		SHSCT		WHSCT		All Sites	
	N	%	N	%	N	%	N	%	N	%	N	%
Total clients enrolled	183	-	110	-	138	-	193	-	289	-	913	-
Total clients form scheduled	183	-	109	-	138	-	193	-	289	-	912	-
Clients with income data recorded	171	93.4	99	90.8	135	97.8	177	91.7	275	95.2	857	94.0
Less than £1,600 per year	54	31.4	13	13.1	24	17.8	67	38.1	97	35.3	255	29.8
£1,600 - £3,099 per year	35	20.3	17	17.2	29	21.5	31	17.6	75	27.3	187	21.8
£3,100 - £4,699 per year	22	12.8	10	10.1	8	5.9	26	14.8	21	7.6	87	10.2
£4,700 - £6,199 per year	7	4.1	5	5.1	6	4.4	11	6.3	20	7.3	49	5.7
£6,200 - £7,799 per year	10	5.8	4	4.0	6	4.4	3	1.7	11	4.0	34	4.0
£7,800 - £10,399 per year	9	5.2	4	4.0	5	3.7	4	2.3	9	3.3	31	3.6
£10,400 - £12,999 per year	7	4.1	5	5.1	4	3.0	12	6.8	13	4.7	41	4.8
£13,000 - £15,599 per year	3	1.7	4	4.0	4	3.0	3	1.7	5	1.8	19	2.2
£15,600 - £18,199 per year	1	0.6	1	1.0	2	1.5	1	0.6	1	0.4	6	0.7
£18,200 - £20,799 per year	3	1.7	-	0.0	1	0.7	2	1.1	1	0.4	7	0.8
£20,800 - £25,999 per year	-	0.0	2	2.0	-	0.0	3	1.7	3	1.1	8	0.9
£26,000 - £31,199 per year	1	0.6	-	0.0	-	0.0	-	0.0	-	0.0	1	0.1
£31,200 - £36,399 per year	-	0.0	-	0.0	2	1.5	-	0.0	-	0.0	2	0.2
£36,400 or more per year	-	0.0	-	0.0	1	0.7	-	0.0	-	0.0	1	0.1
Client unable/unwilling to give information on income	19	11.0	34	34.3	43	31.9	14	8.0	19	6.9	129	15.1
<£13,000 per year	144	84.2	58	58.6	82	60.7	154	87.0	246	89.5	684	79.8
Income Source												

Income based on household	18	10.5	11	11.1	13	9.6	17	9.6	25	9.1	84	9.8
Income based on self only	71	41.5	56	56.6	79	58.5	65	36.7	87	31.6	358	41.8
Income based on household or self only is missing	82	48.0	32	32.2	43	31.9	95	53.7	163	59.3	415	48.4
All income from benefits	67	39.2	45	45.5	56	41.5	36	20.3	106	38.5	310	36.2

All sites

- Annual income was recorded for 94% (N= 857) clients at intake.
- 79.8% (N=684) had an annual income of less than £13,000. However, given the level of completeness (i.e. 48.4% (N=415) missing data) in relation to the source of income as household or self only it is not possible to make comparisons against currently available poverty indicators.
- Relative poverty can be defined as living in a household with an income below 60% of the average income. Annual income in N.I, before housing costs, was £23,504 per year in 2016/2017 and 60% of this is £14,102. Approximately 18% of the population in Northern Ireland were living in relative poverty before housing costs in 2016/2017 (Northern Ireland Poverty Bulletin 2016/2017 [NISRA, 2018]). We have presented comparison data for clients with income under £13,000 due to this most closely matching the £14,102 poverty cut off.
- It is not possible to compare the FNP income data against currently available poverty indicators for the WHSCT, the SHSCT and the BHSCT due to the high level of missing data (between 48% and 60% of data was missing) on whether annual income is based on household income or based on self-only.
- The SEHSCT and the NHSCT had a lower level of missing data on whether annual income was based on household or self only (32.2%; 31.9% missing). However, we are not able to make comparisons with NI comparator data due to the fact that the majority of the clients reported on income based on 'self only' and the NI comparator data is based on household income.
- It is therefore necessary to consider if the income question should be adjusted to focus on household income only. It is also necessary to consider the reasons and possible solutions for such a high level of missing data on this question.

BHSCT

- Annual income was recorded for 93.4% (N=171) clients at intake.
- 84.2% (N=144) had an annual income of less than £13,000. However, given the level of completeness (i.e. 48 % missing data) in relation to the source of income as household or self only it is not possible to make comparisons against currently available poverty indicators.

NHSCT

- Annual income was recorded for 90.8% (N=99) clients at intake.
- 58.6% (N=58) had an annual income of less than £13,000. We are not able to make comparisons with NI comparator data due to the fact that the majority of the clients reported on income based on 'self only' and the NI comparator data is based on household income.

SEHSCT

- Annual income was recorded for 97.8% (N=135) clients at intake.
- 60.7% (N=82) had an annual income of less than £13,000. We are not able to make comparisons with NI comparator data due to the fact that the majority of the clients reported on income based on 'self only' and the NI comparator data is based on household income.

SHSCT

- Annual income was recorded for 91.7% (N=177) clients at intake.
- 87.0% (N=154) had an annual income of less than £13,000. However, given the level of completeness (i.e. 53.7% missing data) in relation to the source of income as household or self only it is not possible to make comparisons against currently available poverty indicators.

WHSCT

- Annual income was recorded for 95.2% (N=275) clients at intake.
- 89.5% (N=246) had an annual income of less than £13,000. However, given the level of completeness (i.e. 59.3% missing data) in relation to the source of income as household or self only it is not possible to make comparisons against currently available poverty indicators.

Ethnicity

Table 16: Ethnicity – All sites

		BHSCT	NHSCT	SEHSCT	SHSCT	WHSCT	All Sites
Total clients enrolled	N	183	110	138	193	289	913
Clients who were scheduled to have reported their ethnicity	N	183	109	138	193	289	912
Clients who have reported their ethnicity	N	171	99	135	177	275	857
	%	93.4	90.8	97.8	91.7	95.2	94.0
White British	N	87	72	84	73	78	394
	%	50.9	72.7	62.2	41.2	28.4	46.0
White Irish	N	73	22	46	70	195	406
	%	42.7	22.2	34.1	39.5	70.9	47.4
White, Other	N	6	2	2	21	1	32
	%	3.5	2.0	1.5	11.9	0.4	3.7
Other ethnic group*	N	5	3	3	13	1	25
	%	2.9	3.0	2.2	7.3	0.4	2.9

*This includes Black or Black British, Other, Mixed, Asian or Asian British and other ethnic group which have all been grouped together due to low numbers and disclosure issues.

All sites

- Overall, ethnicity was recorded for 94% (N=857) of the 912 who were scheduled to have their ethnicity recorded by the report end date (31.12.18).
- For the 857 clients who had their ethnicity recorded, 46% (N= 394) reported that they were White British, 47.4% (N=406) reported that they were White Irish, 3.7% (N=32) reported that they were White Other, 2.9% (N=25) reported that they were Other ethnic group.

By Site

- In BHSCT, 50.9% of clients (N= 87) reported that they were White British, 42.7% (N=73) reported that they were White Irish, 3.5% (N= 6) reported that they were White Other, 2.9% (N= 5) reported that they were Other ethnic group.
- In NHSCT, 72.7% of clients (N= 72) reported that they were White British, 22.2% (N=22) reported that they were White Irish, 2% (N= 2) reported that they were White Other, and 3% (N=3) reported that they were Other ethnic group.

- In SEHSCT, 62.2% of clients (N=84) reported that they were White British, 34.1% (N=46) reported that they were White Irish, 1.5% (N=2) reported that they were White Other, 2.2% (N=3) reported that they were Other ethnic group.
- In SHSCT, 41.2% of clients (N=73) reported that they were White British, 39.5% (N=70) reported that they were White Irish, 11.9% (N=21) reported that they were White Other, 7.3% (N=13) reported that they were Other ethnic group.
- In WHSCT, 28.4% of clients (N=78) reported that they were White British, 70.9% (N=195) reported that they were White Irish, 0.4% (N=1) reported that they were White Other and 0.4% (N= 1) reported that they were Other ethnic group.

Education, Employment and Training (EET)

Intake

Table 17: Education, Employment and Training (EET) Intake – All sites

		BHSCT	NHSCT	SEHSCT	SHSCT	WHSCT	All Sites
Total clients enrolled	N	183	110	138	193	289	913
Clients who were scheduled to have reported their ETT status at intake	N	183	109	138	193	289	912
Clients who have reported their ETT status at intake	N	171	99	135	177	275	857
	%	93.4	90.8	97.8	91.7	95.2	94.0
Clients in Education at intake	N	76	47	57	80	139	399
	%	44.4	47.5	42.2	45.2	50.5	46.6
Clients in employment (currently) at intake	N	29	17	36	31	55	168
	%	17.0	17.2	26.7	17.5	20.0	19.6
Clients in both education and employment	N	12	9	13	17	31	82
	%	7.0	9.1	9.6	9.6	11.3	9.6
Clients in education or employment (EET)	N	93	55	80	94	163	485
	%	54.4	55.6	59.3	53.1	59.3	56.6
Clients not in education or employment (NEET)	N	78	44	55	83	112	372
	%	45.6	44.4	40.7	46.9	40.7	43.4

All sites

- Education and employment status was recorded for 94% (N=857) of clients at intake. At intake 46.6% (N=399) clients were in education, 19.6% (N=168) of clients were in employment and 9.6% (N=82) of clients were in both education and employment.
- At intake, 56.6% (N=485) clients were either in education or employment (EET) and 43.4% (N=372) clients were not in education or employment (NEET).

BHSCT

- Education and employment status was recorded for 93.4% (N=171) of clients at intake. At intake 44.4% (N=76) clients were in education, 17% (N=29) of clients were in employment and 7% (N=12) of clients were in both education and employment.
- At intake, 54.4% (N=93) clients were either in education or employment (EET) and 45.6% (N=78) clients were not in education or employment (NEET).

NHSCT

- Education and employment status was recorded for 90.8% (N=99) of clients at intake. At intake 47.5% (N=47) clients were in education, 17.2% (N=17) of clients were in employment and 9.1% (N=9) of clients were in both education and employment.
- At intake, 55.6% (N=55) clients were either in education or employment (EET) and 44.4% (N=44) clients were not in education or employment (NEET).

SEHSCT

- Education and employment status was recorded for 97.8% (N=135) of clients at intake. At intake 42.2% (N=57) clients were in education, 26.7% (N=36) of clients were in employment and 9.6% (N=13) of clients were in both education and employment.
- At intake, 59.3% (N=80) clients were either in education or employment (EET) and 40.7% (N=55) clients were not in education or employment (NEET).

SHSCT

- Education and employment status was recorded for 91.7% (N=177) of clients at intake. At intake 45.2% (N=80) clients were in education, 17.5% (N=31) of clients were in employment and 9.6% (N=17) of clients were in both education and employment.
- At intake, 53.1% (N=94) clients were either in education or employment (EET) and 46.9% (N=83) clients were not in education or employment (NEET).

WHSCT

- Education and employment status was recorded for 95.2% (N=275) of clients at intake. At intake 50.5% (N=139) clients were in education, 20% (N=55) of clients were in employment and 11.3% (N=31) of clients were in both education and employment.
- At intake, 59.3% (N=163) clients were either in education or employment (EET) and 40.7% (N=112) clients were not in education or employment (NEET).

24 months

Table 18: Education, Employment and Training (EET) 24 Months – All sites

		BHSCT	NHSCT	SEHSCT	SHSCT	WHSCT	All Sites
Clients active at 24 months	N	102	41	54	126	186	509
Clients who have reported their ETT status at 24 months	N	91	31	53	98	157	430
	%	89.2	75.6	98.1	77.8	84.4	84.5
Clients in Education at 24 months	N	27	9	13	31	53	133
	%	29.7	29.0	24.5	31.6	33.8	30.9
Clients in employment (currently) at 24 months	N	24	4	14	17	32	91
	%	26.4	12.9	26.4	17.3	20.4	21.2
Clients in both education and employment	N	10	2	4	4	14	34
	%	11.0	6.5	7.5	4.1	8.9	7.9
Clients in education or employment (EET)	N	41	11	23	44	71	190
	%	45.1	35.5	43.4	44.9	45.2	44.2
Clients not in education or employment (NEET)	N	50	20	30	54	86	240
	%	54.9	64.5	56.6	55.1	54.8	55.8

*There are fewer active clients at 24 months in NHSCT and SEHSCT as these sites commenced in 2015.

All sites

- Education and employment status was recorded for 84.5% (N=430) of clients at 24 months. At 24 months 30.9% (N=133) clients were in education, 21.2% (N=91) of clients were in employment and 7.9% (N=34) of clients were in both education and employment.
- At 24 months, 44.2% (N=190) clients were either in education or employment (EET) and 55.8% (N=240) clients were not in education or employment (NEET).

BHSCT

- Education and employment status was recorded for 89.2% (N=91) of clients at 24 months. At 24 months, 29.7% (N=27) clients were in education, 26.4% (N=24) of clients were in employment and 11% (N=10) of clients were in both education and employment.
- At 24 months, 45.1% (N=41) clients were either in education or employment (EET) and 54.9% (N=50) clients were not in education or employment (NEET).

NHSCT

- Education and employment status was recorded for 75.6% (N=31) of clients at 24 months. At 24 months, 29% (N=9) clients were in education, 12.9% (N=4) clients were in employment and 6.5% (N=2) clients were in both education and employment.
- At 24 months, 35.5% (N=11) clients were either in education or employment (EET) and 64.5% (N=20) clients were not in education or employment (NEET).

SEHSCT

- Education and employment status was recorded for 98.1% (N= 53) of clients at 24 months. At 24 months, 24.5% (N=13) clients were in education, 26.4% (N=14) of clients were in employment and 7.5% (N=4) of clients were in both education and employment.
- At 24 months, 43.4% (N=23) clients were either in education or employment (EET) and 56.6% (N=30) clients were not in education or employment (NEET).

SHSCT

- Education and employment status was recorded for 77.8% (N=98) of clients at 24 months. At 24 months 31.6% (N=31) clients were in education, 17.3% (N=17) of clients were in employment and 4.1% (N=4) of clients were in both education and employment.
- At 24 months, 44.9% (N=44) clients were either in education or employment (EET) and 55.1% (N=54) clients were not in education or employment (NEET).

WHSCT

- Education and employment status was recorded for 84.4% (N=157) of clients at 24 months. At 24 months, 33.8% (N=53) clients were in education, 20.4% (N=32) of clients were in employment and 8.9% (N=14) of clients were in both education and employment.
- At 24 months, 45.2% (N=71) clients were either in education or employment (EET) and 54.8% (N=86) clients were not in education or employment (NEET).

Programme Activity and Fidelity

Dosage

Pregnancy Dosage

Table 19: Dosage Pregnancy Stage – All sites by year enrolled

Year enrolled	Clients	Pregnancy Completer	Active Pregnancy Completer	Scheduled Visits	Actual Visits*	Scheduled Visits	Actual Visits	Expected visits Achieved Given Gestational age**	Expected visits Achieved using 14 expected Visits***
	N	N	N	N	N	Mean N	Mean N	Mean %	Mean %
2010	7	7	7	66	33	9.4	4.7	52.0	33.7
2011	76	76	76	965	695	12.7	9.1	72.0	65.3
2012	35	35	35	436	388	12.5	11.1	90.0	79.2
2013	138	138	135	1752	1491	13.0	11.0	85.1	78.9
2014	116	116	111	1408	1111	12.7	10.0	79.2	71.5
2015	80	80	78	944	801	12.1	10.3	85.3	73.4
2016	202	202	200	2368	1872	11.8	9.4	79.2	66.9
2017	132	132	129	1464	1079	11.3	8.4	73.8	59.7
2018	127	74	74	791	625	10.7	8.4	79.5	60.3
Total	913	860	845	10194	8095	12.1	9.6	79.5	68.4

*Actual visits refer to completed valid visits. **The number of expected (scheduled) visits during the pregnancy stage is calculated for each client using the gestational age at enrolment and the EDD (estimated date of delivery). The total number of expected visits is adjusted if the client delivers before any remaining scheduled pregnancy visits. *** There are a maximum of 14 visits scheduled during the pregnancy stage.

This table reports on pregnancy dosage overall (all sites) for active clients (N=845) who had completed the pregnancy stage by 31.12.2018.

All sites

- 845 active clients completed the pregnancy stage by the end of 31.12.2018.
- Overall, clients were scheduled to have an average of 12.1 visits and they completed an average of 9.6 valid visits in pregnancy.
- Overall to date for all clients who enrolled since 2010, 79.5% of expected visits (based on gestational age) and 68.4% of expected visits (based on 14 expected visits) were achieved by the end of 2018.
- For clients who enrolled in 2018, 79.5% of expected visits (based on gestational age) and 60.3% of expected visits (based on 14 expected visits) were achieved by the end of 2018.

Table 20: Dosage Pregnancy Stage – BHSCT by year enrolled

Year Enrolled	Clients	Pregnancy completers	Active pregnancy completers	Scheduled Visits	Actual Visits*	Scheduled Visits	Actual Visits	Expected visits Achieved Given Gestational age**	Expected visits Achieved using 14 expected visits***
	N	N	N	N	N	Mean N	Mean N	Mean %	Mean %
2013	43	43	41	519	417	12.7	10.2	80.8	72.6
2014	43	43	40	519	404	13.0	10.1	77.6	72.1
2015	8	8	8	89	75	11.1	9.4	82.6	67.0
2016	43	43	43	480	333	11.2	7.7	70.5	55.3
2017	25	25	25	257	178	10.3	7.1	68.8	50.9
2018	21	17	17	161	129	9.5	7.6	81.3	54.2
BHSCT	183	179	174	2025	1536	11.6	8.8	75.9	63.1

*Actual visits refer to completed valid visits. **The number of expected (scheduled) visits during the pregnancy stage is calculated for each client using the gestational age at enrolment and the EDD (estimated date of delivery). The total number of expected visits is adjusted if the client delivers before any remaining scheduled pregnancy visits. *** There are a maximum of 14 visits scheduled during the pregnancy stage.

This table reports on pregnancy dosage (BHSCT) for active clients (N=174) who had completed the pregnancy stage by 31.12.2018.

BHSCT

- 174 active clients in the BHSCT completed the pregnancy stage by the end of 31.12.2018.
- Overall, clients in BHSCT were scheduled to have an average of 11.6 visits and they completed an average of 8.8 valid visits in pregnancy.
- Overall to date for all clients who enrolled since 2013, 75.9% of expected visits (based on gestational age) and 63.1 % of expected visits (based on 14 expected visits) were achieved by the end of 2018 in the BHSCT.
- For clients who enrolled in 2018, 81.3% of expected visits (based on gestational age) and 54.2% of expected visits (based on 14 expected visits) were achieved by the end of 2018.

Table 21: Dosage Pregnancy Stage – NHSCT by year enrolled

Year Enrolled	Clients	Pregnancy completers	Active pregnancy completers	Scheduled Visits	Actual Visits*	Scheduled Visits	Actual Visits	Expected visits Achieved Given Gestational age**	Expected visits Achieved using 14 expected visits***
	N	N	N	N	N	Mean N	Mean N	Mean %	Mean %
2015	19	19	19	204	179	10.7	9.4	90.4	67.3
2016	41	41	41	440	333	10.7	8.1	76.4	58.0
2017	28	28	27	247	188	9.1	7.0	75.6	49.7
2018	22	13	13	105	87	8.1	6.7	83.8	47.8
NHSCT	110	101	100	996	787	10.0	7.9	79.8	56.2

*Actual visits refer to completed valid visits. **The number of expected (scheduled) visits during the pregnancy stage is calculated for each client using the gestational age at enrolment and the EDD (estimated date of delivery). The total number of expected visits is adjusted if the client delivers before any remaining scheduled pregnancy visits. *** There are a maximum of 14 visits scheduled during the pregnancy stage.

This table reports on pregnancy dosage (NHSCT) for active clients (N= 100) who had completed the pregnancy stage by 31.12.2018.

NHSCT

- 100 active clients in the NHSCT completed the pregnancy stage by the end of 31.12.2018.
- Overall, clients in NHSCT were scheduled to have an average of 10 visits and they completed an average of 7.9 valid visits in pregnancy.
- Overall to date for all clients who enrolled since 2015, 79.8% of expected visits (based on gestational age) and 56.2% of expected visits (based on 14 expected visits) were achieved by the end of 2018 in the NHSCT.
- For clients who enrolled in 2018, 83.8% of expected visits (based on gestational age) and 47.8% of expected visits (based on 14 expected visits) were achieved by the end of 2018.

Table 22: Dosage Pregnancy Stage – SEHSCT by year enrolled

Year Enrolled	Clients	Pregnancy completers	Active pregnancy completers	Scheduled Visits	Actual Visits*	Scheduled Visits	Actual Visits	Expected visits Achieved Given Gestational age**	Expected visits Achieved using 14 expected visits***
	N	N	N	N	N	Mean N	Mean N	Mean %	Mean %
2014	1	1	1	14	8	14.0	8.0	57.1	57.1
2015	21	21	20	257	258	12.9	12.9	101.2	92.1
2016	58	58	57	730	665	12.8	11.7	91.2	83.3
2017	18	18	18	230	190	12.8	10.6	84.3	75.4
2018	40	16	16	197	169	12.3	10.6	85.0	75.4
SEHSCT	138	114	112	1428	1290	12.8	11.5	90.7	82.3

*Actual visits refer to completed valid visits. **The number of expected (scheduled) visits during the pregnancy stage is calculated for each client using the gestational age at enrolment and the EDD (estimated date of delivery). The total number of expected visits is adjusted if the client delivers before any remaining scheduled pregnancy visits. *** There are a maximum of 14 visits scheduled during the pregnancy stage.

This table reports on pregnancy dosage (SEHSCT) for active clients (N= 112) who had completed the pregnancy stage by 31.12.2018.

SEHSCT

- 112 active clients in the SEHSCT completed the pregnancy stage by the end of 31.12.2018.
- Overall, clients in SEHSCT were scheduled to have an average of 12.8 visits and they completed an average of 11.5 valid visits in pregnancy.
- Overall to date for all clients who enrolled since 2014, 90.7% of expected visits (based on gestational age) and 82.3% of expected visits (based on 14 expected visits) were achieved by the end of 2018 in the SEHSCT.
- For clients who enrolled in 2018, 85% of expected visits (based on gestational age) and 75.4% of expected visits (based on 14 expected visits) were achieved by the end of 2018.

Table 23: Dosage Pregnancy Stage – SHSCT by year enrolled

Year Enrolled	Clients	Pregnancy completers	Active pregnancy completers	Scheduled Visits	Actual Visits*	Scheduled Visits	Actual Visits	Expected visits Achieved Given Gestational age**	Expected visits Achieved using 14 expected visits***
	N	N	N	N	N	Mean N	Mean N	Mean %	Mean %
2012	13	13	13	161	153	12.4	11.8	95.6	84.1
2013	72	72	71	923	831	13.0	11.7	90.1	83.6
2014	23	23	23	285	240	12.4	10.4	85.1	74.5
2015	5	5	4	43	27	10.8	6.8	57.6	48.2
2016	32	32	31	353	302	11.4	9.7	84.7	69.6
2017	20	20	20	221	159	11.1	8.0	72.6	56.8
2018	28	20	20	238	187	11.9	9.4	78.7	66.8
SHSCT	193	185	182	2224	1899	12.2	10.4	85.0	74.5

*Actual visits refer to completed valid visits. **The number of expected (scheduled) visits during the pregnancy stage is calculated for each client using the gestational age at enrolment and the EDD (estimated date of delivery). The total number of expected visits is adjusted if the client delivers before any remaining scheduled pregnancy visits. *** There are a maximum of 14 visits scheduled during the pregnancy stage.

This table reports on pregnancy dosage (SHSCT) for active clients (N=182) who had completed the pregnancy stage by 31.12.2018.

SHSCT

- 182 active clients in the SHSCT completed the pregnancy stage by the end of 31.12.2018.
- Overall, clients in SHSCT were scheduled to have an average of 12.2 visits and they completed an average of 10.4 valid visits in pregnancy.
- Overall to date for all clients who enrolled since 2012, 85% of expected visits (based on gestational age) and 74.5% of expected visits (based on 14 expected visits) were achieved by the end of 2018 in the SHSCT.
- For clients who enrolled in 2018, 78.7% of expected visits (based on gestational age) and 66.8% of expected visits (based on 14 expected visits) were achieved by the end of 2018.

Table 24: Dosage Pregnancy Stage – WHSCT by year enrolled

Year Enrolled	Clients	Pregnancy completers	Active pregnancy completers	Scheduled Visits	Actual Visits*	Scheduled Visits	Actual Visits	Expected visits Achieved Given Gestational age**	Expected visits Achieved using 14 expected visits***
	N	N	N	N	N	Mean N	Mean N	Mean %	Mean %
2010	7	7	7	66	33	9.4	4.7	52.0	33.7
2011	76	76	76	965	695	12.7	9.1	72.0	65.3
2012	22	22	22	275	235	12.5	10.7	86.6	76.3
2013	23	23	23	310	243	13.5	10.6	77.2	75.5
2014	49	49	47	590	459	12.6	9.8	78.1	69.8
2015	27	27	27	351	262	13.0	9.7	74.9	69.3
2016	28	28	28	365	239	13.0	8.5	65.9	61.0
2017	41	41	39	509	364	13.1	9.3	71.7	66.7
2018	16	8	8	90	53	11.3	6.6	59.9	47.3
WHSCT	289	281	277	3521	2583	12.7	9.3	73.4	66.6

*Actual visits refer to completed valid visits. **The number of expected (scheduled) visits during the pregnancy stage is calculated for each client using the gestational age at enrolment and the EDD (estimated date of delivery). The total number of expected visits is adjusted if the client delivers before any remaining scheduled pregnancy visits. *** There are a maximum of 14 visits scheduled during the pregnancy stage.

This table reports on pregnancy dosage (WHSCT) for active clients (N=277) who had completed the pregnancy stage by 31.12.2018.

WHSCT

- 277 active clients in the WHSCT completed the pregnancy stage by the end of 31.12.2018.
- Overall, clients in WHSCT were scheduled to have an average of 12.7 visits and they completed an average of 9.3 valid visits in pregnancy.
- Overall to date for all clients who enrolled since 2010, 73.4% of expected visits (based on gestational age) and 66.6% of expected visits (based on 14 expected visits) were achieved by the end of 2018 in the WHSCT.
- For clients who enrolled in 2018, 59.9% of expected visits (based on gestational age) and 47.3% of expected visits (based on 14 expected visits) were achieved by the end of 2018.

Infancy Dosage

Table 25: Dosage Infancy Stage – All sites by year enrolled

Year Enrolled	Clients Enrolled	Infancy Completer	Active Infancy Completer	Scheduled Visits	Actual visits	Scheduled Visits	Actual visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2010	7	7	5	140	92	28.0	18.4	65.7
2011	76	76	69	1932	1365	28.0	19.8	70.7
2012	35	35	32	896	758	28.0	23.7	84.6
2013	138	138	126	3528	2666	28.0	21.3	75.6
2014	116	116	106	2941	2105	27.7	19.9	71.6
2015	80	80	73	2044	1504	28.0	20.6	73.6
2016	202	202	174	4872	3327	28.0	19.1	68.3
2017	132	80	75	2100	1336	28.0	17.8	63.6
Total	786	734	660	18453	13153	28.0	20.0	71.3

This table reports on infancy dosage (all sites) for active clients (N= 660) who had completed the infancy stage by 31.12.2018.

All sites

- 660 active clients completed the infancy stage by the end of 31.12.2018.
- Overall, clients were scheduled to have an average of 28 visits and they completed an average of 20 valid visits in infancy.
- Overall to date for all clients who enrolled since 2010, 71.3% of expected visits were achieved by the end of 2018.
- For clients who enrolled in 2017, 63.6% of expected infancy visits were achieved by the end of 2018.

Table 26: Dosage Infancy Stage – BHSCT by year enrolled

Year Enrolled	Clients Enrolled	Infancy Completer	Active Infancy Completer	Scheduled Visits	Actual visits	Scheduled Visits	Actual visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2013	43	43	37	1036	734	28.0	20.4	70.8
2014	43	43	39	1073	760	27.5	19.5	70.8
2015	8	8	8	224	176	28.0	22.0	78.6
2016	43	43	37	1036	638	28.0	17.2	61.6
2017	25	11	11	308	161	28.0	14.6	52.3
BHSCT	162	148	132	3677	2469	27.9	18.8	67.1

This table reports on infancy dosage (BHSCT) for active clients (N=132) who had completed the infancy stage by 31.12.2018.

BHSCT

- 132 active clients in the BHSCT completed the infancy stage by the end of 31.12.2018.
- Overall, clients in BHSCT were scheduled to have an average of 27.9 visits and they completed an average of 18.8 valid visits in infancy.
- Overall to date for all clients who enrolled since 2013, 67.1% of expected visits were achieved by the end of 2018 in the BHSCT.
- For clients who enrolled in 2017, 52.3% of expected infancy visits were achieved by the end of 2018.

Table 27: Dosage Infancy Stage – NHSCT by year enrolled

Year Enrolled	Clients Enrolled	Infancy Completer	Active Infancy Completer	Scheduled Visits	Actual visits	Scheduled Visits	Actual visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2015	19	19	17	476	369	28.0	21.7	77.5
2016	41	41	34	952	680	28.0	20.0	71.4
2017	28	20	18	504	323	28.0	17.9	64.1
NHSCT	88	80	69	1932	1372	28.0	19.9	71.0

This table reports on infancy dosage (NHSCT) for active clients (N=69) who had completed the infancy stage by 31.12.2018.

NHSCT

- 69 active clients in the NHSCT completed the infancy stage by the end of 31.12.2018.
- Overall, clients in NHSCT were scheduled to have an average of 28 visits and they completed an average of 19.9 valid visits in infancy.
- Overall to date for all clients who enrolled since 2015, 71% of expected visits were achieved by the end of 2018 in the NHSCT.
- For clients who enrolled in 2017, 64.1% of expected infancy visits were achieved by the end of 2018.

Table 28: Dosage Infancy Stage – SEHSCT by year enrolled

Year Enrolled	Clients Enrolled	Infancy Completer	Active Infancy Completer	Scheduled Visits	Actual visits	Scheduled Visits	Actual visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2014	1	1	1	28	15	28	15.0	53.6
2015	21	21	19	532	411	28	21.6	77.3
2016	58	58	52	1456	1076	28	20.7	73.9
2017	18	15	15	420	294	28	19.6	70.0
SEHSCT	98	95	87	2436	1796	28	20.6	73.7

This table reports on infancy dosage (SEHSCT) for active clients (N=87) who had completed the infancy stage by 31.12.2018.

SEHSCT

- 87 active clients in the SEHSCT completed the infancy stage by the end of 31.12.2018.
- Overall, clients in SEHSCT were scheduled to have an average of 28 visits and they completed an average of 20.6 valid visits in infancy.
- Overall to date for all clients who enrolled since 2014, 73.7% of expected visits were achieved by the end of 2018 in the SEHSCT.
- For clients who enrolled in 2017, 70% of expected infancy visits were achieved by the end of 2018.

Table 29: Dosage Infancy Stage – SHSCT by year enrolled

Year Enrolled	Clients Enrolled	Infancy Completer	Active Infancy Completer	Scheduled Visits	Actual visits	Scheduled Visits	Actual visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2012	13	13	12	336	282	28	23.5	83.9
2013	72	72	68	1904	1497	28	22.0	78.6
2014	23	23	22	616	437	28	19.9	70.9
2015	5	5	4	112	96	28	24.0	85.7
2016	32	32	29	812	571	28	19.7	70.3
2017	20	12	12	336	203	28	16.9	60.4
SHSCT	165	157	147	4116	3086	28	21.0	75.0

This table reports on infancy dosage (SHSCT) for active clients (N=147) who had completed the infancy stage by 31.12.2018.

SHSCT

- 147 active clients in the SHSCT completed the infancy stage by the end of 31.12.2018.
- Overall, clients in SHSCT were scheduled to have an average of 28 visits and they completed an average of 21 valid visits in infancy.
- Overall to date for all clients who enrolled since 2012, 75% of expected visits were achieved by the end of 2018 in the SHSCT.
- For clients who enrolled in 2017, 60.4% of expected infancy visits were achieved by the end of 2018.

Table 30: Dosage Infancy Stage – WHSCT by year enrolled

Year Enrolled	Clients Enrolled	Infancy Completer	Active Infancy Completer	Scheduled Visits	Actual visits	Scheduled Visits	Actual visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2010	7	7	5	140	92	28.0	18.4	65.7
2011	76	76	69	1932	1365	28.0	19.8	70.7
2012	22	22	20	560	476	28.0	23.8	85.0
2013	23	23	21	588	435	28.0	20.7	74.0
2014	49	49	44	1224	893	27.8	20.3	73.2
2015	27	27	25	700	452	28.0	18.1	64.6
2016	28	28	22	616	362	28.0	16.5	58.8
2017	41	22	19	532	355	28.0	18.7	66.7
WHSCT	273	254	225	6292	4430	28.0	19.7	70.5

This table reports on infancy dosage (WHSCT) for active clients (N= 225) who had completed the infancy stage by 31.12.2018.

WHSCT

- 225 active clients in the WHSCT completed the infancy stage by the end of 31.12.2018.
- Overall, clients in WHSCT were scheduled to have an average of 28 visits and they completed an average of 19.7 valid visits in infancy.
- Overall to date for all clients who enrolled since 2010, 70.5% of expected visits were achieved by the end of 2018 in the WHSCT.
- For clients who enrolled in 2017, 66.7% of expected infancy visits were achieved by the end of 2018.

Toddlerhood Dosage

Table 31: Dosage Toddlerhood Stage – All sites by year enrolled

Year Enrolled	Clients Enrolled	Toddlerhood Completer	Active Completer	Scheduled Visits	Actual Visits	Scheduled Visits	Actual Visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2010	7	7	5	110	71	22.0	14.2	64.5
2011	76	76	64	1408	1059	22.0	16.5	75.2
2012	35	35	31	679	514	21.9	16.6	75.4
2013	138	138	117	2574	1611	22.0	13.8	62.6
2014	116	116	99	2173	1282	21.9	12.9	59.0
2015	80	80	68	1496	893	22.0	13.1	59.7
2016	202	136	104	2288	1404	22.0	13.5	61.4
Total	654	588	488	10728	6834	22.0	14.0	63.7

This table reports on toddlerhood dosage (all sites) for active clients (N=488) who had completed the toddlerhood stage by 31.12.2018.

All sites

- 488 active clients completed the toddlerhood stage by the end of 31.12.18
- Overall, clients were scheduled to have an average of 22 visits and they completed an average of 14 valid visits in toddlerhood.
- Overall to date for all clients who enrolled since 2010, 63.7% of expected visits were achieved by the end of 2018.
- For clients who enrolled in 2016, 61.4% of expected toddlerhood visits were achieved by the end of 2018.

Table 32: Dosage Toddlerhood Stage – BHSCT by year enrolled

Year Enrolled	Clients Enrolled	Toddlerhood Completer	Active Completer	Scheduled Visits	Actual Visits	Scheduled Visits	Actual Visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2013	43	43	35	770	509	22	14.5	66.1
2014	43	43	36	792	448	22	12.4	56.6
2015	8	8	7	154	116	22	16.6	75.3
2016	43	23	17	374	214	22	12.6	57.2
Total	137	117	95	2090	1287	22	13.5	61.6

This table reports on toddlerhood dosage (BHSCT) for active clients (N=95) who had completed the toddlerhood stage by 31.12.2018.

BHSCT

- 95 clients in the BHSCT completed the toddlerhood stage by the end of 31.12.2018.
- Overall, clients in BHSCT were scheduled to have an average of 22 visits and they completed an average of 13.5 valid visits in toddlerhood.
- Overall to date for all clients who enrolled since 2013, 61.6% of expected visits were achieved by the end of 2018 in the BHSCT.
- For clients who enrolled in 2016, 57.2% of expected toddlerhood visits were achieved by the end of 2018.

Table 33: Dosage Toddlerhood Stage – NHSCT by year enrolled

Year Enrolled	Clients Enrolled	Toddlerhood Completer	Active Completer	Scheduled Visits	Actual Visits	Scheduled Visits	Actual Visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2015	19	19	16	352	205	22	12.8	58.2
2016	41	33	22	484	275	22	12.5	56.8
Total	60	52	38	836	480	22	12.6	57.4

This table reports on toddlerhood dosage (NHSCT) for active clients (N=38) who had completed the toddlerhood stage by 31.12.2018.

NHSCT

- 38 clients in the NHSCT completed the toddlerhood stage by the end of 31.12.2018.
- Overall, clients in NHSCT were scheduled to have an average of 22 visits and they completed an average of 12.6 valid visits in toddlerhood.
- Overall to date for all clients who enrolled since 2015, 57.4% of expected visits were achieved by the end of 2018 in the NHSCT.
- For clients who enrolled in 2016, 56.8% of expected toddlerhood visits were achieved by the end of 2018.

Table 34: Dosage Toddlerhood Stage – SEHSCT by year enrolled

Year Enrolled	Clients Enrolled	Toddlerhood Completer	Active Completer	Scheduled Visits	Actual Visits	Scheduled Visits	Actual Visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2014	1	1	0	NA	NA	NA	NA	NA
2015	21	21	17	374	227	22	13.4	60.7
2016	58	41	34	748	499	22	14.7	66.7
Total	80	63	51	1122	726	22	14.2	64.7

This table reports on toddlerhood dosage (SEHSCT) for active clients (N=51) who had completed the toddlerhood stage by 31.12.2018.

SEHSCT

- 51 clients in the SEHSCT completed the toddlerhood stage by the end of 31.12.2018.
- Overall, clients in SEHSCT were scheduled to have an average of 22 visits and they completed an average of 14.2 valid visits in infancy.
- Overall to date for all clients who enrolled since 2014, 64.7% of expected visits were achieved by the end of 2018 in the SEHSCT.
- For clients who enrolled in 2016, 66.7% of expected toddlerhood visits were achieved by the end of 2018.

Table 35: Dosage Toddlerhood Stage – SHSCT by year enrolled

Year Enrolled	Clients Enrolled	Toddlerhood Completer	Active Completer	Scheduled Visits	Actual Visits	Scheduled Visits	Actual Visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2012	13	13	12	261	187	21.8	15.6	70.9
2013	72	72	62	1364	826	22.0	13.3	60.6
2014	23	23	21	457	274	21.8	13.0	60.1
2015	5	5	4	88	78	22.0	19.5	88.6
2016	32	28	23	506	350	22.0	15.2	69.2
Total	145	141	122	2676	1715	21.9	14.1	64.0

This table reports on toddlerhood dosage (SHSCT) for active clients (N=122) who had completed the toddlerhood stage by 31.12.2018.

SHSCT

- 122 clients in the SHSCT completed the toddlerhood stage by the end of 31.12.2018.
- Overall, clients in SHSCT were scheduled to have an average of 21.9 visits and they completed an average of 14.1 valid visits in toddlerhood.
- Overall to date for all clients who enrolled since 2012, 64% of expected visits were achieved by the end of 2018 in the SHSCT.
- For clients who enrolled in 2016, 69.2% of expected toddlerhood visits were achieved by the end of 2018.

Table 36: Dosage Toddlerhood Stage – WHSCT by year enrolled

Year Enrolled	Clients Enrolled	Toddlerhood Completer	Active Completer	Scheduled Visits	Actual Visits	Scheduled Visits	Actual Visits	Expected Visits Achieved
	N	N	N	N	N	Mean N	Mean N	Mean %
2010	7	7	5	110	71	22	14.2	64.5
2011	76	76	64	1408	1059	22	16.5	75.2
2012	22	22	19	418	327	22	17.2	78.2
2013	23	23	20	440	276	22	13.8	62.7
2014	49	49	42	924	560	22	13.3	60.6
2015	27	27	24	528	267	22	11.1	50.6
2016	28	11	8	176	66	22	8.3	37.5
Total	232	215	182	4004	2626	22	14.4	65.6

This table reports on toddlerhood dosage (WHSCT) for active clients (N=182) who had completed the toddlerhood stage by 31.12.2018.

WHSCT

- 182 clients in the WHSCT completed the toddlerhood stage by the end of 31.12.2018.
- Overall, clients in WHSCT were scheduled to have an average of 22 visits and they completed an average of 14.4 valid visits in toddlerhood.
- Overall to date for all clients who enrolled since 2010, 65.6% of expected visits were achieved by the end of 2018 in the WHSCT.
- For clients who enrolled in 2016, 37.5% of expected toddlerhood visits were achieved by the end of 2018.

Visit Duration

Pregnancy Duration

Table 37: Pregnancy Duration – All sites by year enrolled

Year Enrolled	Clients N	Valid Visits N	Visit Duration			
			Mean	Minimum	Maximum	Standard Deviation
2010	7	33	70.6	30	120	17.6
2011	76	695	64.3	15	120	12.6
2012	35	388	75.1	20	333	25.3
2013	135	1491	71.7	15	240	18.0
2014	111	1111	65.1	15	120	13.0
2015	78	801	77.0	30	210	23.8
2016	200	1872	71.8	15	240	19.5
2017	129	1079	67.7	20	210	18.1
2018	74	625	68.4	15	195	21.0
Total	845	8095	70.1	15	333	19.1

This table provides overall pregnancy visit duration by year of enrolment for all active clients (N= 845) who had completed the pregnancy stage by 31.12.2018.

All sites

- Overall, the average visit duration for valid pregnancy visits (N= 8095 visits) for pregnancy completers (N=845) was 1 hour and 10 minutes. This ranged from 15 minutes to 5 hours and 33 minutes. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.

Table 38: Pregnancy Duration – By site by year enrolled

Site	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
BHSCT	174	1536	67.3	15	240	18.9
NHSCT	100	787	75.5	30	210	19.7
SEHSCT	112	1290	73.3	15	210	22.9
SHSCT	182	1899	75.1	15	240	20.0
WHSCT	277	2583	64.8	15	333	13.8
Total	845	8095	70.1	15	333	19.1

This table provides pregnancy visit duration by site for all active clients (N= 845) who had completed the pregnancy stage by 31.12.2018.

All sites

- In BHSCT, the average visit duration for valid pregnancy visits (N= 1536 visits) for pregnancy completers (N= 174) was one hour and seven minutes. This ranged from 15 minutes to four hours. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.
- In NHSCT, the average visit duration for valid pregnancy visits (N= 787 visits) for pregnancy completers (N= 100) was one hour and 15 minutes. This ranged from 30 minutes to three hours and 30 minutes. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.
- In SEHSCT, the average visit duration for valid pregnancy visits (N=1290 visits) for pregnancy completers (N= 112) was one hour and 13 minutes. This ranged from 15 minutes to three hours and 30 minutes. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.
- In SHSCT, the average visit duration for valid pregnancy visits (N= 1899 visits) for pregnancy completers (N= 182) was one hour and 15 minutes. This ranged from 15 minutes to four hours. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.
- In WHSCT, the average visit duration for valid pregnancy visits (N= 2583 visits) for pregnancy completers (N= 277) was one hour and four minutes. This ranged from 15 minutes to five hours and 33 minutes. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.

Table 39: Pregnancy Duration – BHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N		N	Mean	Minimum	Maximum
2013	41	417	66.9	40	150	15.7
2014	40	404	63.9	22	120	12.6
2015	8	75	75.1	30	180	23.7
2016	43	333	66.4	15	240	19.2
2017	25	178	70.9	20	210	23.0
2018	17	129	72.3	30	180	28.9
Total	174	1536	67.3	15	240	18.9

This table provides pregnancy visit duration (BHSCT) for all active clients (N= 174) who had completed the pregnancy stage by 31.12.2018.

BHSCT

- In BHSCT, the average visit duration for valid pregnancy visits (N= 1536 visits) for pregnancy completers (N= 174) was one hour and seven minutes. This ranged from 15 minutes to four hours. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.

Table 40: Pregnancy Duration – NHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2015	19	179	88.8	30	210	22.2
2016	41	333	77.0	30	120	17.4
2017	27	188	66.7	30	120	14.8
2018	13	87	61.6	30	90	11.6
Total	100	787	75.5	30	210	19.7

This table provides pregnancy visit duration (NHSCT) for all active clients (N= 100) who had completed the pregnancy stage by 31.12.2018.

NHSCT

- In NHSCT, the average visit duration for valid pregnancy visits (N= 787 visits) for pregnancy completers (N= 100) was one hour and 15 minutes. This ranged from 30 minutes to three hours and 30 minutes. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.

Table 41: Pregnancy Duration – SEHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2014	1	8	78.8	60	120	22.3
2015	20	258	82.1	30	210	27.1
2016	57	665	73.6	15	180	21.5
2017	18	190	66.8	30	150	19.5
2018	16	169	65.5	15	195	20.3
Total	112	1290	73.3	15	210	22.9

This table provides pregnancy visit duration (SEHSCT) for all active clients (N= 112) who had completed the pregnancy stage by 31.12.2018.

SEHSCT

- In SEHSCT, the average visit duration for valid pregnancy visits (N=1290 visits) for pregnancy completers (N= 112) was one hour and 13 minutes. This ranged from 15 minutes to three hours and 30 minutes. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.

Table 42: Pregnancy Duration – SHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2012	13	153	89.6	30	165	23.8
2013	71	831	76.6	20	240	19.2
2014	23	240	69.6	30	120	14.3
2015	4	27	75.7	60	120	17.9
2016	31	302	72.0	15	140	19.1
2017	20	159	70.8	30	180	22.0
2018	20	187	72.2	30	150	19.6
Total	182	1899	75.1	15	240	20.0

This table provides pregnancy visit duration (SHSCT) for all active clients (N= 182) who had completed the pregnancy stage by 31.12.2018.

SHSCT

- In SHSCT, the average visit duration for valid pregnancy visits (N= 1899 visits) for pregnancy completers (N= 182) was one hour and 15 minutes. This ranged from 15 minutes to four hours. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.

Table 43: Pregnancy Duration – WHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2010	7	33	70.6	30	120	17.6
2011	76	695	64.3	15	120	12.6
2012	22	235	65.6	20	333	21.4
2013	23	243	62.9	15	120	10.1
2014	47	459	63.6	15	120	11.9
2015	27	262	64.4	30	120	14.2
2016	28	239	67.1	45	135	14.2
2017	39	364	65.8	30	150	13.3
2018	8	53	65.3	60	100	10.7
Total	277	2583	64.8	15	333	13.8

This table provides pregnancy visit duration (WHSCT) for all active clients (N= 277) who had completed the pregnancy stage by 31.12.2018.

WHSCT

- In WHSCT, the average visit duration for valid pregnancy visits (N= 2583 visits) for pregnancy completers (N= 277) was one hour and four minutes. This ranged from 15 minutes to five hours and 33 minutes. The fidelity stretch goal for length of visits is that, on average, pregnancy visits should be 60 minutes or more.

Infancy Duration

Table 44: Infancy duration – All sites by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2010	5	92	60.9	15	240	22.0
2011	69	1365	60.7	15	150	11.7
2012	32	758	68.1	15	270	19.4
2013	126	2666	68.8	15	270	18.8
2014	106	2105	65.2	15	240	17.4
2015	73	1504	71.5	15	330	23.1
2016	174	3327	66.7	15	360	21.2
2017	75	1336	65.4	15	325	20.8
Total	660	13153	66.7	15	360	19.7

This table provides overall infancy visit duration by year of enrolment for all active clients (N= 660) who had completed the infancy stage by 31.12.2018.

All sites

- Overall, the average visit duration for valid infancy visits (N= 13153 visits) for infancy completers (N= 660) was one hour and six minutes. This ranged from 15 minutes to six hours. The fidelity stretch goal for length of visits is that, on average, infancy visits should be 60 minutes or more.

Table 45: Infancy duration – By site by year enrolled

Site	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
BHSCT	132	2469	67.7	15	360	23.2
NHSCT	69	1372	72.1	20	240	19.0
SEHSCT	87	1796	65.7	15	300	21.0
SHSCT	147	3086	70.7	15	360	22.6
WHSCT	225	4430	62.1	15	270	12.7
Total	660	13153	66.7	15	360	19.7

This table provides infancy visit duration by site for all active clients (N= 660) who had completed the infancy stage by 31.12.2018.

All sites

- In BHSCT, the average visit duration for valid infancy visits (N= 2469 visits) for infancy completers (N= 132) was one hour and seven minutes. This ranged from 15 minutes to six hours.
- In NHSCT, the average visit duration for valid infancy visits (N= 1372 visits) for infancy completers (N=69) was one hour and 12 minutes. This ranged from 20 minutes to four hours.
- In SEHSCT, the average visit duration for valid infancy visits (N= 1796 visits) for infancy completers (N=87) was one hour and five minutes. This ranged from 15 minutes to five hours.
- In SHSCT, the average visit duration for valid infancy visits (N= 3086 visits) for infancy completers (N=147) was one hour and ten minutes. This ranged from 15 minutes to six hours.
- In WHSCT, the average visit duration for valid infancy visits (N= 4430 visits) for infancy completers (N=225) was one hour and two minutes. This ranged from 15 minutes to four hours and 30 minutes.

Table 46: Infancy duration – BHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2013	37	734	67.1	15	270	20.1
2014	39	760	67.8	25	240	22.5
2015	8	176	73.1	30	330	29.9
2016	37	638	67.4	15	360	25.0
2017	11	161	65.8	15	150	23.3
Total	132	2469	67.7	15	360	23.2

This table provides infancy visit duration (BHSCT) for all active clients (N= 132) who had completed the infancy stage by 31.12.2018.

BHSCT

- In BHSCT, the average visit duration for valid infancy visits (N= 2469 visits) for infancy completers (N= 132) was one hour and seven minutes. This ranged from 15 minutes to six hours.

Table 47: Infancy duration – NHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2015	17	369	81.2	60	240	20.8
2016	34	680	69.8	20	120	16.5
2017	18	323	66.6	30	120	18.1
Total	69	1372	72.1	20	240	19.0

This table provides infancy visit duration (NHSCT) for all active clients (N= 69) who had completed the infancy stage by 31.12.2018.

NHSCT

- In NHSCT, the average visit duration for valid infancy visits (N= 1372 visits) for infancy completers (N=69) was one hour and 12 minutes. This ranged from 20 minutes to four hours.

Table 48: Infancy duration – SEHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2014	1	15	77.0	30	120	25.9
2015	19	411	70.5	30	300	26.4
2016	52	1076	64.3	15	195	18.4
2017	15	294	63.5	30	180	20.4
Total	87	1796	65.7	15	300	21.0

This table provides infancy visit duration (SEHSCT) for all active clients (N=87) who had completed the infancy stage by 31.12.2018.

SEHSCT

- In SEHSCT, the average visit duration for valid infancy visits (N= 1796 visits) for infancy completers (N=87) was one hour and five minutes. This ranged from 15 minutes to five hours.

Table 49: Infancy duration – SHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2012	12	282	78.8	15	270	25.2
2013	68	1497	71.7	20	195	19.5
2014	22	437	66.4	15	130	16.7
2015	4	96	76.9	15	240	26.0
2016	29	571	66.3	15	360	26.2
2017	12	203	70.6	30	325	32.2
Total	147	3086	70.7	15	360	22.6

This table provides infancy visit duration (SHSCT) for all active clients (N=147) who had completed the infancy stage by 31.12.2018.

SHSCT

- In SHSCT, the average visit duration for valid infancy visits (N= 3086 visits) for infancy completers (N=147) was one hour and ten minutes. This ranged from 15 minutes to six hours.

Table 50: Infancy duration – WHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2010	5	92	60.9	15	240	22.0
2011	69	1365	60.7	15	150	11.7
2012	20	476	61.7	15	120	10.9
2013	21	435	61.2	15	120	9.1
2014	44	893	62.2	15	150	11.0
2015	25	452	62.8	30	180	12.2
2016	22	362	67.7	30	270	20.5
2017	19	355	62.9	30	150	11.4
Total	225	4430	62.1	15	270	12.7

This table provides infancy visit duration (WHSCT) for all active clients (N=225) who had completed the infancy stage by 31.12.2018.

WHSCT

- In WHSCT, the average visit duration for valid infancy visits (N= 4430 visits) for infancy completers (N=225) was one hour and two minutes. This ranged from 15 minutes to four hours and 30 minutes.

Toddlerhood Duration

Table 51: Toddlerhood Duration – All sites by year of enrolment

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2010	5	71	64.6	40	120	13.9
2011	64	1059	61.0	15	150	11.6
2012	31	514	64.5	15	390	20.0
2013	117	1611	69.6	15	300	23.3
2014	99	1282	66.5	15	480	22.3
2015	68	893	66.7	15	150	19.2
2016	104	1404	64.8	15	300	22.9
Total	488	6834	65.9	15	480	20.9

This table provides toddlerhood visit duration by site for all active clients (N= 488) who had completed the toddlerhood stage by 31.12.2018.

All sites

- Overall, the average visit duration for valid toddlerhood visits (N= 6834 visits) for toddlerhood completers (N=488) was one hour and five minutes. This ranged from 15 minutes to eight hours. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.

Table 52: Toddlerhood Duration – By site by year enrolled

Site	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
BHSCT	95	1287	73.5	15	480	30.9
NHSCT	38	480	70.1	30	120	17.2
SEHSCT	51	726	59.6	15	180	20.7
SHSCT	122	1715	67.3	15	390	21.7
WHSCT	182	2626	62.2	15	150	11.6
Total	488	6834	65.9	15	480	20.9

All sites

- In BHSCT, the average visit duration for valid toddlerhood visits (N= 1287 visits) for toddlerhood completers (N= 95) was one hour and 13 minutes. This ranged from 15 minutes to eight hours. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.
- In NHSCT, the average visit duration for valid toddlerhood visits (N= 480 visits) for toddlerhood completers (N=38) was one hour and 10 minutes. This ranged from 30 minutes to two hours. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.
- In SEHSCT, the average visit duration for valid toddlerhood visits (N= 726 visits) for toddlerhood completers (N=51) was 59 minutes. This ranged from 15 minutes to three hours. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.
- In SHSCT, the average visit duration for valid toddlerhood visits (N= 1715 visits) for toddlerhood completers (N=122) was one hour and seven minutes. This ranged from 15 minutes to six hours and 30 minutes. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.
- In WHSCT, the average visit duration for valid toddlerhood visits (N= 2626 visits) for toddlerhood completers (N=182) was one hour and two minutes. This ranged from 15 minutes to two hours and 30 minutes. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.

Table 53: Toddlerhood Duration – BHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2013	35	509	74.9	15	300	32.3
2014	36	448	71.5	20	480	32.3
2015	7	116	74.0	15	120	24.9
2016	17	214	73.9	15	180	27.2
Total	95	1287	73.5	15	480	30.9

This table provides toddlerhood visit duration (BHSCT) for all active clients (N= 95) who had completed the toddlerhood stage by 31.12.2018.

BHSCT

- In BHSCT, the average visit duration for valid toddlerhood visits (N= 1287 visits) for toddlerhood completers (N= 95) was one hour and 13 minutes. This ranged from 15 minutes to eight hours. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.

Table 54: Toddlerhood Duration – NHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2015	16	205	74.7	30	120	16.9
2016	22	275	66.6	30	120	16.6
Total	38	480	70.1	30	120	17.2

This table provides toddlerhood visit duration (NHSCT) for all active clients (N= 38) who had completed the toddlerhood stage by 31.12.2018.

NHSCT

- In NHSCT, the average visit duration for valid toddlerhood visits (N= 480 visits) for toddlerhood completers (N=38) was one hour and 10 minutes. This ranged from 30 minutes to two hours. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.

Table 55: Toddlerhood Duration – SEHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2015	17	227	58.8	15	150	20.8
2016	34	499	60.0	20	180	20.7
Total	51	726	59.6	15	180	20.7

This table provides toddlerhood visit duration (SEHSCT) for all active clients (N=51) who had completed the toddlerhood stage by 31.12.2018.

SEHSCT

- In SEHSCT, the average visit duration for valid toddlerhood visits (N= 726 visits) for toddlerhood completers (N=51) was 59 minutes. This ranged from 15 minutes to three hours. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.

Table 56: Toddlerhood Duration – SHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2012	12	187	70.4	30	390	29.1
2013	62	826	68.5	16	150	18.3
2014	21	274	66.2	15	150	18.5
2015	4	78	64.3	30	105	17.1
2016	23	350	64.4	15	300	26.7
Total	122	1715	67.3	15	390	21.7

This table provides toddlerhood visit duration (SHSCT) for all active clients (N=122) who had completed the toddlerhood stage by 31.12.2018.

SHSCT

- In SHSCT, the average visit duration for valid toddlerhood visits (N= 1715 visits) for toddlerhood completers (N=122) was one hour and seven minutes. This ranged from 15 minutes to six hours and 30 minutes. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more

Table 57: Toddlerhood Duration – WHSCT by year enrolled

Year Enrolled	Clients	Valid Visits	Visit Duration			
	N	N	Mean	Minimum	Maximum	Standard Deviation
2010	5	71	64.6	40	120	13.9
2011	64	1059	61.0	15	150	11.6
2012	19	327	61.1	15	120	10.7
2013	20	276	63.1	30	120	12.0
2014	42	560	62.5	30	120	10.2
2015	24	267	64.9	45	150	12.9
2016	8	66	66.5	45	120	14.1
Total	182	2626	62.2	15	150	11.6

This table provides toddlerhood visit duration (WHSCT) for all active clients (N=182) who had completed the toddlerhood stage by 31.12.2018.

WHSCT

- In WHSCT, the average visit duration for valid toddlerhood visits (N= 2626 visits) for toddlerhood completers (N=182) was one hour and two minutes. This ranged from 15 minutes to two hours and 30 minutes. The fidelity stretch goal for length of visits is that, on average, toddlerhood visits should be 60 minutes or more.

Visit Content

Visit Content Pregnancy Completers

Table 58: Visit content – Overall active pregnancy completers by site (until end of 2018)

Site	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (23-25%)	Personal Health (35-40%)	Environmental Health (5-7%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
BHSCT	1536	13.2	14.1	26.4	35.9	10.5	100
NHSCT	787	12.7	13.7	26.6	39.5	7.6	100
SEHSCT	1290	9.9	16.2	29.8	35.2	9.0	100
SHSCT	1899	13.1	14.2	26.7	38.6	7.2	100
WHSCT	2583	11.2	16.2	28.6	34.1	9.9	100
Total	8095	12.0	15.1	27.7	36.2	9.0	100

Table 59: Visit content – All sites active pregnancy completers by year enrolled (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (23-25%)	Personal Health (35-40%)	Environmental Health (5-7%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2010	33	9.8	15.9	22.6	41.5	10.2	100
2011	695	10.9	14.6	29.3	35.8	9.3	100
2012	388	12.2	15.9	26.9	35.2	9.8	100
2013	1491	12.9	15.6	25.9	38.0	7.6	100
2014	1111	12.3	14.9	26.7	36.6	9.5	100
2015	801	11.4	15.1	27.6	35.9	10.0	100
2016	1872	11.5	14.6	29.1	35.8	9.0	100
2017	1079	12.4	15.1	27.8	35.7	9.0	100
2018	625	11.7	15.6	28.7	34.3	9.6	100
Total	8095	12.0	15.1	27.7	36.2	9.0	100

These tables provides data on visit content for pregnancy completers for all sites for all active clients (N= 845) who had completed the pregnancy stage by 31.12.2018.

All sites

- Overall (N=8095 valid visits), life course development (12%) was within the visit content goal (10-15%), family and friends (15.1%) was just above the visit content goal (10-15%), maternal role (27.7%) was above the visit content goal (23-25%), personal health (36.2%) was within the visit content goal (35-40%) and environmental health (9%) was above the visit content goal (5-7%).
- In BHSCT (N=1536 valid visits), life course development (13.2%) was within the visit content goal (10-15%), family and friends (14.1%) was within the visit content goal (10-15%), maternal role (26.4%) was above the visit content goal (23-25%), personal health (35.9%) was within the visit content goal (35-40%) and environmental health (10.5%) was above the visit content goal (5-7%).
- In NHSCT (N=787 valid visits), life course development (12.7%) was within the visit content goal (10-15%), family and friends (13.7%) was within the visit content goal (10-15%), maternal role (26.6%) was above the visit content goal (23-25%), personal health (39.5%) was within the visit content goal (35-40%) and environmental health (7.6%) was just above the visit content goal (5-7%).
- In SEHSCT (N=1290 valid visits), life course development (9.9%) was just below the visit content goal (10-15%), family and friends (16.2%) was above the visit content goal (10-15%), maternal role (29.8%) was above the visit content goal (23-25%), personal health (35.2%) was within the visit content goal (35-40%) and environmental health (9%) was above the visit content goal (5-7%).
- In SHSCT (N=1899 valid visits), life course development (13.1%) was within the visit content goal (10-15%), family and friends (14.2%) was within the visit content goal (10-15%), maternal role (26.7%) was above the visit content goal (23-25%), personal health (38.6%) was within the visit content goal (35-40%) and environmental health (7.2%) was just above the visit content goal (5-7%).
- In WHSCT (N=2583 valid visits), life course development (11.2%) was within the visit content goal (10-15%), family and friends (16.2%) was above the visit content goal (10-15%), maternal role (28.6%) was above the visit content goal (23-25%), personal health (34.1%) was just below the visit content goal (35-40%) and environmental health (9.9%) was above the visit content goal (5-7%).

Table 60: Visit content – BHSCT active pregnancy completers by year enrolled (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (23-25%)	Personal Health (35-40%)	Environmental Health (5-7%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2013	417	12.6	16.5	25.0	37.8	8.0	100
2014	404	13.3	13.0	25.3	38.3	10.2	100
2015	75	14.3	11.6	27.3	33.3	13.5	100
2016	333	14.2	13.3	26.8	32.9	13.1	100
2017	178	14.1	13.4	27.4	34.7	10.3	100
2018	129	10.4	14.2	30.9	32.9	11.6	100
Total	1536	13.2	14.1	26.4	35.9	10.5	100

This table provides data on visit content for pregnancy completers (BHSCT) for all active clients (N= 174) who had completed the pregnancy stage by 31.12.2018.

BHSCT

- In BHSCT (N=1536 valid visits), life course development (13.2%) was within the visit content goal (10-15%), family and friends (14.1%) was within the visit content goal (10-15%), maternal role (26.4%) was above the visit content goal (23-25%), personal health (35.9%) was within the visit content goal (35-40%) and environmental health (10.5%) was above the visit content goal (5-7%).

Table 61: Visit content – NHSCT active pregnancy completers by year enrolled (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (23-25%)	Personal Health (35-40%)	Environmental Health (5-7%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2015	179	12.6	14.1	26.4	38.7	8.2	100
2016	333	12.5	13.7	27.1	39.4	7.4	100
2017	188	13.2	13.5	25.6	40.4	7.3	100
2018	87	12.6	13.3	27.1	39.3	7.8	100
Total	787	12.7	13.7	26.6	39.5	7.6	100

This table provides data on visit content for pregnancy completers (NHSCT) for all active clients (N= 100) who had completed the pregnancy stage by 31.12.2018.

NHSCT

- In NHSCT (N=787 valid visits), life course development (12.7%) was within the visit content goal (10-15%), family and friends (13.7%) was within the visit content goal (10-15%), maternal role (26.6%) was above the visit content goal (23-25%), personal health (39.5%) was within the visit content goal (35-40%) and environmental health (7.6%) was just above the visit content goal (5-7%).

Table 62: Visit content – SEHSCT active pregnancy completers by year enrolled (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (23-25%)	Personal Health (35-40%)	Environmental Health (5-7%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2014	8	5.6	28.8	25.6	33.8	6.3	100
2015	258	9.4	14.7	27.1	39.4	9.4	100
2016	665	9.4	16.1	31.0	36.0	7.6	100
2017	190	11.5	16.4	29.8	30.8	11.5	100
2018	169	10.9	18.1	29.1	30.5	11.4	100
Total	1290	9.9	16.2	29.8	35.2	9.0	100

This table provides data on visit content for pregnancy completers (SEHSCT) for all active clients (N= 112) who had completed the pregnancy stage by 31.12.2018.

SEHSCT

- In SEHSCT (N=1290 valid visits), life course development (9.9%) was just below the visit content goal (10-15%), family and friends (16.2%) was above the visit content goal (10-15%), maternal role (29.8%) was above the visit content goal (23-25%), personal health (35.2%) was within the visit content goal (35-40%) and environmental health (9%) was above the visit content goal (5-7%).

Table 63: Visit content – SHSCT active pregnancy completers by year enrolled (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (23-25%)	Personal Health (35-40%)	Environmental Health (5-7%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2012	153	12.7	15.3	24.8	39.1	8.1	100
2013	831	13.3	14.7	25.5	39.7	6.8	100
2014	240	13.3	14.0	25.5	39.9	7.3	100
2015	27	14.6	14.6	26.7	38.1	5.9	100
2016	302	12.4	12.4	30.5	36.5	8.2	100
2017	159	13.5	14.2	28.7	36.7	6.9	100
2018	187	13.3	14.5	27.6	37.1	7.5	100
Total	1899	13.1	14.2	26.7	38.6	7.2	100

This table provides data on visit content for pregnancy completers (SHSCT) for all active clients (N= 182) who had completed the pregnancy stage by 31.12.2018.

SHSCT

- In SHSCT (N=1899 valid visits), life course development (13.1%) was within the visit content goal (10-15%), family and friends (14.2%) was within the visit content goal (10-15%), maternal role (26.7%) was above the visit content goal (23-25%), personal health (38.6%) was within the visit content goal (35-40%) and environmental health (7.2%) was just above the visit content goal (5-7%).

Table 64: Visit content – WHSCT active pregnancy completers by year enrolled (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (23-25%)	Personal Health (35-40%)	Environmental Health (5-7%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2010	33	9.8	15.9	22.6	41.5	10.2	100
2011	695	10.9	14.6	29.3	35.8	9.3	100
2012	235	11.9	16.3	28.3	32.6	10.8	100
2013	243	11.6	17.1	29.0	32.7	9.5	100
2014	459	11.1	16.8	28.6	33.3	10.2	100
2015	262	11.4	17.1	28.9	31.2	11.4	100
2016	239	11.3	16.3	28.2	33.8	10.4	100
2017	364	11.3	16.3	27.7	35.8	8.9	100
2018	53	10.7	19.1	28.2	32.5	9.6	100
Total	2583	11.2	16.2	28.6	34.1	9.9	100

This table provides data on visit content for pregnancy completers (WHSCT) for all active clients (N= 277) who had completed the pregnancy stage by 31.12.2018.

WHSCT

- In WHSCT (N=2583 valid visits), life course development (11.2%) was within the visit content goal (10-15%), family and friends (16.2%) was above the visit content goal (10-15%), maternal role (28.6%) was above the visit content goal (23-25%), personal health (34.1%) was just below the visit content goal (35-40%) and environmental health (9.9%) was above the visit content goal (5-7%).

Visit Content Infancy completers

Table 65: Visit content – active infancy completers by site (until end of 2018)

Site	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (45-50%)	Personal Health (14-20%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
BHSCT	2469	13.2	13.6	40.9	20.1	12.2	100
NHSCT	1372	11.2	11.6	46.8	20.5	9.9	100
SEHSCT	1796	9.6	15	40.9	22.4	12.1	100
SHSCT	3086	11.1	11.4	44.9	22.3	10.4	100
WHSCT	4430	10.4	14.6	41.7	22.6	10.6	100
Total	13153	11.1	13.4	42.7	21.8	11.0	100

Table 66: Visit content - All sites active infancy completers by year of enrolment (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (45-50%)	Personal Health (14-20%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2010	92	10.3	13	43.2	23.2	10.4	100
2011	1365	10.2	14.5	43.8	21.7	9.8	100
2012	758	11.5	13.4	42.5	22.1	10.6	100
2013	2666	11.4	12.4	44.1	21.5	10.6	100
2014	2105	11.2	13.6	42.6	21.6	11.0	100
2015	1504	11.1	13.6	42	22.6	10.7	100
2016	3327	11.2	13.7	41.5	21.8	11.8	100
2017	1336	10.6	13.1	43.3	21.7	11.3	100
Total	13153	11.1	13.4	42.7	21.8	11.0	100

These tables provide data on visit content for infancy completers for all sites for all active clients (N= 660) who had completed the infancy stage by 31.12.2018.

All sites

- Overall (N=13153 valid visits), life course development (11.1%) was within the visit content goal (10-15%), family and friends (13.4%) was within the visit content goal (10-15%), maternal role (42.7%) was below the visit content goal (45-50%), personal health (21.8%) was above the visit content goal (14-20%) and environmental health (11%) was above the visit content goal (7-10%).
- In BHSCT (N=2469 valid visits), life course development (13.2%) was within the visit content goal (10-15%), family and friends (13.6%) was within the visit content goal (10-15%), maternal role (40.9%) was below the visit content goal (45-50%), personal health (20.1%) was just above the visit content goal (14-20%) and environmental health (12.2%) was above the visit content goal (7-10%).
- In NHSCT (N=1372 valid visits), life course development (11.2%) was within the visit content goal (10-15%), family and friends (11.6%) was within the visit content goal (10-15%), maternal role (46.8%) was within the visit content goal (45-50%), personal health (20.5%) was just above the visit content goal (14-20%) and environmental health (9.9%) was within the visit content goal (7-10%).
- In SEHSCT (N=1796 valid visits), life course development (9.6%) was just below the visit content goal (10-15%), family and friends (15%) was within the visit content goal (10-15%), maternal role (40.9%) was below the visit content goal (45-50%), personal health (22.4%) was above the visit content goal (14-20%) and environmental health (12.1%) was above the visit content goal (7-10%).
- In SHSCT (N=3086 valid visits), life course development (11.1%) was within the visit content goal (10-15%), family and friends (11.4%) was within the visit content goal (10-15%), maternal role (44.9%) was just below the visit content goal (45-50%), personal health (22.3%) was above the visit content goal (14-20%) and environmental health (10.4%) was just above the visit content goal (7-10%).
- In WHSCT (N=4430 valid visits), life course development (10.4%) was within the visit content goal (10-15%), family and friends (14.6%) was within the visit content goal (10-15%), maternal role (41.7%) was below the visit content goal (45-50%), personal health (22.6%) was above the visit content goal (14-20%) and environmental health (10.6%) was just above the visit content goal (7-10%).

Table 67: Visit content – BHSCT active infancy completers by year of enrolment (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (45-50%)	Personal Health (14-20%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2013	734	12.8	13.6	41.9	20.8	10.9	100
2014	760	12.6	13.3	42.7	19.6	11.7	100
2015	176	13.9	14.1	38.0	22.3	11.7	100
2016	638	14.1	14.1	38.1	19.5	14.3	100
2017	161	13.2	13.4	42.4	18.8	12.3	100
Total	2469	13.2	13.6	40.9	20.1	12.2	100

This table provides data on visit content for infancy completers (BHSCT) for all active clients (N= 132) who had completed the infancy stage by 31.12.2018.

BHSCT

- In BHSCT (N=2469 valid visits), life course development (13.2%) was within the visit content goal (10-15%), family and friends (13.6%) was within the visit content goal (10-15%), maternal role (40.9%) was below the visit content goal (45-50%), personal health (20.1%) was just above the visit content goal (14-20%) and environmental health (12.2%) was above the visit content goal (7-10%).

Table 68: Visit content – NHSCT active infancy completers by year of enrolment (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (45-50%)	Personal Health (14-20%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2015	369	11.6	11.7	46.1	21.2	9.4	100
2016	680	11.1	11.5	47.6	20.1	9.7	100
2017	323	10.8	11.8	45.9	20.4	10.9	100
Total	1372	11.2	11.6	46.8	20.5	9.9	100

This table provides data on visit content for infancy completers (NHSCT) for all active clients (N=69) who had completed the infancy stage by 31.12.2018.

NHSCT

- In NHSCT (N=1372 valid visits), life course development (11.2%) was within the visit content goal (10-15%), family and friends (11.6%) was within the visit content goal (10-15%), maternal role (46.8%) was within the visit content goal (45-50%), personal health (20.5%) was just above the visit content goal (14-20%) and environmental health (9.9%) was within the visit content goal (7-10%).

Table 69: Visit content – SEHSCT active infancy completers by year of enrolment (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (45-50%)	Personal Health (14-20%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2014	15	13.3	20.0	35.7	19.0	12.0	100
2015	411	9.5	13.4	43.4	23.3	10.5	100
2016	1076	9.8	15.5	39.9	22.3	12.5	100
2017	294	8.7	15.0	41.4	22.0	13.0	100
Total	1796	9.6	15.0	40.9	22.4	12.1	100

This table provides data on visit content for infancy completers (SEHSCT) for all active clients (N= 87) who had completed the infancy stage by 31.12.2018.

SEHSCT

- In SEHSCT (N=1796 valid visits), life course development (9.6%) was just below the visit content goal (10-15%), family and friends (15%) was within the visit content goal (10-15%), maternal role (40.9%) was below the visit content goal (45-50%), personal health (22.4%) was above the visit content goal (14-20%) and environmental health (12.1%) was above the visit content goal (7-10%).

Table 70: Visit content – SHSCT active infancy completers by year of enrolment (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (45-50%)	Personal Health (14-20%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2012	282	11.6	12.3	45.7	20.4	10.1	100
2013	1497	10.8	11.2	45.9	21.9	10.2	100
2014	437	10.9	11.8	44.8	21.4	11.2	100
2015	96	13.9	13.5	40.9	22.1	9.7	100
2016	571	11.1	11.0	43.1	24.4	10.4	100
2017	203	12.0	10.8	43.2	23.5	10.5	100
Total	3086	11.1	11.4	44.9	22.3	10.4	100

This table provides data on visit content for infancy completers (SHSCT) for all active clients (N= 147) who had completed the infancy stage by 31.12.2018.

SHSCT

- In SHSCT (N=3086 valid visits), life course development (11.1%) was within the visit content goal (10-15%), family and friends (11.4%) was within the visit content goal (10-15%), maternal role (44.9%) was just below the visit content goal (45-50%), personal health (22.3%) was above the visit content goal (14-20%) and environmental health (10.4%) was just above the visit content goal (7-10%).

Table 71: Visit content – WHSCT active infancy completers by year of enrolment (until end of 2018)

Year Enrolled	Visits	Life Course Development (10-15%)	Family and Friends (10-15%)	Maternal Role (45-50%)	Personal Health (14-20%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2010	92	10.3	13.0	43.2	23.2	10.4	100
2011	1365	10.2	14.5	43.8	21.7	9.8	100
2012	476	11.4	14.0	40.5	23.1	11.0	100
2013	435	11.0	14.7	41.7	21.3	11.2	100
2014	893	10.0	14.7	41.6	23.4	10.4	100
2015	452	10.6	15.2	39.0	23.3	11.9	100
2016	362	10.7	16.0	37.8	23.9	11.7	100
2017	355	9.9	14.1	43.0	22.8	10.2	100
Total	4430	10.4	14.6	41.7	22.6	10.6	100

This table provides data on visit content for infancy completers (WHSCT) for all active clients (N=225) who had completed the infancy stage by 31.12.2018.

WHSCT

- In WHSCT (N=4430 valid visits), life course development (10.4%) was within the visit content goal (10-15%), family and friends (14.6%) was within the visit content goal (10-15%), maternal role (41.7%) was below the visit content goal (45-50%), personal health (22.6%) was above the visit content goal (14-20%) and environmental health (10.6%) was just above the visit content goal (7-10%).

Visit Content Toddlerhood Completers

Table 72: Visit Content – Toddlerhood completers by site (up to end of December 2018)

Site	Visits	Life Course Development (18-20%)	Family and Friends (10-15%)	Maternal Role (40-45%)	Personal Health (10-15%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
BHSCT	1287	16.6	13.0	38.6	19.0	12.9	100
NHSCT	480	17.0	12.4	44.1	15.7	10.8	100
SEHSCT	726	12.2	16.1	37.0	19.4	15.3	100
SHSCT	1715	15.3	12.9	42.1	19.1	10.6	100
WHSCT	2626	13.4	15.7	39.7	19.8	11.4	100
Total	6834	14.6	14.3	40.1	19.2	11.9	100

Table 73: Visit Content – Toddlerhood (All sites) completers by year of enrolment (up to end of December 2018)

Year Enrolled	Visits	Life Course Development (18-20%)	Family and Friends (10-15%)	Maternal Role (40-45%)	Personal Health (10-15%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2010	71	14.5	15.4	42.6	16.8	10.8	100
2011	1059	13.9	15.6	40.2	18.7	11.5	100
2012	514	13.0	14.5	40.1	21.4	11.0	100
2013	1611	15.7	13.1	41.4	18.5	11.3	100
2014	1282	14.6	14.6	40.2	19.4	11.2	100
2015	893	14.4	14.8	38.6	19.0	13.3	100
2016	1404	14.4	13.9	39.4	19.4	12.8	100
Total	6834	14.6	14.3	40.1	19.2	11.9	100

These tables provide data on visit content for toddlerhood completers for all sites for all active clients (N= 488) who had completed the toddlerhood stage by 31.12.2018.

All sites

- Overall (N= 6834 valid visits), life course development (14.6%) was below the visit content goal (18-20%), family and friends (14.3%) was within the visit content goal (10-15%), maternal role (40.1%) was within the visit content goal (40-45%), personal health (19.2%) was above the visit content goal (10-15%) and environmental health (11.9%) was above the visit content goal (7-10%).
- In BHSCT (N= 1287 valid visits), life course development (16.6%) was below the visit content goal (18-20%), family and friends (13.0%) was within the visit content goal (10-15%), maternal role (38.6%) was below the visit content goal (40-45%), personal health (19%) was above the visit content goal (10-15%) and environmental health (12.9%) was above the visit content goal (7-10%).
- In NHSCT (N= 480 valid visits), life course development (17.0%) was below the visit content goal (18-20%), family and friends (12.4%) was within the visit content goal (10-15%), maternal role (44.1%) was within the visit content goal (40-45%), personal health (15.7%) was just above the visit content goal (10-15%) and environmental health (10.8%) was just above the visit content goal (7-10%).
- In SEHSCT (N= 726 valid visits), life course development (12.2%) was below the visit content goal (18-20%), family and friends (16.1%) was above the visit content goal (10-15%), maternal role (37%) was below the visit content goal (40-45%), personal health (19.4%) was above the visit content goal (10-15%) and environmental health (15.3%) was above the visit content goal (7-10%).
- In SHSCT (N= 1715 valid visits), life course development (15.3%) was below the visit content goal (18-20%), family and friends (12.9%) was within the visit content goal (10-15%), maternal role (42.1%) was within the visit content goal (40-45%), personal health (19.1%) was above the visit content goal (10-15%) and environmental health (10.6%) was just above the visit content goal (7-10%).
- In WHSCT (N= 2626 valid visits), life course development (13.4%) was below the visit content goal (18-20%), family and friends (15.7%) was just above the visit content goal (10-15%), maternal role (39.7%) was just below the visit content goal (40-45%), personal health (19.8%) was above the visit content goal (10-15%) and environmental health (11.4%) was above the visit content goal (7-10%).

Table 74: Visit Content – BHSCT toddlerhood completers by year of enrolment (up to end of December 2018)

Year Enrolled	Visits	Life Course Development (18-20%)	Family and Friends (10-15%)	Maternal Role (40-45%)	Personal Health (10-15%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2013	509	16.8	12.3	39.7	18.6	12.6	100
2014	448	16.2	13.5	39.3	19.1	12.0	100
2015	116	16.6	13.8	34.7	20.0	14.9	100
2016	214	16.9	12.9	36.9	18.9	14.3	100
Total	1287	16.6	13.0	38.6	19.0	12.9	100

This table provides data on visit content for toddlerhood completers (BHSCT) for all active clients (N= 95) who had completed the toddlerhood stage by 31.12.2018.

BHSCT

- In BHSCT (N= 1287 valid visits), life course development (16.6%) was below the visit content goal (18-20%), family and friends (13.0%) was within the visit content goal (10-15%), maternal role (38.6%) was below the visit content goal (40-45%), personal health (19%) was above the visit content goal (10-15%) and environmental health (12.9%) was above the visit content goal (7-10%).

Table 75: Visit Content – NHSCT toddlerhood completers by year of enrolment (up to end of December 2018)

Year Enrolled	Visits	Life Course Development (18-20%)	Family and Friends (10-15%)	Maternal Role (40-45%)	Personal Health (10-15%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2015	205	16.6	12.7	44.6	15.7	10.3	100
2016	275	17.2	12.2	43.7	15.7	11.1	100
Total	480	17.0	12.4	44.1	15.7	10.8	100

This table provides data on visit content for toddlerhood completers (NHSCT) for all active clients (N= 38) who had completed the toddlerhood stage by 31.12.2018.

NHSCT

- In NHSCT (N= 480 valid visits), life course development (17.0%) was below the visit content goal (18-20%), family and friends (12.4%) was within the visit content goal (10-15%), maternal role (44.1%) was within the visit content goal (40-45%), personal health (15.7%) was just above the visit content goal (10-15%) and environmental health (10.8%) was just above the visit content goal (7-10%).

Table 76: Visit Content – SEHSCT toddlerhood completers by year of enrolment (up to end of December 2018)

Year Enrolled	Visits	Life Course Development (18-20%)	Family and Friends (10-15%)	Maternal Role (40-45%)	Personal Health (10-15%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2015	227	13.0	14.9	38.1	17.8	16.2	100
2016	499	11.8	16.6	36.5	20.1	14.9	100
Total	726	12.2	16.1	37.0	19.4	15.3	100

This table provides data on visit content for toddlerhood completers (SEHSCT) for all active clients (N= 51) who had completed the toddlerhood stage by 31.12.2018.

SEHSCT

- In SEHSCT (N= 726 valid visits), life course development (12.2%) was below the visit content goal (18-20%), family and friends (16.1%) was above the visit content goal (10-15%), maternal role (37%) was below the visit content goal (40-45%), personal health (19.4%) was above the visit content goal (10-15%) and environmental health (15.3%) was above the visit content goal (7-10%).

Table 77: Visit Content – SHSCT toddlerhood completers by year of enrolment (up to end of December 2018)

Year Enrolled	Visits	Life Course Development (18-20%)	Family and Friends (10-15%)	Maternal Role (40-45%)	Personal Health (10-15%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2012	187	13.4	13.6	42.3	20.3	10.5	100
2013	826	15.7	13.1	42.5	18.2	10.6	100
2014	274	16.4	13.6	41.8	17.2	11.0	100
2015	78	14.8	11.7	41.3	21.3	10.9	100
2016	350	14.5	11.7	41.7	21.8	10.4	100
Total	1715	15.3	12.9	42.1	19.1	10.6	100

This table provides data on visit content for toddlerhood completers (SHSCT) for all active clients (N= 122) who had completed the toddlerhood stage by 31.12.2018.

SHSCT

- In SHSCT (N= 1715 valid visits), life course development (15.3%) was below the visit content goal (18-20%), family and friends (12.9%) was within the visit content goal (10-15%), maternal role (42.1%) was within the visit content goal (40-45%), personal health (19.1%) was above the visit content goal (10-15%) and environmental health (10.6%) was just above the visit content goal (7-10%).

Table 78: Visit Content – WHSCT toddlerhood completers by year of enrolment (up to end of December 2018)

Year Enrolled	Visits	Life Course Development (18-20%)	Family and Friends (10-15%)	Maternal Role (40-45%)	Personal Health (10-15%)	Environmental Health (7-10%)	Total
	N	Mean %	Mean %	Mean %	Mean %	Mean %	Mean %
2010	71	14.5	15.4	42.6	16.8	10.8	100
2011	1059	13.9	15.6	40.2	18.7	11.5	100
2012	327	12.8	15.1	38.8	22.1	11.2	100
2013	276	13.9	14.4	41.5	19.1	11.1	100
2014	560	12.5	15.9	40.2	20.7	10.7	100
2015	267	12.9	17.5	35.2	21.4	13.0	100
2016	66	14.1	15.9	40.2	18.7	11.1	100
Total	2626	13.4	15.7	39.7	19.8	11.4	100

This table provides data on visit content for toddlerhood completers (WHSCT) for all active clients (N= 182) who had completed the toddlerhood stage by 31.12.2018.

WHSCT

- In WHSCT (N= 2626 valid visits), life course development (13.4%) was below the visit content goal (18-20%), family and friends (15.7%) was just above the visit content goal (10-15%), maternal role (39.7%) was just below the visit content goal (40-45%), personal health (19.8%) was above the visit content goal (10-15%) and environmental health (11.4%) was above the visit content goal (7-10%).

Father Engagement

Table 79: Father Engagement – All sites and by stage for active completers by 31.12.2018

Fathers Engagement**	Overall %	BHSCT %	NHSCT %	SEHSCT %	SHSCT %	WHSCT %
Pregnancy Completers*	16.6	14.2	14.6	23.2	14.9	16.5
	(N=1340/8095)	(N=218/1536)	(N=115/787)	(N=299/1290)	(N=283/1899)	(N=425/2583)
Infancy Completers*	16.5	13.4	12.8	19.2	16.9	18.1
	(N=2172/13153)	(N=330/2469)	(N=176/1372)	(N=344/1796)	(N=522/3086)	(N=800/4430)
Toddlerhood Completers*	12.7	12.0	8.5	12.5	11.2	14.9
	(N=871/6834)	(N=155/1287)	(N=41/480)	(N=91/726)	(N=192/1715)	(N=392/2626)

*Active Stage Completers **Attendance at visits

All sites

- Overall, father engagement was 16.6% for pregnancy completers, 16.5% for infancy completers and 12.7% for toddlerhood completers.
- For the BHSCT, father engagement was 14.2% for pregnancy completers, 13.4% for infancy completers and 12% for toddlerhood completers.
- For the NHSCT, father engagement was 14.6% for pregnancy completers, 12.8% for infancy completers and 8.5% for toddlerhood completers.
- For the SEHSCT, father engagement was 23.2% for pregnancy completers, 19.2% for infancy completers and 12.5% for toddlerhood completers.
- For the SHSCT, father engagement was 14.9% for pregnancy completers, 16.9% for infancy completers and 11.2% for toddlerhood completers.
- For the WHSCT, father engagement was 16.5% for pregnancy completers, 18.1% for infancy completers and 14.9% for toddlerhood completers.

Client Parent Engagement

Table 80: Client parent engagement – All sites and by stage for active completers by 31.12.2018

Clients parents engagement**	Overall %	BHSCT %	NHSCT %	SEHSCT %	SHSCT %	WHSCT %
Pregnancy Completers*	18.7	20.6	11.6	21.6	16.5	20.0
	(N=1516/8095)	(N=317/1536)	(N=91/787)	(N=278/1290)	(N=313/1899)	(N=517/2583)
Infancy Completers*	17.3	16.3	9.9	21.3	16.5	19.0
	(N=2273/13153)	(N=402/2469)	(N=136/1372)	(N=383/1796)	(N=510/3086)	(N=842/4430)
Toddlerhood Completers*	10.8	6.6	4.6	13.5	10.8	13.2
	(N=737/6834)	(N=85/1287)	(N=22/480)	(N=98/726)	(N=186/1715)	(N=346/2626)

*Active Stage Completers **Attendance at visits

All sites

- Overall, client parent engagement was 18.7% for pregnancy completers, 17.3% for infancy completers and 10.8% for toddlerhood completers.
- For the BHSCT, client parent engagement was 20.6% for pregnancy completers, 16.3% for infancy completers and 6.6% for toddlerhood completers
- For the NHSCT, client parent engagement was 11.6% for pregnancy completers, 9.9% for infancy completers and 4.6% for toddlerhood completers
- For the SEHSCT, client parent engagement was 21.6% for pregnancy completers, 21.3% for infancy completers and 13.5% for toddlerhood completers
- For the SHSCT, client parent engagement was 16.5% for pregnancy completers, 16.5% for infancy completers and 10.8% for toddlerhood completers.
- For the WHSCT, client parent engagement was 20% for pregnancy completers, 19% for infancy completers and 13.2% for toddlerhood completers.

Client Retention

Table 81: Client Retention by Year – All sites

Year Enrolled	Total Ever Enrolled	Active Pregnancy Completer		Active Infancy Completer		Active Toddlerhood Completer	
		N	%	N	%	N	%
2010	7	7	100	5	71.4	5	71.4
2011	76	76	100	69	90.8	64	84.2
2012	35	35	100	32	91.4	31	88.6
2013	138	135	97.8	126	91.3	117	84.8
2014	116	111	95.7	106	91.4	99	85.3
2015	80	78	97.5	73	91.3	68	85.0
2016	202	200	99.0	174	86.1	104	51.5
2017	132	129	97.7	75	56.8	0	0.0
2018	127	74	58.3	0	0.0	0	0.0
Total Enrolled*	913	845	92.6	660	72.3	488	53.5
Total Enrolled**	913	860	94.2	734	80.4	588	64.4

*Total Ever Enrolled Active Completers.

** Total Ever Enrolled Active and Inactive Completers.

All sites

- A total of 913 clients were enrolled on the programme and received a valid visit by the end of December 2018.
- To date, 845 active clients have completed the pregnancy stage, 660 active clients have completed the infancy stage and 488 active clients have completed the toddlerhood stage.

Table 82: Client Retention by Year – All sites

Site	Year Enrolled	Total Ever Enrolled	Active Pregnancy Completer		Active Infancy Completer		Active Toddlerhood Completer	
			N	%	N	%	N	%
BHSCT	2013	43	41	95.3	37	86.0	35	81.4
	2014	43	40	93.0	39	90.7	36	83.7
	2015	8	8	100.0	8	100.0	7	87.5
	2016	43	43	100.0	37	86.0	17	39.5
	2017	25	25	100.0	11	44.0	0	0.0
	2018	21	17	81.0	0	0.0	0	0.0
BHSCT Total		183	174	95.1	132	71.7	95	51.6
NHSCT	2015	19	19	100.0	17	89.5	16	84.2
	2016	41	41	100.0	34	82.9	22	53.7
	2017	28	27	96.4	18	64.3	0	0.0
	2018	22	13	59.1	0	0.0	0	0.0
NHSCT Total		110	100	90.9	69	62.7	38	34.5
SEHSCT	2014	1	1	100.0	1	100.0	0	0.0
	2015	21	20	95.2	19	90.5	17	81.0
	2016	58	57	98.3	52	89.7	34	58.6
	2017	18	18	100.0	15	83.3	0	0.0
	2018	40	16	40.0	0	0.0	0	0.0
SEHSCT Total		138	112	81.2	87	63.0	51	37.0
SHSCT	2012	13	13	100.0	12	92.3	12	92.3
	2013	72	71	98.6	68	94.4	62	86.1
	2014	23	23	100.0	22	95.7	21	91.3
	2015	5	4	80.0	4	80.0	4	80.0

	2016	32	31	96.9	29	90.6	23	71.9
	2017	20	20	100.0	12	60.0	0	0.0
	2018	28	20	71.4	0	0.0	0	0.0
SHSCT Total		193	182	94.3	147	76.6	122	63.5
WHSCT	2010	7	7	100.0	5	71.4	5	71.4
	2011	76	76	100.0	69	90.8	64	84.2
	2012	22	22	100.0	20	90.9	19	86.4
	2013	23	23	100.0	21	91.3	20	87.0
	2014	49	47	95.9	44	89.8	42	85.7
	2015	27	27	100.0	25	92.6	24	88.9
	2016	28	28	100.0	22	78.6	8	28.6
	2017	41	39	95.1	19	46.3	0	0.0
	2018	16	8	50.0	0	0.0	0	0.0
WHSCT Total		289	277	95.8	225	77.9	182	63.0
All site total		913	845	92.6	660	72.3	488	53.5

BHSCT

- A total of 183 clients were enrolled on the programme and received a valid visit by the end of December 2018.
- To date, 174 active clients have completed the pregnancy stage, 132 active clients have completed the infancy stage and 95 clients have completed the toddlerhood stage.

NHSCT

- A total of 110 clients were enrolled on the programme and received a valid visit by the end of December 2018.
- To date, 100 active clients have completed the pregnancy stage, 69 active clients have completed the infancy stage and 38 clients have completed the toddlerhood stage.

SEHSCT

- A total of 138 clients were enrolled on the programme and received a valid visit by the end of December 2018.
- To date, 112 active clients have completed the pregnancy stage, 87 active clients have completed the infancy stage and 51 clients have completed the toddlerhood stage.

SHSCT

- A total of 193 clients were enrolled on the programme and received a valid visit by the end of December 2018.
- To date, 182 active clients have completed the pregnancy stage, 147 active clients have completed the infancy stage and 122 clients have completed the toddlerhood stage.

WHSCT

- A total of 289 clients were enrolled on the programme and received a valid visit by the end of December 2018.
- To date, 277 active clients have completed the pregnancy stage, 225 active clients have completed the infancy stage and 182 clients have completed the toddlerhood stage.

Client Attrition

Table 83: Attrition by Stage and Cumulative Attrition for Stage Completers (All sites)

Client Retention	Total ever enrolled and stage completers (including inactive completers)	Active Completer*	% those enrolled who complete the programme**	Leaver/Inactive during stage***	% attrition during stage****
Total Ever Enrolled	913	NA	NA	NA	NA
Pregnancy Completer	860	845	98.3	15	1.7
Infancy Completer	734	660	89.9	63	8.6
Toddlerhood Completer	588	488	83.0	41	7.0
Programme Completer	588	488	83.0	41	7.0

*For active stage completers

**% of those enrolled who complete the programme for the specified stage of the programme.

***There were two returners in the infancy phase and four returners in the toddlerhood phase.

**** For those ever enrolled who were stage completers.

All sites

- Overall, 98.3% (N= 845 of 860) active clients completed the pregnancy stage, 89.9% (N= 660 of 734) active clients completed the infancy stage and 83% (N= 488 of 588) active clients completed the toddlerhood phase.

Table 84: Total Attrition by Stage for Stage Completers (All sites)

Total attrition by stage for stage completers	N (%) Inactive/Leavers
Attrition during pregnancy	15 (1.7% [15/860])
Attrition during infancy	63 (8.6% [63/734])
Attrition during toddlerhood	41 (7% [41/588])

All sites

Overall, of the 913 clients ever enrolled, attrition by the 31.12.2018 is as follows:

- 15 (1.7%) left during pregnancy (11 were leavers and 4 became inactive);
- 63 (8.6%) left during infancy (29 were leavers and 34 became inactive); and
- 41 (7%) left during toddlerhood (17 were leavers and 27 became inactive, one of these clients became inactive, returned to the programme and then left the programme, 2 further clients became inactive and then left in toddlerhood).

Table 85: Total Attrition by Stage for Stage Completers BHSCT

Total attrition by stage for stage completers	N (%) Inactive/Leavers
Attrition during pregnancy	5 (2.8% [5/179])
Attrition during infancy	12 (8.1% [12/148])
Attrition during toddlerhood	8 (6.8% [8/117])

BHSCT

Overall, of the 183 clients ever enrolled, attrition by the 31.12.2018 is as follows:

- 5 (2.8%) left or became inactive during pregnancy (3 were leavers and 2 became inactive);
- 12 (8.1%) left or became inactive during infancy (6 were leavers and 6 became inactive);
- 8 (6.8%) left or became inactive during toddlerhood (3 were leavers and 5 became inactive).

Table 86: Total Attrition by Stage for Stage Completers NHSCT

Total attrition by stage for stage completers	N (%) Inactive/Leavers
Attrition during pregnancy	1 (1% [1/101])
Attrition during infancy	11 (13.8% [11/80])
Attrition during toddlerhood	5 (9.6% [5/52])

NHSCT

Overall, of the 110 clients ever enrolled, attrition by the 31.12.2018 is as follows:

- 1 (1%) left during pregnancy
- 11 (13.8%) left during infancy (4 were leavers and 7 became inactive).
- 5 (9.6%) left or became inactive during toddlerhood (3 were leavers and 3 became inactive, one client became inactive and then left).

Table 87: Total Attrition by Stage for Stage Completers SEHSCT

Total attrition by stage for stage completers	N (%) Inactive/Leavers
Attrition during pregnancy	2 (1.8% [2/114])
Attrition during infancy	6 (6.3% [6/95])
Attrition during toddlerhood	5 (7.9% [5/63])

SEHSCT

Overall, of the 138 clients ever enrolled, attrition by the 31.12.2018 is as follows:

- 2 (1.8%) left during pregnancy.
- 6 (6.3%) left during infancy.
- 5 (7.9%) left during toddlerhood.

Table 88: Total Attrition by Stage for Stage Completers SHSCT

Total attrition by stage for stage completers	N (%) Inactive/Leavers
Attrition during pregnancy	3 (1.6% [3/185])
Attrition during infancy	7 (4.5% [7/157])
Attrition during toddlerhood	11 (7.8% [11/141])

SHSCT

Overall, of the 193 clients ever enrolled, attrition by the 31.12.2018 is as follows:

- 3 (1.6%) left or became inactive during pregnancy (2 were leavers and 1 became inactive);
- 7 (4.5%) left or became inactive during infancy (4 were leavers and 3 became inactive); and
- 11 (7.8%) left or became inactive during toddlerhood (2 were leavers and 10 became inactive, one client became inactive and then left).

Table 89: Total Attrition by Stage for Stage Completers WHSCT

Total attrition by stage for stage completers	N (%) Inactive/Leavers
Attrition during pregnancy	4 (1.4% [4/281])
Attrition during infancy	27 (10.6% [27/254])
Attrition during toddlerhood	12 (5.6% [12/215])

WHSCT

Overall, of the 289 clients ever enrolled, attrition by the 31.12.2018 is as follows:

- 4 (1.4%) left or became inactive during pregnancy (3 were leavers and 1 became inactive).
- 27 (10.6%) left or became inactive during infancy (9 were leavers and 18 became inactive).
- 12 (5.6%) left or became inactive during toddlerhood (4 were leavers and 9 became inactive, one client became inactive and then left).

Retention Pregnancy

Table 90: Client Retention by Site and by Year – Pregnancy Stage

Site	Year Enrolled	Completer	Active Completer		Inactive/Leaver*		Leaver		Inactive	
		N	N	%	N	%	N	%	N	%
BHSCT	2013	43	41	95.3	2	4.7	2	4.7	0	0.0
	2014	43	40	93.0	3	7.0	1	2.3	2	4.7
	2015	8	8	100.0	0	0.0	0	0.0	0	0.0
	2016	43	43	100.0	0	0.0	0	0.0	0	0.0
	2017	25	25	100.0	0	0.0	0	0.0	0	0.0
	2018	17	17	100.0	0	0.0	0	0.0	0	0.0
	Total		179	174	97.2	5	2.8	3	1.7	2
NHSCT	2015	19	19	100.0	0	0.0	0	0.0	0	0.0
	2016	41	41	100.0	0	0.0	0	0.0	0	0.0
	2017	28	27	96.4	1	3.6	1	3.6	0	0.0
	2018	13	13	100.0	0	0.0	0	0.0	0	0.0
	Total		101	100	99.0	1	1.0	1	1.0	0
SEHSCT	2014	1	1	100.0	0	0.0	0	0.0	0	0.0
	2015	21	20	95.2	1	4.8	1	4.8	0	0.0
	2016	58	57	98.3	1	1.7	1	1.7	0	0.0
	2017	18	18	100.0	0	0.0	0	0.0	0	0.0
	2018	16	16	100.0	0	0.0	0	0.0	0	0.0
	Total		114	112	98.2	2	1.8	2	1.8	0
SHSCT	2012	13	13	100.0	0	0.0	0	0.0	0	0.0
	2013	72	71	98.6	1	1.4	1	1.4	0	0.0
	2014	23	23	100.0	0	0.0	0	0.0	0	0.0
	2015	5	4	80.0	1	20.0	0	0.0	1	20.0
	2016	32	31	96.9	1	3.1	1	3.1	0	0.0

	2017	20	20	100.0	0	0.0	0	0.0	0	0.0
	2018	20	20	100.0	0	0.0	0	0.0	0	0.0
	Total	185	182	98.4	3	1.6	2	1.1	1	0.5
WHSCT	2010	7	7	100.0	0	0.0	0	0.0	0	0.0
	2011	76	76	100.0	0	0.0	0	0.0	0	0.0
	2012	22	22	100.0	0	0.0	0	0.0	0	0.0
	2013	23	23	100.0	0	0.0	0	0.0	0	0.0
	2014	49	47	95.9	2	4.1	1	2.0	1	2.0
	2015	27	27	100.0	0	0.0	0	0.0	0	0.0
	2016	28	28	100.0	0	0.0	0	0.0	0	0.0
	2017	41	39	95.1	2	4.9	2	4.9	0	0.0
	2018	8	8	100.0	0	0.0	0	0.0	0	0.0
	Total	281	277	98.6	4	1.4	3	1.1	1	0.4
All site total		860	845	98.3	15	1.7	11	1.3	4	0.5

*Those who became inactive/left during the stage.

Retention Infancy

Table 91: Client Retention by Site and by Year – Infancy Stage

Site	Year Enrolled	Completer	Active Completer		Inactive/Leaver*		Leaver		Inactive	
		N	N	%	N	%	N	%	N	%
BHSCT	2013	43	37	86.0	4	9.3	2	4.7	2	4.7
	2014	43	39	90.7	2	4.7	2	4.7	0	0.0
	2015	8	8	100.0	0	0.0	0	0.0	0	0.0
	2016	43	37	86.0	6	14.0	2	4.7	4	9.3
	2017	11	11	100.0	0	0.0	0	0.0	0	0.0
	Total	148	132	89.2	12	8.1	6	4.1	6	4.1
NHSCT	2015	19	17	89.5	2	10.5	1	5.3	1	5.3
	2016	41	34	82.9	7	17.1	3	7.3	4	9.8
	2017	20	18	90.0	2	10.0	0	0.0	2	10.0
		Total	80	69	86.3	11	13.8	4	5.0	7
SEHSCT	2014	1	1	100.0	0	0.0	0	0.0	0	0.0
	2015	21	19	90.5	1	4.8	1	4.8	0	0.0
	2016	58	52	89.7	5	8.6	5	8.6	0	0.0
	2017	15	15	100.0	0	0.0	0	0.0	0	0.0
		Total	95	87	91.6	6	6.3	6	6.3	0
SHSCT	2012	13	12	92.3	1	7.7	0	0.0	1	7.7
	2013	72	68	94.4	3	4.2	1	1.4	2	2.8
	2014	23	22	95.7	1	4.3	1	4.3	0	0.0
	2015	5	4	80.0	0	0.0	0	0.0	0	0.0
	2016	32	29	90.6	2	6.3	2	6.3	0	0.0
	2017	12	12	100.0	0	0.0	0	0.0	0	0.0
		Total	157	147	93.6	7	4.5	4	2.5	3
WHSCT	2010	7	5	71.4	2	28.6	0	0.0	2	28.6
	2011	76	69	90.8	7	9.2	1	1.3	6	7.9
	2012	22	20	90.9	2	9.1	1	4.5	1	4.5
	2013	23	21	91.3	2	8.7	1	4.3	1	4.3
	2014	49	44	89.8	4	8.2	2	4.1	2	4.1
	2015	27	25	92.6	2	7.4	1	3.7	1	3.7

	2016	28	22	78.6	6	21.4	3	10.7	3	10.7
	2017	22	19	86.4	2	9.1	0	0.0	2	9.1
	Total	254	225	88.6	27	10.6	9	3.5	18	7.1
All Site total		734	660	89.9	63	8.6	29	4.0	34	4.6

*Those who became inactive/left during the stage.

Retention Toddlerhood

Table 92: Client Retention by Site and by Year – Toddlerhood Stage

Site	Year Enrolled	Completer	Active Completer		Inactive/Leaver*		Leaver		Inactive	
		N	N	%	N	%	N	%	N	%
BHSCT	2013	43	35	81.4	2	4.7	0	0.0	2	4.7
	2014	43	36	83.7	3	7.0	1	2.3	2	4.7
	2015	8	7	87.5	1	12.5	1	12.5	0	0.0
	2016	23	17	73.9	2	8.7	1	4.3	1	4.3
	Total	117	95	81.2	8	6.8	3	2.6	5	4.3
NHSCT	2015	19	16	84.2	1	5.3	1	5.3	0	0.0
	2016	33	22	66.7	4	12.1	2	6.1	3	9.1
	Total	52	38	73.1	5	9.6	3	5.8	3	5.8
SEHSCT	2014	1	0	0.0	1	100.0	1	100.0	0	0.0
	2015	21	17	81.0	2	9.5	2	9.5	0	0.0
	2016	41	34	82.9	2	4.9	2	4.9	0	0.0
	Total	63	51	81.0	5	7.9	5	7.9	0	0.0
SHSCT	2012	13	12	92.3	1	7.7	0	0.0	1	7.7
	2013	72	62	86.1	6	8.3	1	1.4	5	6.9
	2014	23	21	91.3	2	8.7	0	0.0	2	8.7
	2015	5	4	80.0	0	0.0	0	0.0	0	0.0
	2016	28	23	82.1	2	7.1	1	3.6	2	7.1
	Total	141	122	86.5	11	7.8	2	1.4	10	7.1
WHSCT	2010	7	5	71.4	0	0.0	0	0.0	0	0.0
	2011	76	64	84.2	6	7.9	1	1.3	6	7.9
	2012	22	19	86.4	1	4.5	1	4.5	0	0.0
	2013	23	20	87.0	1	4.3	0	0.0	1	4.3
	2014	49	42	85.7	2	4.1	1	2.0	1	2.0

	2015	27	24	88.9	1	3.7	1	3.7	0	0.0
	2016	11	8	72.7	1	9.1	0	0.0	1	9.1
	Total	215	182	84.7	12	5.6	4	1.9	9	4.2
All site total		588	488	83.0	41	7.0	17	2.9	27	4.6

*Those who became inactive/left during the stage.

Overall Retention

- 98.3% (845 active clients of 860 clients scheduled to complete the stage) clients completed the pregnancy stage.
- 89.9% (660 active clients of 734 clients scheduled to complete the stage) clients completed the infancy stage.
- 83% (488 active clients of 588 clients scheduled to complete the stage) clients completed the toddlerhood stage.

Short-Term Outcomes

Hospital Anxiety and Depression Scale (HADs)

Table 93: HADS* Anxiety – All sites

Anxiety		BHSCT	NHSCT	SEHSCT	SHSCT	WHSCT	All sites
Number of active clients reaching 36 weeks pregnancy	N	177	102	116	187	281	863
Clients with HADs data recorded in pregnancy	N	158	80	109	160	229	736
	%	89.3	78.4	94.0	85.6	81.5	85.3
Unlikely presence of anxiety pregnancy	N	117	64	90	139	177	587
	%	74.1	80.0	82.6	86.9	77.3	79.8
Borderline presence of anxiety pregnancy	N	24	12	11	11	30	88
	%	15.2	15.0	10.1	6.9	13.1	12.0
Probable presence of anxiety pregnancy	N	11	2	4	9	18	44
	%	7.0	2.5	3.7	5.6	7.9	6.0
High likelihood of presence of anxiety pregnancy	N	6	2	4	-	4	16
	%	3.8	2.5	3.7	-	1.7	2.2

Table 94: HADS Depression – All sites

Depression		BHSCT	NHSCT	SEHSCT	SHSCT	WHSCT	All sites
Number of active clients reaching 36 weeks pregnancy	N	177	102	116	187	281	863
Clients with HADs data recorded in pregnancy	N	158	80	109	158	229	734
	%	89.3	78.4	94.0	85.4	81.5	85.1
Unlikely presence of depression pregnancy	N	142	67	102	149	204	664
	%	89.9	83.8	93.6	94.3	89.1	90.5
Borderline presence of depression pregnancy	N	10	6	4	7	20	47
	%	6.3	7.5	3.7	4.4	8.7	6.4
Probable presence of depression pregnancy	N	5	7	3	1	4	20
	%	3.2	8.8	2.8	0.6	1.7	2.7
High likelihood of presence of depression pregnancy	N	1	-	-	-	1	2

	%	0.6	-	-	-	0.4	0.3
--	---	-----	---	---	---	-----	-----

* The Hospital Anxiety and Depression Scale (HADS) measures anxiety and depression [Zigmond A.S. & Snaith R.P. (1983).The Hospital Anxiety and Depression Scale. Acta Psychiatr Scand; 67:361–370]. **2 clients recorded an answer for anxiety but answer was missing for depression

Overall

Anxiety

- Overall, anxiety (HADS) scores were recorded for 85.3% (N=736) clients in pregnancy.
- Of these, 79.8% (N=587) were found to have an unlikely presence of anxiety, 12% (N=88) were found to have borderline presence of anxiety, 6% (N=44) were found to have probable presence of anxiety and 2.2% (N=16) were found to have a high likelihood of anxiety in pregnancy.

Depression

- Overall, depression (HADS) scores were recorded for 85.1% (N=734) clients in pregnancy.
- Of these, 90.5% (N=664) were found to have an unlikely presence of depression, 6.4% (N=47) were found to have borderline presence of depression, 2.7% (N=20) were found to have probable presence of depression and 0.3% (N=2) clients were found to have a high likelihood of depression in pregnancy.

BHSCT

Anxiety

- Overall, anxiety (HADS) scores were recorded for 89.3% (N=158) clients in pregnancy.
- Of these, 74.1% (N=117) were found to have an unlikely presence of anxiety, 15.2% (N=24) were found to have borderline presence of anxiety, 7% (N=11) were found to have probable presence of anxiety and 3.8% (N=6) were found to have a high likelihood of anxiety in pregnancy.

Depression

- Overall, depression (HADS) scores were recorded for 89.3% (N=158) clients in pregnancy.
- Of these, 89.9% (N=142) were found to have an unlikely presence of depression, 6.3% (N=10) were found to have borderline presence of depression, 3.2% (N=5) were found to have probable presence of depression and 0.6% (N=1) clients were found to have a high likelihood of depression in pregnancy.

NHSCT

Anxiety

- Overall, anxiety (HADs) scores were recorded for 78.4% (N=80) clients in pregnancy.
- Of these, 80% (N=64) were found to have an unlikely presence of depression, 15% (N=12) were found to have borderline presence of depression, 2.5% (N=2) were found to have probable presence of depression and 2.5% (N=2) clients were found to have a high likelihood of depression in pregnancy.

Depression

- Overall, depression (HADs) scores were recorded for 78.4% (N=80) clients in pregnancy.
- Of these, 83.8% (N=67) were found to have an unlikely presence of depression, 7.5% (N=6) were found to have borderline presence of depression, 8.8% (N=7) were found to have probable presence of depression and no clients were found to have a high likelihood of depression in pregnancy.

SEHSCT

Anxiety

- Overall, anxiety (HADs) scores were recorded for 94% (N=109) clients in pregnancy.
- Of these, 82.6% (N=90) were found to have an unlikely presence of anxiety, 10.1% (N=11) were found to have borderline presence of anxiety, 3.7% (N=4) were found to have probable presence of anxiety and 3.7% (N=4) were found to have a high likelihood of anxiety in pregnancy.

Depression

- Overall, depression (HADs) scores were recorded for 94% (N=109) clients in pregnancy.
- Of these, 93.6% (N=102) were found to have an unlikely presence of depression, 3.7% (N=4) were found to have borderline presence of depression, 2.8% (N=3) were found to have probable presence of depression and no clients were found to have a high likelihood of depression in pregnancy.

SHSCT

Anxiety

- Overall, anxiety (HADs) scores were recorded for 85.6% (N=160) clients in pregnancy.
- Of these, 86.9% (N=139) were found to have an unlikely presence of anxiety, 6.9% (N=11) were found to have borderline presence of anxiety, 5.6% (N=9) were found to have probable presence of anxiety and no clients were found to have a high likelihood of anxiety in pregnancy.

Depression

- Overall, depression (HADs) scores were recorded for 85.4% (N=158) clients in pregnancy.
- Of these, 94.3% (N=149) were found to have an unlikely presence of depression, 4.4% (N=7) were found to have borderline presence of depression, 0.6% (N=1) were found to have probable presence of depression and no clients were found to have a high likelihood of depression in pregnancy.

WHSCT

Anxiety

- Overall, anxiety (HADs) scores were recorded for 81.5% (N=229) clients in pregnancy.
- Of these, 77.3% (N=177) were found to have an unlikely presence of anxiety, 13.1% (N=30) were found to have borderline presence of anxiety, 7.9% (N=18) were found to have probable presence of anxiety and 1.7% (N=4) were found to have a high likelihood of anxiety in pregnancy.

Depression

- Overall, depression (HADs) scores were recorded for 81.5% (N=229) clients in pregnancy.
- Of these, 89.1% (N=204) were found to have an unlikely presence of depression, 8.7% (N=20) were found to have borderline presence of depression, 1.7% (N=4) were found to have probable presence of depression and 0.4% (N=1) clients were found to have a high likelihood of depression in pregnancy.

Breastfeeding

Table 95: Breastfeeding – All sites

	Total
Total number enrolled	913
Initiation	
Total number of active clients with infants	827
Number (%) of clients with breastfeeding initiation status recorded	785 (94.9%)
Number (%) of clients initiating breastfeeding	345 (43.9%)
6 weeks Infancy	
Total number of active clients with infants reaching 6 weeks infancy	804
Number (%) of clients with status recorded at 6 weeks infancy	745 (92.7%)
Number (%) of clients breastfeeding at 6 weeks infancy	60 (8.1%)
6 months Infancy	
Total number of active clients with infants reaching 6 months infancy	744
Number (%) of clients with status recorded at 6 months infancy	628 (84.4%)
Number (%) of clients breastfeeding at 6 months infancy	19 (3%)
Exclusive breastfeeding	
Number (%) of clients with exclusive breastfeeding status recorded at 6 weeks or 6 months	763 (94.9%)
Average duration clients exclusively breastfed their infant (weeks)	3.9 weeks

All sites

- Breastfeeding initiation was recorded for 94.9% (N=785) of clients with infants. Of these 43.9% (N=345) reported initiating breastfeeding.
- Breastfeeding status was recorded for 92.7% (N=745) of clients who had reached 6 weeks infancy. Of these, 8.1% (N=60) reported that they were still breastfeeding at 6 weeks.
- Breastfeeding status was recorded for 84.4% (N=628) of clients who had reached 6 months infancy. Of these, 3% (N=19) reported that they were still breastfeeding at 6 months.
- Exclusive breastfeeding status at either 6 weeks or 6 months was recorded for 94.9% (N=763) clients. For these clients, the average duration that clients exclusively breastfed their infant was 3.9 weeks.

Table 96: Breastfeeding – BHSCT and all sites

	BHSCT	Total
Total number enrolled	183	913
Initiation		
Total number of active clients with infants	169	827
Number (%) of clients with breastfeeding initiation status recorded	159 (94.1%)	785 (94.9%)
Number (%) of clients initiating breastfeeding	53 (33.3%)	345 (43.9%)
6 weeks Infancy		
Total number of active clients with infants reaching 6 weeks infancy	165	804
Number (%) of clients with status recorded at 6 weeks infancy	155 (93.9%)	745 (92.7%)
Number (%) of clients breastfeeding at 6 weeks infancy	17 (11%)	60 (8.1%)
6 months Infancy		
Total number of active clients with infants reaching 6 months infancy	153	744
Number (%) of clients with status recorded at 6 months infancy	137 (89.5%)	628 (84.4%)
Number (%) of clients breastfeeding at 6 months infancy	4 (2.9%)	19 (3%)
Exclusive breastfeeding		
Number (%) of clients with exclusive breastfeeding status recorded at 6 weeks or 6 months	157 (95.2%)	763 (94.9%)
Average duration clients exclusively breastfed their infant (weeks)	4.9 weeks	3.9 weeks

BHSCT

- Breastfeeding initiation was recorded for 94.1% (N=159) of clients with infants. Of these 33.3% (N=53) reported initiating breastfeeding.
- Breastfeeding status was recorded for 93.9% (N=155) of clients who had reached 6 weeks infancy. Of these, 11% (N=17) reported that they were still breastfeeding at 6 weeks.
- Breastfeeding status was recorded for 89.5% (N=137) of clients who had reached 6 months infancy. Of these, 2.9% (N=4) reported that they were still breastfeeding at 6 months.
- Exclusive breastfeeding status at either 6 weeks or 6 months was recorded for 95.2% (N=157) clients. For these clients, the average duration that clients exclusively breastfed their infant was 4.9 weeks.

Table 97: Breastfeeding – NHSCT and all sites

	NHSCT	Total
Total number enrolled	110	913
Initiation		
Total number of active clients with infants	96	827
Number (%) of clients with breastfeeding initiation status recorded	91 (94.8%)	785 (94.9%)
Number (%) of clients initiating breastfeeding	39 (42.9%)	345 (43.9%)
6 weeks Infancy		
Total number of active clients with infants reaching 6 weeks infancy	94	804
Number (%) of clients with status recorded at 6 weeks infancy	83 (88.3%)	745 (92.7%)
Number (%) of clients breastfeeding at 6 weeks infancy	6 (7.2%)	60 (8.1%)
6 months Infancy		
Total number of active clients with infants reaching 6 months infancy	86	744
Number (%) of clients with status recorded at 6 months infancy	56 (65.1%)	628 (84.4%)
Number (%) of clients breastfeeding at 6 months infancy	1 (1.8%)	19 (3%)
Exclusive breastfeeding		
Number (%) of clients with exclusive breastfeeding status recorded at 6 weeks or 6 months	83 (88.3%)	763 (94.9%)
Average duration clients exclusively breastfed their infant (weeks)	3.1 weeks	3.9 weeks

NHSCT

- Breastfeeding initiation was recorded for 94.8% (N=91) of clients with infants. Of these 42.9% (N=39) reported initiating breastfeeding.
- Breastfeeding status was recorded for 88.3% (N=83) of clients who had reached 6 weeks infancy. Of these, 7.2% (N=6) reported that they were still breastfeeding at 6 weeks.
- Breastfeeding status was recorded for 65.1% (N=56) of clients who had reached 6 months infancy. Of these, 1.8% (N=1) reported that they were still breastfeeding at 6 months.
- Exclusive breastfeeding status at either 6 weeks or 6 months was recorded for 88.3% (N=83) clients. For these clients, the average duration that clients exclusively breastfed their infant was 3.1 weeks.

Table 98: Breastfeeding – SEHSCT and all sites

	SEHSCT	Total
Total number enrolled	138	913
Initiation		
Total number of active clients with infants	110	827
Number (%) of clients with breastfeeding initiation status recorded	108 (98.2%)	785 (94.9%)
Number (%) of clients initiating breastfeeding	64 (59.3%)	345 (43.9%)
6 weeks Infancy		
Total number of active clients with infants reaching 6 weeks infancy	105	804
Number (%) of clients with status recorded at 6 weeks infancy	103 (98.1%)	745 (92.7%)
Number (%) of clients breastfeeding at 6 weeks infancy	12 (11.7%)	60 (8.1%)
6 months Infancy		
Total number of active clients with infants reaching 6 months infancy	92	744
Number (%) of clients with status recorded at 6 months infancy	87 (94.6%)	628 (84.4%)
Number (%) of clients breastfeeding at 6 months infancy	4 (4.6%)	19 (3%)
Exclusive breastfeeding		
Number (%) of clients with exclusive breastfeeding status recorded at 6 weeks or 6 months	104 (99%)	763 (94.9%)
Average duration clients exclusively breastfed their infant (weeks)	4.5 weeks	3.9 weeks

SEHSCT

- Breastfeeding initiation was recorded for 98.2% (N=108) of clients with infants. Of these 59.3% (N=64) reported initiating breastfeeding.
- Breastfeeding status was recorded for 98.1% (N=103) of clients who had reached 6 weeks infancy. Of these, 11.7% (N=12) reported that they were still breastfeeding at 6 weeks.
- Breastfeeding status was recorded for 94.6% (N=87) of clients who had reached 6 months infancy. Of these, 4.6% (N=4) reported that they were still breastfeeding at 6 months.
- Exclusive breastfeeding status at either 6 weeks or 6 months was recorded for 99% (N=104) clients. For these clients, the average duration that clients exclusively breastfed their infant was 4.5 weeks.

Table 99: Breastfeeding – SHSCT and all sites

	SHSCT	Total
Total number enrolled	193	913
Initiation		
Total number of active clients with infants	180	827
Number (%) of clients with breastfeeding initiation status recorded	172 (95.6%)	785 (94.9%)
Number (%) of clients initiating breastfeeding	80 (46.5%)	345 (43.9%)
6 weeks Infancy		
Total number of active clients with infants reaching 6 weeks infancy	175	804
Number (%) of clients with status recorded at 6 weeks infancy	167 (95.4%)	745 (92.7%)
Number (%) of clients breastfeeding at 6 weeks infancy	16 (9.6%)	60 (8.1%)
6 months Infancy		
Total number of active clients with infants reaching 6 months infancy	161	744
Number (%) of clients with status recorded at 6 months infancy	143 (88.8%)	628 (84.4%)
Number (%) of clients breastfeeding at 6 months infancy	7 (4.9%)	19 (3%)
Exclusive breastfeeding		
Number (%) of clients with exclusive breastfeeding status recorded at 6 weeks or 6 months	170 (97.1%)	763 (94.9%)
Average duration clients exclusively breastfed their infant (weeks)	4.4 weeks	3.9 weeks

SHSCT

- Breastfeeding initiation was recorded for 95.6% (N=172) of clients with infants. Of these 46.5% (N=80) reported initiating breastfeeding.
- Breastfeeding status was recorded for 95.4% (N=167) of clients who had reached 6 weeks infancy. Of these, 9.6% (N=16) reported that they were still breastfeeding at 6 weeks.
- Breastfeeding status was recorded for 88.8% (N=143) of clients who had reached 6 months infancy. Of these, 4.9% (N=7) reported that they were still breastfeeding at 6 months.
- Exclusive breastfeeding status at either 6 weeks or 6 months was recorded for 97.1% (N=170) clients. For these clients, the average duration that clients exclusively breastfed their infant was 4.4 weeks.

Table 100: Breastfeeding – WHSCT and all sites

	WHSCT	Total
Total number enrolled	289	913
Initiation		
Total number of active clients with infants	272	827
Number (%) of clients with breastfeeding initiation status recorded	255 (93.8%)	785 (94.9%)
Number (%) of clients initiating breastfeeding	109 (42.7%)	345 (43.9%)
6 weeks Infancy		
Total number of active clients with infants reaching 6 weeks infancy	265	804
Number (%) of clients with status recorded at 6 weeks infancy	237 (89.4%)	745 (92.7%)
Number (%) of clients breastfeeding at 6 weeks infancy	9 (3.8%)	60 (8.1%)
6 months Infancy		
Total number of active clients with infants reaching 6 months infancy	252	744
Number (%) of clients with status recorded at 6 months infancy	205 (81.3%)	628 (84.4%)
Number (%) of clients breastfeeding at 6 months infancy	3 (1.5%)	19 (3%)
Exclusive breastfeeding		
Number (%) of clients with exclusive breastfeeding status recorded at 6 weeks or 6 months	249 (94%)	763 (94.9%)
Average duration clients exclusively breastfed their infant (weeks)	3 weeks	3.9 weeks

WHSCT

- Breastfeeding initiation was recorded for 93.8% (N=255) of clients with infants. Of these 42.7% (N=109) reported initiating breastfeeding.
- Breastfeeding status was recorded for 89.4% (N=237) of clients who had reached 6 weeks infancy. Of these, 3.8% (N=9) reported that they were still breastfeeding at 6 weeks.
- Breastfeeding status was recorded for 81.3% (N=205) of clients who had reached 6 months infancy. Of these, 1.5% (N=3) reported that they were still breastfeeding at 6 months.
- Exclusive breastfeeding status at either 6 weeks or 6 months was recorded for 94% (N=249) clients. For these clients, the average duration that clients exclusively breastfed their infant was 3 weeks.

Tobacco Use

Table 101: Change in self-reported smoking during pregnancy (intake to 36 weeks) – All sites

		BHSCT	NHSCT	SEHSCT	SHSCT	WHSCT	Total
Number of clients enrolled	N	183	110	138	193	289	913
Clients with information on smoking during pregnancy recorded at intake	N	173	99	135	180	267	854
	%	94.5	90.0	97.8	93.3	92.4	93.5
Clients who smoked at any time during the pregnancy (intake)	N	78	47	52	69	103	349
	%	45.1	47.5	38.5	38.3	38.6	40.9
Clients who smoked in the last 48h recorded at intake	N	173	99	135	180	267	854
	%	94.5	90.0	97.8	93.3	92.4	93.5
Clients who smoked in the last 48h at intake	N	53	24	33	45	79	234
	%	30.6	24.2	24.4	25.0	29.6	27.4
Mean number of cigarettes smoked in last 48h at intake	N	10.7	9.8	8.2	9.1	11.2	10.1
Mean number of cigarettes smoked in last 48h at intake among women who smoked more than 5 cigarettes per day	N	15.5	17.8	16.8	17.3	15.7	16.1
Number of active clients reaching 36 weeks' gestation	N	177	102	116	187	281	863
Clients with information on smoking during pregnancy recorded at 36weeks	N	158	84	109	160	229	740
	%	89.3	82.4	94.0	85.6	81.5	85.7
Clients who smoked at any time during the pregnancy (36 weeks)	N	69	39	43	61	81	293
	%	43.7	46.4	39.4	38.1	35.4	39.6
Clients who smoked in the last 48h recorded at 36 weeks	N	158	84	109	160	229	740
	%	89.3	82.4	94.0	85.6	81.5	85.7
Clients who smoked in the last 48h at 36 weeks	N	42	17	24	34	51	168
	%	26.6	20.2	22.0	21.3	22.3	22.7
Mean number of cigarettes smoked in last 48h at intake	N	9.3	8.0	7.9	6.9	11.2	9.1
Mean number of cigarettes smoked in last 48h at intake among women who smoked more than 5 cigarettes per day	N	16.7	13.3	12.5	17.0	16.3	15.7
Number of active clients at 36 weeks pregnancy with information on number of cigarettes smoked in the last 48h recorded at both intake and 36 weeks*	N	158	82	109	157	221	727
% change in number of active clients smoking in the last 48h recorded at both intake and 36 weeks*	%	-4.4	-2.4	-3.7	-5.1	-6.8	-5.0

% change in proportion of clients smoking in last 48h between intake and 36 weeks*	%	-14.3	-10.5	-14.3	-19.5	-23.8	-18.0
% of clients smoking fewer cigarettes in last 48h at 36weeks pregnancy*	%	19.0	15.9	12.8	18.5	19.5	17.7
% of clients smoking fewer cigarettes in last 48h at 36 weeks pregnancy than at intake among women who smoked more than 5 cigarettes per day at intake*	%	66.7	87.5	66.7	78.6	79.5	75.3
Clients who smoked more at 36weeks than at intake*	N	17	3	10	7	16	53
	%	10.8	3.7	9.2	4.5	7.2	7.3
Number of smokers at 36 weeks who denied at intake*	N	6	2	2	1	1	12
	%	3.8	2.4	1.8	0.6	0.5	1.7

*These figures are for the subgroup of clients who had data recorded at both intake and 36 weeks.

All sites

Intake

- Information on smoking during pregnancy was recorded for 93.5% (N= 854) of clients at intake. Of these, 40.9 (N=349) reported that they had smoked at some time during pregnancy at intake.
- Information on smoking during the last 48 hours was recorded for 93.5% (N= 854) of clients at intake. Of these, 27.4% (N=234) reported that they had smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 10.1 for those who smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 16.1 for those who smoked more than 5 cigarettes per day during the last 48 hours at intake.

36 weeks pregnancy

- Information on smoking during pregnancy was recorded for 85.7% (N= 740) of clients at 36 weeks pregnancy. Of these, 39.6% (N=293) reported that they had smoked at some time during the pregnancy at 36 weeks pregnancy
- Information on smoking during the last 48 hours was recorded for 85.7% (N= 740) of clients at 36 weeks pregnancy. Of these, 22.7% (N=168) reported that they had smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 9.1 for those who smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 15.7 for those who smoked more than 5 cigarettes per day during the last 48 hours at 36 weeks pregnancy.

Intake and 36 weeks

- 727 clients had information on smoking status recorded at both intake and 36 weeks pregnancy. For these clients:
 - a -5% change in smoking in the last 48 hours was achieved between intake and 36 weeks.
 - a -18% change in the proportion of clients smoking in in the last 48 hours was achieved between intake and 36 weeks.
 - 17.7% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks.
 - 75.3% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks for clients who smoked more than 5 cigarettes per day at intake.
 - 7.3% (N=53) clients smoked more at 36 weeks than at intake.
 - 1.7% (N=12) clients smoked at 36 weeks who denied smoking at intake.

BHSCT

Intake

- Information on smoking during pregnancy was recorded for 94.5% (N=173) of clients at intake. Of these, 45.1% (N=78) reported that they had smoked at some time during pregnancy at intake.
- Information on smoking during the last 48 hours was recorded for 94.5% (N=173) of clients at intake. Of these, 30.6% (N=53) reported that they had smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 10.7 for those who smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 15.5 for those who smoked more than 5 cigarettes per day during the last 48 hours at intake.

36 weeks pregnancy

- Information on smoking during pregnancy was recorded for 89.3% (N=158) of clients at 36 weeks pregnancy. Of these, 43.7% (N=69) reported that they had smoked at some time during the pregnancy at 36 weeks pregnancy
- Information on smoking during the last 48 hours was recorded for 89.3% (N=158) of clients at 36 weeks pregnancy. Of these, 26.6% (N=42) reported that they had smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 9.3 for those who smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 16.7 for those who smoked more than 5 cigarettes per day during the last 48 hours at 36 weeks pregnancy.

Intake and 36 weeks

- 158 clients had information on smoking status recorded at both intake and 36 weeks pregnancy. For these clients:
 - a -4.4 % change in smoking in the last 48 hours was achieved between intake and 36 weeks.
 - a -14.3 % change in the proportion of clients smoking in in the last 48 hours was achieved between intake and 36 weeks.
 - 19% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks.
 - 66.7% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks for clients who smoked more than 5 cigarettes per day at intake.
 - 10.8% (N=17) clients smoked more at 36 weeks than at intake.
 - 3.8% (N=6) clients smoked at 36 weeks who denied smoking at intake

NHSCT

Intake

- Information on smoking during pregnancy was recorded for 90% (N=99) of clients at intake. Of these, 47.5% (N=47) reported that they had smoked at some time during pregnancy at intake.
- Information on smoking during the last 48 hours was recorded for 90% (N=99) of clients at intake. Of these, 24.2% (N=24) reported that they had smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 9.8 for those who smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 17.8 for those who smoked more than 5 cigarettes per day during the last 48 hours at intake.

36 weeks pregnancy

- Information on smoking during pregnancy was recorded for 82.4% (N=84) of clients at 36 weeks pregnancy. Of these, 46.4% (N=39) reported that they had smoked at some time during the pregnancy at 36 weeks pregnancy
- Information on smoking during the last 48 hours was recorded for 82.4% (N=84) of clients at 36 weeks pregnancy. Of these, 20.2% (N=17) reported that they had smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 8 for those who smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 13.3 for those who smoked more than 5 cigarettes per day during the last 48 hours at 36 weeks pregnancy.

Intake and 36 weeks

- 82 clients had information on smoking status recorded at both intake and 36 weeks pregnancy. For these clients:
 - a –2.4% change in smoking in the last 48 hours was achieved between intake and 36 weeks.
 - a –10.5% change in the proportion of clients smoking in in the last 48 hours was achieved between intake and 36 weeks.
 - 15.9% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks.
 - 87.5% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks for clients who smoked more than 5 cigarettes per day at intake.
 - 3.7% (N=3) clients smoked more at 36 weeks than at intake.
 - 2.4% (N=2) clients smoked at 36 weeks who denied smoking at intake

SEHSCT

Intake

- Information on smoking during pregnancy was recorded for 97.8% (N=135) of clients at intake. Of these, 38.5% (N=52) reported that they had smoked at some time during pregnancy at intake.
- Information on smoking during the last 48 hours was recorded for 97.8% (N=135) of clients at intake. Of these, 24.4% (N=33) reported that they had smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 8.2 for those who smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 16.8 for those who smoked more than 5 cigarettes per day during the last 48 hours at intake.

36 weeks pregnancy

- Information on smoking during pregnancy was recorded for 94% (N=109) of clients at 36 weeks pregnancy. Of these, 39.4% (N=43) reported that they had smoked at some time during the pregnancy at 36 weeks pregnancy
- Information on smoking during the last 48 hours was recorded for 94% (N=109) of clients at 36 weeks pregnancy. Of these, 22% (N=24) reported that they had smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 7.9 for those who smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 12.5 for those who smoked more than 5 cigarettes per day during the last 48 hours at 36 weeks pregnancy.

Intake and 36 weeks

- 109 clients had information on smoking status recorded at both intake and 36 weeks pregnancy. For these clients:
 - a -3.7 % change in smoking in the last 48 hours was achieved between intake and 36 weeks.
 - a -14.3 % change in the proportion of clients smoking in in the last 48 hours was achieved between intake and 36 weeks.
 - 12.8% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks.
 - 66.7% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks for clients who smoked more than 5 cigarettes per day at intake.
 - 9.2% (N=10) clients smoked more at 36 weeks than at intake.
 - 1.8% (N=2) clients smoked at 36 weeks who denied smoking at intake

SHSCT

Intake

- Information on smoking during pregnancy was recorded for 93.3% (N=180) of clients at intake. Of these, 38.3% (N=69) reported that they had smoked at some time during pregnancy at intake.
- Information on smoking during the last 48 hours was recorded for 93.3% (N=180) of clients at intake. Of these, 25% (N=45) reported that they had smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 9.1 for those who smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 17.3 for those who smoked more than 5 cigarettes per day during the last 48 hours at intake.

36 weeks pregnancy

- Information on smoking during pregnancy was recorded for 85.6% (N=160) of clients at 36 weeks pregnancy. Of these, 38.1% (N=61) reported that they had smoked at some time during the pregnancy at 36 weeks pregnancy
- Information on smoking during the last 48 hours was recorded for 85.6% (N=160) of clients at 36 weeks pregnancy. Of these, 21.3% (N=34) reported that they had smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 6.9 for those who smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 17 for those who smoked more than 5 cigarettes per day during the last 48 hours at 36 weeks pregnancy.

Intake and 36 weeks

- 157 clients had information on smoking status recorded at both intake and 36 weeks pregnancy. For these clients:
 - a -5.1 % change in smoking in the last 48 hours was achieved between intake and 36 weeks.
 - a -19.5 % change in the proportion of clients smoking in in the last 48 hours was achieved between intake and 36 weeks.
 - 18.5% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks.
 - 78.6% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks for clients who smoked more than 5 cigarettes per day at intake.
 - 4.5% (N=7) clients smoked more at 36 weeks than at intake.
 - 0.6% (N=1) clients smoked at 36 weeks who denied smoking at intake.

WHST

Intake

- Information on smoking during pregnancy was recorded for 92.4% (N=267) of clients at intake. Of these, 38.6% (N=103) reported that they had smoked at some time during pregnancy at intake.
- Information on smoking during the last 48 hours was recorded for 92.4% (N=267) of clients at intake. Of these, 29.6% (N=79) reported that they had smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 11.2 for those who smoked during the last 48 hours at intake.
- The mean number of cigarettes smoked was 15.7 for those who smoked more than 5 cigarettes per day during the last 48 hours at intake.

36 weeks pregnancy

- Information on smoking during pregnancy was recorded for 81.5% (N=229) of clients at 36 weeks pregnancy. Of these, 35.4% (N=81) reported that they had smoked at some time during the pregnancy at 36 weeks pregnancy
- Information on smoking during the last 48 hours was recorded for 81.5% (N=229) of clients at 36 weeks pregnancy. Of these, 22.3% (N=51) reported that they had smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 11.2 for those who smoked during the last 48 hours at 36 weeks pregnancy.
- The mean number of cigarettes smoked was 16.3 for those who smoked more than 5 cigarettes per day during the last 48 hours at 36 weeks pregnancy.

Intake and 36 weeks

- 221 clients had information on smoking status recorded at both intake and 36 weeks pregnancy. For these clients:
 - a -6.8 % change in smoking in the last 48 hours was achieved between intake and 36 weeks.
 - a -23.8% change in the proportion of clients smoking in in the last 48 hours was achieved between intake and 36 weeks.
 - 19.5% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks.
 - 79.5% of clients smoked fewer cigarettes in the last 48 hours between intake and 36 weeks for clients who smoked more than 5 cigarettes per day at intake.
 - 7.2% (N=16) clients smoked more at 36 weeks than at intake.
 - 0.5% (N=1) client smoked at 36 weeks who denied smoking at intake.

Alcohol Use

Table 102: Change in self-reported alcohol use during pregnancy (intake to 36 weeks) – All sites

		BHSCT	NHSCT	SEHSCT	SHSCT	WHSCT	All Sites
Number of active clients enrolled	N	183	110	138	193	289	913
Clients with information on alcohol use during pregnancy recorded at intake	N	173	99	135	180	267	854
	%	94.5	90.0	97.8	93.3	92.4	93.5
Clients who drank alcohol in last 14 days at intake	N	1	0	3	5	3	12
	%	0.6	0.0	2.2	2.8	1.1	1.4
Clients who drank alcohol excessively in last 14 days at intake	N	0	0	1	0	2	3
	%	0	0	0.7	0	0.7	0.4
Number of active clients reaching 36 weeks gestation	N	177	102	116	187	281	863
Clients with information on alcohol use recorded at 36 weeks pregnancy	N	158	84	109	160	229	740
	%	89.3	82.4	94.0	85.6	81.5	85.7
Clients who drank alcohol in last 14 days at 36 weeks	N	2	0	0	0	4	6
	%	1.3	0	0	0	1.7	0.8
Clients who drank alcohol excessively in last 14 days at 36 weeks	N	0	0	0	0	2	2
	%	0	0	0	0	0.9	0.3
Number of active clients with information on alcohol status recorded at both intake and 36 weeks*	N	158	82	109	157	221	727
% change in proportion of clients drinking alcohol in last 14 days between intake and 36 weeks* [§]	%	100	-	-100	-100	0	-44.4
% change in number of clients drinking alcohol in last 14 days between intake and 36 weeks*	%	0.6	0	-0.9	-2.5	0	-0.6
% change in proportion of clients drinking alcohol excessively in last 14 days between intake and 36 weeks*	%	-	-	-	-	-50	-50
% change in number of clients drinking alcohol excessively in last 14 days between intake and 36 weeks*	%	0	0	0	0	-0.5	-0.1
Number of clients who started drinking between intake and 36 weeks*	N	1	0	0	0	2	3
	%	0.6	-	-	-	0.9	0.4

Number of clients who stopped drinking between intake and 36 weeks*	N	0	0	1	4	2	7
	%	0	0	0.9	2.5	0.9	1.0

*These figures are for the subgroup of clients who had data recorded at both intake and 36 weeks. Please note the small numbers of clients who reported drinking alcohol when interpreting the figures for % change in proportion of clients drinking alcohol.

All sites

Intake

- Information on alcohol use was recorded at intake for 93.5% of clients (N=854) at intake. Of these, 1.4% (N=12) reported drinking alcohol in the last 14 days at intake. 0.4% (N=3) clients reported drinking excessively in the last 14 days at intake.

36 weeks pregnancy

- Information on alcohol use was recorded at 36 weeks pregnancy for 85.7% (N=740) clients. Of these, 0.8% (N=6) reported drinking alcohol in the last 14 days at 36 weeks. 0.3% (N=2) clients reported drinking excessively in the last 14 days at 36 weeks pregnancy.

Intake and 36 weeks

- Information on alcohol use was recorded at both intake and 36 weeks for 727 clients. For these clients, there was:
 - 44.4% change in the proportion of clients drinking alcohol between intake and 36 weeks.
 - 0.6% change in the number of clients drinking alcohol between intake and 36 weeks.
 - 50% change in the proportion of clients drinking alcohol excessively between intake and 36 weeks.
 - 0.1% change in the number of clients drinking alcohol excessively between intake and 36 weeks.
 - 0.4% (N=3) clients started drinking between intake and 36 weeks.
 - 1% (N=7) clients stopped drinking between intake and 36 weeks.
- Please note the small numbers of clients who reported drinking alcohol when interpreting the figures for % change in proportion of clients drinking alcohol.

BHSCT

Intake

- Information on alcohol use was recorded at intake for 94.5% of clients (N=173) at intake. Of these, 0.6% (N=1) reported drinking alcohol in the last 14 days at intake. No clients reported drinking excessively in the last 14 days at intake.

36 weeks pregnancy

- Information on alcohol use was recorded at 36 weeks pregnancy for 89.3% (N=158) clients. Of these, 1.3% (N=2) reported drinking alcohol in the last 14 days at 36 weeks. No clients reported drinking excessively in the last 14 days at 36 weeks pregnancy.

Intake and 36 weeks

- Information on alcohol use was recorded at both intake and 36 weeks for 158 clients. For these clients, there was:
 - 100% change in the proportion of clients drinking alcohol between intake and 36 weeks.
 - 0.6% change in the number of clients drinking alcohol between intake and 36 weeks.
 - 0.6% (N=1) client started drinking between intake and 36 weeks.
- Please note the small numbers of clients who reported drinking alcohol when interpreting the figures for % change in proportion of clients drinking alcohol.

NHSCT

Intake

- Information on alcohol use was recorded at intake for 90% of clients (N=99) at intake. Of these, no clients reported drinking alcohol in the last 14 days at intake. No clients reported drinking excessively in the last 14 days at intake.

36 weeks pregnancy

- Information on alcohol use was recorded at 36 weeks pregnancy for 82.4% (N=84) clients. Of these, no reported drinking alcohol in the last 14 days at 36 weeks. No clients reported drinking excessively in the last 14 days at 36 weeks pregnancy.

Intake and 36 weeks

- Information on alcohol use was recorded at both intake and 36 weeks for 82 clients. For these clients, there was no change (0%) in the number or proportion of clients drinking alcohol between intake and 36 weeks.
- Please note the small numbers of clients who reported drinking alcohol when interpreting the figures for % change in proportion of clients drinking alcohol.

SEHSCT

Intake

- Information on alcohol use was recorded at intake for 97.8% of clients (N=135) at intake. Of these, 2.2% (N=3) reported drinking alcohol in the last 14 days at intake. 0.7% (N=1) clients reported drinking excessively in the last 14 days at intake.

36 weeks pregnancy

- Information on alcohol use was recorded at 36 weeks pregnancy for 94% (N=109) clients. Of these, no clients reported drinking alcohol in the last 14 days at 36 weeks. No clients reported drinking excessively in the last 14 days at 36 weeks pregnancy.

Intake and 36 weeks

- Information on alcohol use was recorded at both intake and 36 weeks for 109 clients. For these clients, there was:
 - -100% change in the proportion of clients drinking alcohol between intake and 36 weeks.
 - -0.9% change in the number of clients drinking alcohol between intake and 36 weeks.
 - No clients started drinking between intake and 36 weeks.
 - 1 (0.9%) client stopped drinking between intake and 36 weeks.
- Please note the small numbers of clients who reported drinking alcohol when interpreting the figures for % change in proportion of clients drinking alcohol.

SHSCT

Intake

- Information on alcohol use was recorded at intake for 93.3% of clients (N=180) at intake. Of these, 2.8% (N=5) reported drinking alcohol in the last 14 days at intake. No clients reported drinking excessively in the last 14 days at intake.

36 weeks pregnancy

- Information on alcohol use was recorded at 36 weeks pregnancy for 85.6% (N=160) clients. Of these, No reported drinking alcohol in the last 14 days at 36 weeks. No clients reported drinking excessively in the last 14 days at 36 weeks pregnancy.

Intake and 36 weeks

- Information on alcohol use was recorded at both intake and 36 weeks for 157 clients. For these clients, there was:

- -100% change in the proportion of clients drinking alcohol between intake and 36 weeks.
- -2.5% change in the number of clients drinking alcohol between intake and 36 weeks.
- No clients started drinking between intake and 36 weeks.
- 2.5% (N=4) clients stopped drinking between intake and 36 weeks.
- Please note the small numbers of clients who reported drinking alcohol when interpreting the figures for % change in proportion of clients drinking alcohol.

WHSCCT

Intake

- Information on alcohol use was recorded at intake for 92.4% of clients (N=267) at intake. Of these, 1.1% (N=3) reported drinking alcohol in the last 14 days at intake. 0.7% (N=2) clients reported drinking excessively in the last 14 days at intake.

36 weeks pregnancy

- Information on alcohol use was recorded at 36 weeks pregnancy for 81.5% (N=229) clients. Of these, 1.7% (N=4) reported drinking alcohol in the last 14 days at 36 weeks. 0.9% (N=2) clients reported drinking excessively in the last 14 days at 36 weeks pregnancy.

Intake and 36 weeks

- Information on alcohol use was recorded at both intake and 36 weeks for 221 clients. For these clients, there was:
 - No change in the number or proportion of clients drinking alcohol between intake and 36 weeks.
 - -50% change in the proportion of clients drinking alcohol excessively between intake and 36 weeks.
 - -0.5% change in the number of clients drinking alcohol excessively between intake and 36 weeks.
 - 0.9% (N=2) clients started drinking between intake and 36 weeks.
 - 0.9% (N=2) clients stopped drinking between intake and 36 weeks.
- Please note the small numbers of clients who reported drinking alcohol when interpreting the figures for % change in proportion of clients drinking alcohol.

Illegal Drug Use

Table 103: Change in self-reported drug use during pregnancy (intake to 36 weeks) – All sites

		BHSCT	NHSCT	SEHSCT	SHSCT	WHSCT	All Sites
Number of active clients enrolled	N	183	110	138	193	289	913
Clients with information on illegal drug use during pregnancy recorded at intake	N	173	99	135	180	267	854
	%	94.5	90.0	97.8	93.3	92.4	93.5
Clients who reported using illegal drugs in last 14 days at intake	N	0	0	1	0	0	1
	%	0.0	0.0	0.7	0.0	0	0.1
Number of active clients reaching 36 weeks gestation	N	177	102	116	187	281	863
Clients with information on illegal drug use during pregnancy recorded at 36 weeks	N	158	84	109	160	229	740
	%	89.3	82.4	94.0	85.6	81.5	85.7
Clients who reported using illegal drugs in last 14 days at 36 weeks	N	1	0	0	0	0	1
	%	0.6	0.0	0.0	0.0	0.0	0.1
Number of active clients with information on drug use recorded at both intake and 36 weeks*	N	158	82	109	157	221	727
Clients who stopped between intake and 36 weeks *	N	NA	NA	NA	NA	NA	NA
	%	NA	NA	NA	NA	NA	NA
% Change in number of clients with illegal drug use in last 14 days recorded at intake and 36 weeks* [§]	%	0.6	0	0	0	0	0.1
% Change in proportion of clients with illegal drug use in last 14 days recorded at intake and 36 weeks*	%	-	-	-	-	-	-

*These figures are for the subgroup of clients who had data recorded at both intake and 36 weeks. [§]The % change in the number of clients who reported illegal drug use between intake and 36 weeks must be viewed in the context of the very low numbers of clients who reported using illegal drugs for the subgroup of clients who had data recorded at both time points.

All sites

- Information on illegal drug use was recorded for 93.5% (N=854) clients at intake and 85.7% (N=740) clients at 36 weeks pregnancy. 0.1% (N=1) client reported illegal drug use during pregnancy or the last 14 days at intake. 0.1% (N=1) client reported illegal drug use in last 14 days at 36 weeks.

- Information on illegal drug use was recorded for 727 clients at both intake and 36 weeks. For these clients, there was a 0.1% change in the number of clients who reported using illegal drugs between intake and 36 weeks.

BHSCT

- Information on illegal drug use was recorded for 94.5% (N=173) clients at intake and 89.3% (N=158) clients at 36 weeks pregnancy. No clients reported illegal drug use in last 14 days at intake. 0.6% (N=1) client reported illegal drug use at 36 weeks pregnancy.
- Information on illegal drug use was recorded at both intake and 36 weeks for 158 clients. For these clients, there was: 0.6% change in the number of clients with illegal drug use in the last 14 days recorded between intake and 36 weeks.

NHSCT

- Information on illegal drug use was recorded for 90% (N=99) clients at intake and 82.4% (N=84) clients at 36 weeks pregnancy.
- No clients reported illegal drug use in last 14 days at intake or at 36 weeks pregnancy.

SEHSCT

- Information on illegal drug use was recorded for 97.8% (N=135) clients at intake and 94% (N=109) clients at 36 weeks pregnancy. 0.7% (N=1) client reported illegal drug use in last 14 days at intake. No clients reported illegal drug use at 36 weeks pregnancy.
- Information on illegal drug use was recorded at both intake and 36 weeks for 109 clients for whom there was no reported illegal drug use at intake or 36 weeks.

SHSCT

- Information on illegal drug use was recorded for 93.3% (N=180) clients at intake and 85.6% (N=160) clients at 36 weeks pregnancy. No clients reported illegal drug use in last 14 days at intake or at 36 weeks pregnancy.
- Information on illegal drug use was recorded at both intake and 36 weeks for 157 clients for whom there was no reported illegal drug use at intake or 36 weeks.

WHSCT

- Information on illegal drug use was recorded for 92.4% (N=267) clients at intake and 81.5% (N=229) clients at 36 weeks pregnancy. No clients reported illegal drug use in last 14 days at intake or at 36 weeks pregnancy.
- Information on illegal drug use was recorded at both intake and 36 weeks for 221 clients for whom there was no reported illegal drug use at intake or 36 weeks.

Mastery

Mastery at Intake

Table 104: Mastery at intake – All sites

Site	Total number enrolled	Number of Clients with Mastery* recorded - intake	% Clients with mastery recorded - intake	Clients with Low Mastery - intake	% Clients with Low Mastery - intake	Clients with Not Low Mastery - intake	% Clients with Not Low Mastery - intake	Mean Mastery Score - intake	Min Mastery Score - intake	Max Mastery Score - intake
BHSCT	183	168	91.8	31	18.5	137	81.5	22.1	12	28
NHSCT	110	99	90.0	26	26.3	73	73.7	21.5	9	28
SEHSCT	138	135	97.8	24	17.8	111	82.2	22.4	12	28
SHSCT	193	178	92.2	23	12.9	155	87.1	21.9	13	28
WHSCT	289	274	94.8	36	13.1	238	86.9	22.0	13	28
All sites	913	854	93.5	140	16.4	714	83.6	22.0	9	28

* Mastery is a form of perceived personal control, a sense of control over the events in one's life [Pearlin, L. and Schooler, C. (1978).The structure of coping. J Health Soc Behav, 19, 2-21].

All sites

- Overall, mastery was recorded for 93.5% (N=854) of clients at intake. The mean mastery score was 22 (Range= 9 - 28). At intake 16.4% clients (N=140) had low mastery and 83.6% (N=714) of clients did not have low mastery.

BHSCT

- In BHSCT mastery was recorded for 91.8% (N=168) of clients at intake. The mean mastery score was 22.1 (Range= 12 – 28). At intake 18.5% clients (N=31) had low mastery and 81.5% (N=137) of clients did not have low mastery.

NHSCT

- In NHSCT mastery was recorded for 90% (N=99) of clients at intake. The mean mastery score was 21.5 (Range= 9 – 28). At intake 26.3% clients (N=26) had low mastery and 73.7% (N=73) of clients did not have low mastery.

SEHSCT

- In SEHSCT mastery was recorded for 97.8% (N=135) of clients at intake. The mean mastery score was 22.4 (Range= 12 – 28). At intake 17.8% clients (N=24) had low mastery and 82.2% (N=111) of clients did not have low mastery.

SHSCT

- In SHSCT mastery was recorded for 92.2% (N=178) of clients at intake. The mean mastery score was 21.9 (Range= 13 – 28). At intake 12.9% clients (N=23) had low mastery and 87.1% (N=155) of clients did not have low mastery.

WHSCT

- In WHSCT mastery was recorded for 94.8% (N=274) of clients at intake. The mean mastery score was 22 (Range= 13 – 28). At intake 13.1% clients (N=36) had low mastery and 86.9% (N=238) of clients did not have low mastery.

Mastery at 24 Months

Table 105: Mastery at 24 Months – All sites

Site	Number of Clients active at 24mths	Scheduled	Number of Clients with Mastery recorded - 24mths	% Clients with mastery recorded - 24mths	Clients with Low Mastery - 24mths	% Clients with Low Mastery - 24mths	Clients with Not Low Mastery - 24mth	% Clients with Not Low Mastery - 24mth	Mean Mastery Score - 24mths	Min Mastery Score - 24mths	Max Mastery Score - 24mths	Number of clients with Mastery recorded at intake and 24mths	% Change in number of clients with low mastery between intake to 24mths	Number of clients with higher mastery score (improvement) at 24mths than at intake	% of clients with higher mastery score (improvement) at 24mths than at intake
BHSCT	102	102	91	89.2	16	17.6	75	82.4	22.9	15	28	85	1.2	43	50.6
NHSCT	41	41	31	75.6	3	9.7	28	90.3	23.1	17	28	31	-16.1	17	54.8
SEHSCT	54	54	50	92.6	4	8.0	46	92.0	24.2	17	28	49	-4.1	30	61.2
SHSCT	126	126	95	75.4	9	9.5	86	90.5	22.2	15	28	90	-1.1	37	41.1
WHSCT	185	172	110	64.0	16	14.5	94	85.5	22.0	13	28	107	-0.9	37	34.6
All sites	508	495	377	76.2	48	12.7	329	87.3	22.7	13	28	362	-2.2	164	45.3

* The personal belief questions were scheduled for 495 active clients at 24 months (13 of the 508 active clients commenced the programme before the personal belief questions were introduced to be scheduled at 24 months). A total of 417 clients (of the 495 active clients) completed the personal belief questions. However, the total number of clients with Mastery recorded is 377 because 40 clients were excluded from the analysis due the client marked as 'unable to answer' due to learning difficulties/ language barrier for at least one of the seven questions.

All Sites

- Overall, mastery was recorded for 76.2% (N=377) of clients at 24 months. The mean mastery score was 22.7 (Range= 13 - 28). At 24 months 12.7% clients (N=48) had low mastery and 87.3% (N=329) of clients did not have low mastery.
- There was a reduction (-2.2%) in the number of clients who had low mastery from intake to 24 months for clients who had mastery recorded at both time points (N=362). There was also an increase in the number of clients who had higher mastery scores from intake to 24 months as 45.3% (N=164) of clients across all sites had a higher mastery score at 24 months than at intake.

BHSCT

- In BHSCT, mastery was recorded for 89.2% (N=91) of clients at 24 months. The mean mastery score was 22.9 (Range= 15 - 28). At 24 months 17.6% clients (N=16) had low mastery and 82.4% (N=75) of clients did not have low mastery.
- There was an increase (1.2%) in the number of clients who had low mastery from intake to 24 months for clients who had mastery recorded at both time points (N=85). There was also an increase in the number of clients who had higher mastery scores from intake to 24 months as 50.6% (N=43) of clients had a higher mastery score at 24 months than at intake.

NHSCT

- In NHSCT, mastery was recorded for 75.6% (N=31) of clients at 24 months. The mean mastery score was 23.1 (Range= 17 – 28). At 24 months 9.7% clients (N=3) had low mastery and 90.3% (N=28) of clients did not have low mastery.
- There was a reduction (-16.1%) in the number of clients who had low mastery from intake to 24 months for clients who had mastery recorded at both time points (N=31). There was also an increase in the number of clients who had higher mastery scores from intake to 24 months as 54.8% (N=17) of clients had a higher mastery score at 24 months than at intake.

SEHSCT

- In SEHSCT, mastery was recorded for 92.6% (N=50) of clients at 24 months. The mean mastery score was 24.2 (Range= 17 - 28). At 24 months 8% clients (N=4) had low mastery and 92% (N=46) of clients did not have low mastery.
- There was a reduction (-4.1%) in the number of clients who had low mastery from intake to 24 months for clients who had mastery recorded at both time points (N=49). There was also an increase in the number of clients who had higher mastery scores from intake to 24 months as 61.2% (N=30) of clients had a higher mastery score at 24 months than at intake.

SHSCT

- In SHSCT, mastery was recorded for 75.4% (N=95) of clients at 24 months. The mean mastery score was 22.2 (Range= 15 - 28). At 24 months 9.5% (N=9) clients had low mastery and 90.5% (N=86) of clients did not have low mastery.
- There was a reduction (-1.1%) in the number of clients who had low mastery from intake to 24 months for clients who had mastery recorded at both time points (N=90). There was also an increase in the number of clients who had high mastery scores from intake to 24 months as 41.1% (N=37) of clients across all sites had a higher mastery score at 24 months than at intake.

WHSCT

- In WHSCT, mastery was recorded for 64% (N=110) of clients at 24 months. The mean mastery score was 22 (Range= 13 - 28). At 24 months 14.5% (N=16) clients had low mastery and 85.5% (N=94) of clients did not have low mastery.

- There was a reduction (-0.9%) in the number of clients who had low mastery from intake to 24 months for clients who had mastery recorded at both time points (N=107). There was also an increase in the number of clients who had high mastery scores from intake to 24 months as 34.6% (N=37) of clients across all sites had a higher mastery score at 24 months than at intake.

Hospital Admissions due to injury or ingestion

Admissions Due to Injury/Ingestion Birth to 6 Months

Table 106: Admissions Birth to 6 Months – All sites

Site	Active clients reaching 6 months infancy	Number of infants reaching 6 months (including twins)	Infants with hospital admissions due to injury/ingestion between birth and 6 months data recorded		Hospital admissions due to injury/ingestion between birth and 6 months	Infants with 1 hospital admission due to injury/ingestion between birth and 6 months		Infants with 2 or more hospital admissions due to injury/ingestion between birth and 6 months		Mean hospital attendances due to injury/ingestion between birth and 6 months
			N	%		N	%	N	%	
BHSCT	153	154	138	89.6	1	1	0.7	0	0	0.007
NHSCT	86	86	56	65.1	0	0	0.0	0	0	0
SEHSCT	92	93	88	94.6	1	1	1.1	0	0	0.011
SHSCT	161	163	145	89.0	0	0	0.0	0	0	0
WHSCT	252	252	205	81.3	0	0	0.0	0	0	0
All sites	744	748	632	84.5	2	2	0.3	0	0	0.003

This table reports on hospital admissions due to injury or ingestion for infants reaching 6 months (N= 748) by 31.12.18.

All sites

- Overall, 748 infants reached 6 months infancy and information on hospital admissions was recorded for 84.5% (N=632) of these clients.
- Two infants (0.3%) were recorded with having 1 hospital admission due to injury/ingestion between birth and 6 months.
- Overall, the mean hospital admission due to injury or ingestion between birth and 6 months is 0.003.

BHSCT

- In the BHSCT, 154 infants reached 6 months infancy and information on hospital admissions was recorded for 89.6% (N=138) of these clients.
- One infant (0.7%) was recorded with having 1 hospital admission due to injury/ingestion between birth and 6 months.
- Overall, in the BHSCT, the mean hospital attendance due to injury/ingestion between birth and 6 months was 0.007.

NHSCT

- In the NHSCT, 86 infants reached 6 months infancy and information on hospital admissions was recorded for 65.1% (N=56) of these clients. There were no hospital admissions recorded between birth and 6 months.

SEHSCT

- In the SEHSCT, 93 infants reached 6 months infancy and information on hospital admissions was recorded for 94.6% (N=88) of these clients.
- One infant (1.1%) was recorded with having 1 hospital admission due to injury/ingestion between birth and 6 months.
- Overall, in the SEHSCT, the mean hospital attendance due to injury/ingestion between birth and 6 months was 0.011

SHSCT

- In the SHSCT, 163 infants reached 6 months infancy and information on hospital admissions was recorded for 89% (N=145) of these clients. There were no hospital admissions recorded between birth and 6 months.

WHSCT

- In the WHSCT, 252 infants reached 6 months infancy and information on hospital admissions was recorded for 81.3% (N=205) of these clients. There were no hospital admissions recorded between birth and 6 months.

Admissions Due to Injury/Ingestion Birth to 12 Months

Table 107: Admissions Birth to 12 Months – All sites

Site	Active clients reaching 12 months infancy	Number of infants reaching 12 months (including twins)	Infants with hospital admissions due to injury/ingestion between birth and 12 months data recorded		Hospital admissions due to injury/ingestion between birth and 12 months	Infants with 1 hospital admission due to injury/ingestion between birth and 12 months		Infants with 2 or more hospital admissions due to injury/ingestion between birth and 12 months		Mean hospital attendances due to injury/ingestion between birth and 12 months
			N	%		N	%	N	%	
BHSCT	136	137	122	89.1	0	0	0	0	0	0
NHSCT	68	68	41	60.3	0	0	0	0	0	0
SEHSCT	88	89	79	88.8	0	0	0	0	0	0
SHSCT	147	149	110	73.8	1	1	0.9	0	0	0.01
WHSCT	228	228	196	86.0	2	2	1.0	0	0	0.01
All sites	667	671	548	81.7	3	3	0.5	0	0	0.01

This table reports on hospital admissions due to injury or ingestion for infants reaching 12 months (N=671) by 31.12.18.

All sites

- Overall, 671 infants reached 12 months infancy and information on hospital admissions was recorded for 81.7% (N=548) of these clients.
- Three infants (0.5%) had 1 hospital admission due to injury or ingestion between birth and 12 months.
- No infants recorded 2 or more hospital admission due to injury or ingestion between birth and 12 months.
- Overall, the mean hospital admission due to injury or ingestion between birth and 12 months is 0.01.

BHSCT

- 137 infants reached 12 months by the end of December 2018 and information on hospital admissions was recorded for 89.1% (N=122) of these clients.
- No infants recorded any hospital admissions due to injury or ingestion between birth and 12 months.

NHSCT

- 68 infants reached 12 months by the end of December 2018 and information on hospital admissions was recorded for 60.3% (N=41) of these clients.
- No infants recorded any hospital admissions due to injury or ingestion between birth and 12 months.

SEHSCT

- 89 infants reached 12 months by the end of December 2018 and information on hospital admissions was recorded for 88.8% (N=79) of these clients.
- No infants recorded any hospital admissions due to injury or ingestion between birth and 12 months.

SHSCT

- 149 infants reached 12 months by the end of December 2018 and information on hospital admissions was recorded for 73.8% (N=110) of these clients.
- One infant (0.9%) recorded having 1 hospital admission due to injury or ingestion between birth and 12 months.
- Overall, in the SHSCT, the mean hospital attendance due to injury/ingestion between birth and 12 months was 0.01.

WHsCT

- 228 infants reached 12 months by the end of December 2018 and information on hospital admissions was recorded for 86% (N=196) of these clients.
- Two infants (1%) were recorded with having 1 hospital admission due to injury or ingestion between birth and 12 months.
- Overall, in the WHsCT, the mean hospital attendance due to injury/ingestion between birth and 12 months was 0.01.

Admissions 12 to 18 Months

Table 108: Admissions 12 to 18 Months – All sites

Site	Active clients reaching 18 months infancy	Number of infants reaching 18 months (including twins)	Infants with hospital admissions due to injury/ingestion between 12 and 18 months data recorded		Hospital admissions due to injury/ingestion between 12 and 18 months	Infants with 1 hospital admission due to injury/ingestion between 12 and 18 months		Infants with 2 or more hospital admissions due to injury/ingestion between 12 and 18 months		Mean hospital attendances due to injury/ingestion between 12 and 18 months
			N	%		N	%	N	%	
BHSCT	118	119	105	88.2	2	2	1.9	0	0	0.02
NHSCT	54	54	35	64.8	0	0	0.0	0	0	0
SEHSCT	69	69	63	91.3	0	0	0.0	0	0	0
SHSCT	133	134	71	53.0	0	0	0.0	0	0	0
WHSCT	204	204	160	78.4	6	4	2.5	1	0.6	0.04
All sites	578	580	434	74.8	8	6	1.4	1	0.2	0.02

This table reports on hospital admissions due to injury or ingestion for infants reaching 18 months (N=580) by 31.12.18.

All sites

- Overall, 580 infants reached 18 months toddlerhood and information on hospital admissions was recorded for 74.8% (N=434) of these clients.
- Six infants (1.4%) were recorded with having 1 hospital admission due to injury or ingestion between 12 and 18 months.
- One infant (0.2%) was recorded with having 2 or more hospital admissions due to injury or ingestion between 12 and 18 months.
- Overall, the mean hospital admissions due to injury or ingestion between 12 and 18 months is 0.02.

BHSCT

- 119 infants reached 18 months toddlerhood and information on hospital admissions was recorded for 88.2% (N=105) of these clients.

- Two infants (1.9%) were recorded with having 1 hospital admission due to injury or ingestion between 12 and 18 months.
- No infants recorded 2 or more hospital admission due to injury or ingestion between 12 and 18 months.
- Overall, the mean hospital admissions due to injury or ingestion between 12 and 18 months is 0.02.

NHSCT

- 54 infants reached 18 months toddlerhood and information on hospital admissions was recorded for 64.8% (N=35) of these clients.
- No infants had any hospital admission due to injury or ingestion between 12 and 18 months.

SEHSCT

- 69 infants reached 18 months toddlerhood and information on hospital admissions was recorded for 91.3% (N=63) of these clients.
- No infants had any hospital admission due to injury or ingestion between 12 and 18 months.

SHSCT

- 134 infants reached 18 months toddlerhood and information on hospital admissions was recorded for 53% (N=71) of these clients.
- No infants had any hospital admission due to injury or ingestion between 12 and 18 months.

WHSCT

- 204 infants reached 18 months toddlerhood and information on hospital admissions was recorded for 78.4% (N=160) of these clients.
- Four infants (2.5%) were recorded with having 1 hospital admission due to injury or ingestion between 12 and 18 months.
- One infant (0.6%) was recorded with having 2 or more hospital admission due to injury or ingestion between 12 and 18 months.
- Overall, the mean hospital admissions due to injury or ingestion between 12 and 18 months is 0.04.

Admissions 12 to 24 Months

Table 109: Admissions 12 to 24 Months – All sites

Site	Active clients reaching 24 months infancy	Number of infants reaching 24 months (including twins)	Infants with hospital admissions due to injury/ingestion between 12 and 24 months recorded		Hospital admissions due to injury/ingestion between 12 and 24 months	Infants with 1 hospital admission due to injury/ingestion between 12 and 24 months		Infants with 2 or more hospital admissions due to injury/ingestion between 12 and 24 months		Mean hospital attendances due to injury/ingestion on between 12 and 24 months
			N	%		N	%	N	%	
BHSCT	102	103	89	86.4	1	1	1.1	0	0	0.01
NHSCT	41	41	26	63.4	0	0	0.0	0	0	0
SEHSCT	54	54	49	90.7	1	1	2.0	0	0	0.02
SHSCT	126	127	96	75.6	0	0	0.0	0	0	0
WHSCT	185	185	157	84.9	7	7	4.5	0	0	0.04
All sites	508	510	417	81.8	9	9	2.2	0	0	0.02

This table reports on hospital admissions due to injury or ingestion for infants reaching 24 months (N=510) by 31.12.18.

All sites

- Overall, 510 infants reached 24 months toddlerhood and information on hospital admissions was recorded for 81.8% (N=417) of these clients.
- Nine infants (2.2%) were recorded with having 1 hospital admission due to injury or ingestion between 12 and 24 months.
- No infants recorded 2 or more hospital admission due to injury or ingestion between 12 and 24 months.
- Overall, the mean hospital admissions due to injury or ingestion between 12 and 24 months is 0.02.

BHSCT

- Overall, 103 infants reached 24 months toddlerhood and information on hospital admissions was recorded for 86.4% (N=89) of these clients.

- One infant (1.1%) was recorded with having 1 hospital admission due to injury or ingestion between 12 and 24 months.
- No infants recorded 2 or more hospital admission due to injury or ingestion between 12 and 24 months.
- Overall, the mean hospital admissions due to injury or ingestion between 12 and 24 months is 0.01.

NHSCT

- Overall, 41 infants reached 24 months toddlerhood and information on hospital admissions was recorded for 63.4% (N=26) of these clients.
- No infants had any hospital admission due to injury or ingestion between 12 and 24 months.

SEHSCT

- Overall, 54 infants reached 24 months toddlerhood and information on hospital admissions was recorded for 90.7% (N=49) of these clients.
- One infant (2.0%) was recorded with having 1 hospital admission due to injury or ingestion between 12 and 24 months.
- No infants recorded 2 or more hospital admission due to injury or ingestion between 12 and 24 months.
- Overall, the mean hospital admissions due to injury or ingestion between 12 and 24 months is 0.02.

SHSCT

- Overall, 127 infants reached 24 months toddlerhood and information on hospital admissions was recorded for 75.6% (N=96) of these clients.
- No infants had any hospital admission due to injury or ingestion between 12 and 24 months.

WHSCT

- Overall, 185 infants reached 24 months toddlerhood and information on hospital admissions was recorded for 84.9% (N=157) of these clients.
- Seven infants (4.5%) were recorded with having 1 hospital admission due to injury or ingestion between 12 and 24 months.
- No infants recorded 2 or more hospital admission due to injury or ingestion between 12 and 24 months.
- Overall, the mean hospital admissions due to injury or ingestion between 12 and 24 months is 0.04.

ED attendances due to injury or ingestion

A&E Birth to 6 Months

Table 110: A&E Birth to 6 Months – All sites

Site	Active clients reaching 6 months infancy	Number of infants reaching 6 months (including twins)	Infants with A&E visits due to injury/ingestion between birth and 6 months recorded		A&E visits due to injury/ingestion between birth and 6 months	Infants with 1 A&E visit due to injury/ingestion between birth and 6 months		Infants with 2 or more A&E visits due to injury/ingestion between birth and 6 months		Mean visits to A&E due to injury/ingestion between birth and 6 months
	N	N	N	%	N	N	%	N	%	Mean
BHSCT	153	154	138	89.6	4	4	2.9	0	0	0.03
NHSCT	86	86	56	65.1	4	4	7.1	0	0	0.07
SEHSCT	92	93	88	94.6	3	3	3.4	0	0	0.03
SHSCT	161	163	145	89.0	6	4	2.8	1	0.7	0.04
WHSCT	252	252	205	81.3	11	11	5.4	0	0	0.05
All sites	744	748	632	84.5	28	26	4.1	1	0.2	0.04

This table reports on A&E attendances due to injury or ingestion for infants reaching 6 months (N=748) by 31.12.18.

All sites

- 748 infants reached 6 months by the end of December 2018 and information on A&E visits was recorded for 84.5% (N=632) of these clients.
- Overall, 4.1% (N=26) infants had 1 A&E visit due to injury or ingestion between birth and 6 months.
- Overall, 0.2% (N=1) infant had 2 or more A&E visits due to injury or ingestion between birth and 6 months.
- Overall, the mean visits to A&E due to injury or ingestion between birth and 6 months is 0.04.

BHSCT

- 154 infants reached 6 months by the end of December 2018 and information on A&E visits was recorded for 89.6% (N=138) of these clients.
- Overall, 2.9% (N=4) infants had 1 A&E visit due to injury or ingestion between birth and 6 months.
- No infants had 2 or more A&E visits due to injury or ingestion between birth and 6 months.
- Overall, the mean visits to A&E due to injury or ingestion between birth and 6 months is 0.03.

NHSCT

- 86 infants reached 6 months by the end of December 2018 and information on A&E visits was recorded for 65.1% (N=56) of these clients.
- Overall, 7.1% (N=4) infants had 1 A&E visit due to injury or ingestion between birth and 6 months.
- No infants had 2 or more A&E visits due to injury or ingestion between birth and 6 months.
- Overall, the mean visits to A&E due to injury or ingestion between birth and 6 months is 0.07.

SEHSCT

- 93 infants reached 6 months by the end of December 2018 and information on A&E visits was recorded for 94.6% (N=88) of these clients.
- Overall, 3.4% (N=3) infants had 1 A&E visit due to injury or ingestion between birth and 6 months.
- No infants had 2 or more A&E visits due to injury or ingestion between birth and 6 months.
- Overall, the mean visits to A&E due to injury or ingestion between birth and 6 months is 0.03.

SHSCT

- 163 infants reached 6 months by the end of December 2018 and information on A&E visits was recorded for 89% (N=145) of these clients.
- Overall, 2.8% (N=4) infants had 1 A&E visit due to injury or ingestion between birth and 6 months.
- In the SHSCT, 0.7% (N=1) infants had 2 or more A&E visit due to injury or ingestion between birth and 6 months.
- Overall, in the SHSCT the mean visits to A&E due to injury or ingestion between birth and 6 months are 0.04.

WHSCT

- 252 infants reached 6 months by the end of December 2018 and information on A&E visits was recorded for 81.3% (N=205) of these clients.
- Overall, 5.4% (N=11) infants had 1 A&E visit due to injury or ingestion between birth and 6 months.
- No infants had 2 or more A&E visits due to injury or ingestion between birth and 6 months.

- Overall, the mean visits to A&E due to injury or ingestion between birth and 6 months is 0.05.

A&E Birth to 12 months

Table 111: A&E Birth to 12 months – All sites

Site	Active clients reaching 12 months infancy	Number of infants reaching 12 months (including twins)	Infants with A&E visits due to injury/ingestion between birth and 12 months recorded		A&E visits due to injury/ingestion between birth and 12 months	Infants with 1 A&E visit due to injury/ingestion between birth and 12 months		Infants with 2 or more A&E visits due to injury/ingestion between birth and 12 months		Mean visits to A&E due to injury/ingestion between birth and 12 months
			N	%		N	%	N	%	
BHSCT	136	137	122	89.1	7	5	3.6	1	0.7	0.05
NHSCT	68	68	41	60.3	3	3	4.4	0	0.0	0.05
SEHSCT	88	89	79	88.8	5	5	5.6	0	0.0	0.06
SHSCT	147	149	110	73.8	7	2	1.3	2	1.3	0.05
WHSCT	228	228	196	86.0	17	17	7.5	0	0.0	0.08
All sites	667	671	548	81.7	39	32	4.8	3	0.4	0.06

This table reports on A&E attendances due to injury or ingestion for infants reaching 12 months (N=671) by 31.12.18.

All sites

- 671 infants reached 12 months by the end of December 2018 and information on A&E visits was recorded for 81.7% (N=548) of these clients.
- Overall, 4.8% (N=32) infants had 1 A&E visit due to injury or ingestion between birth and 12 months.
- Overall, 0.4% (N=3) infant had 2 or more A&E visits due to injury or ingestion between birth and 12 months.
- Overall, the mean visits to A&E due to injury or ingestion between birth and 12 months is 0.06.

BHSCT

- 137 infants reached 12 months by the end of December 2018 and information on A&E visits was recorded for 89.1% (N=122) of these clients.

- Overall, 3.6% (N=5) infants had 1 A&E visit due to injury or ingestion between birth and 12 months.
- Overall, 0.7% (N=1) infant had 2 or more A&E visits due to injury or ingestion between birth and 12 months.
- Overall, the mean visits to A&E due to injury or ingestion between birth and 12 months is 0.05.

NHSCT

- 68 infants reached 12 months by the end of December 2018 and information on A&E visits was recorded for 60.3% (N=41) of these clients.
- Overall, 4.4% (N=3) infants had 1 A&E visit due to injury or ingestion between birth and 12 months.
- No infants had 2 or more A&E visits due to injury or ingestion between birth and 12 months.
- Overall, the mean visits to A&E due to injury or ingestion between birth and 12 months is 0.05.

SEHSCT

- 89 infants reached 12 months by the end of December 2018 and information on A&E visits was recorded for 88.8% (N=79) of these clients.
- Overall, 5.6% (N=5) infants had 1 A&E visit due to injury or ingestion between birth and 12 months.
- No infants had 2 or more A&E visits due to injury or ingestion between birth and 12 months.
- Overall, the mean visits to A&E due to injury or ingestion between birth and 12 months is 0.06.

SHSCT

- 149 infants reached 12 months by the end of December 2018 and information on A&E visits was recorded for 73.8% (N=110) of these clients.
- Overall, 1.3% (N=2) infants had 1 A&E visit due to injury or ingestion between birth and 12 months.
- Overall, 1.3% (N=2) infant had 2 or more A&E visits due to injury or ingestion between birth and 12 months.
- Overall, the mean visits to A&E due to injury or ingestion between birth and 12 months is 0.05.

WHSCT

- 228 infants reached 12 months by the end of December 2018 and information on A&E visits was recorded for 86% (N=196) of these clients.
- Overall, 7.5% (N=17) infants had 1 A&E visit due to injury or ingestion between birth and 12 months.
- No infants had 2 or more A&E visits due to injury or ingestion between birth and 12 months.
- Overall, the mean visits to A&E due to injury or ingestion between birth and 12 months is 0.08.

A&E 12 to 18 months

Table 112: A&E 12 to 18 months – All sites

Site	Active clients reaching 18 months infancy	Number of infants reaching 18 months (including twins)	Infants with A&E visits due to injury/ingestion between 12 and 18 months recorded		A&E visits due to injury/ingestion between 12 and 18 months	Infants with 1 A&E visit due to injury/ingestion between 12 and 18 months		Infants with 2 or more A&E visits due to injury/ingestion between 12 and 18 months		Mean visits to A&E due to injury/ingestion between 12 and 18 months
			N	%		N	%	N	%	
BHSCT	118	119	105	88.2	6	3	2.9	1	1.0	0.06
NHSCT	54	54	35	64.8	2	2	5.7	0	0.0	0.06
SEHSCT	69	69	63	91.3	4	1	1.6	1	1.6	0.06
SHSCT	133	134	71	53.0	1	1	1.4	0	0.0	0.01
WHsCT	204	204	160	78.4	17	13	8.1	2	1.3	0.11
All sites	578	580	434	74.8	30	20	4.6	4	0.9	0.07

This table reports on A&E attendances due to injury or ingestion overall for infants reaching 18 months (N=580) by 31.12.18.

All sites

- 580 infants reached 18 months by the end of December 2018 and information on A&E visits was recorded for 74.8% (N=434) of these clients.
- Overall, 4.6% (N=20) infants had 1 A&E visit due to injury or ingestion between 12 and 18 months.
- Overall, 0.9% (N=4) infant had 2 or more A&E visits due to injury or ingestion between 12 and 18 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 18 months is 0.07.

BHSCT

- 119 infants reached 18 months by the end of December 2018 and information on A&E visits was recorded for 88.2% (N=105) of these clients.

- Overall, 2.9% (N=3) infants had 1 A&E visit due to injury or ingestion between 12 and 18 months.
- Overall, 1.0% (N=1) infant had 2 or more A&E visits due to injury or ingestion between 12 and 18 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 18 months is 0.06.

NHSCT

- 54 infants reached 18 months by the end of December 2018 and information on A&E visits was recorded for 64.8% (N=35) of these clients.
- Overall, 5.7% (N=2) infants had 1 A&E visit due to injury or ingestion between 12 and 18 months.
- No infants had 2 or more A&E visits due to injury or ingestion between 12 and 18 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 18 months is 0.06.

SEHSCT

- 69 infants reached 18 months by the end of December 2018 and information on A&E visits was recorded for 91.3% (N=63) of these clients.
- Overall, 1.6% (N=1) infants had 1 A&E visit due to injury or ingestion between 12 and 18 months.
- Overall, 1.6% (N=1) infant had 2 or more A&E visits due to injury or ingestion between 12 and 18 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 18 months is 0.06.

SHSCT

- 134 infants reached 18 months by the end of December 2018 and information on A&E visits was recorded for 53% (N=71) of these clients.
- Overall, 1.4% (N=1) infants had 1 A&E visit due to injury or ingestion between 12 and 18 months.
- No infants had 2 or more A&E visits due to injury or ingestion between 12 and 18 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 18 months is 0.01.

WHSCT

- 204 infants reached 18 months by the end of December 2018 and information on A&E visits was recorded for 78.4% (N=160) of these clients.
- Overall, 8.1% (N=13) infants had 1 A&E visit due to injury or ingestion between 12 and 18 months.
- Overall, 1.3% (N=2) infant had 2 or more A&E visits due to injury or ingestion between 12 and 18 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 18 months is 0.11.

A&E 12 to 24 months

Table 113: A&E 12 to 24 months – All sites

Site	Active clients reaching 24 months infancy	Number of infants reaching 24 months (including twins)	Infants with A&E visits due to injury/ingestion between 12 and 24 months recorded		A&E visits due to injury/ingestion between 12 and 24 months	Infants with 1 A&E visit due to injury/ingestion between 12 and 24 months		Infants with 2 or more A&E visits due to injury/ingestion between 12 and 24 months		Mean visits to A&E due to injury/ingestion between 12 and 24 months
			N	%		N	%	N	%	
BHSCT	102	103	89	86.4	5	3	3.4	1	1.1	0.06
NHSCT	41	41	26	63.4	0	0	0.0	0	0.0	0
SEHSCT	54	54	49	90.7	1	1	2.0	0	0.0	0.02
SHSCT	126	127	96	75.6	5	5	5.2	0	0.0	0.05
WHSCT	185	185	157	84.9	19	15	9.6	2	1.3	0.12
All sites	508	510	417	81.8	30	24	5.8	3	0.7	0.07

This table reports on A&E attendances due to injury or ingestion for infants reaching 24 months (N=510) by 31.12.18.

All sites

- 510 infants reached 24 months by the end of December 2018 and information on A&E visits was recorded for 81.8% (N=417) of these clients.
- Overall, 5.8% (N=24) infants had 1 A&E visit due to injury or ingestion between 12 and 24 months.
- Overall, 0.7% (N=3) infant had 2 or more A&E visits due to injury or ingestion between 12 and 24 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 24 months is 0.07.

BHSCT

- 103 infants reached 24 months by the end of December 2018 and information on A&E visits was recorded for 86.4% (N=89) of these clients.
- Overall, 3.4% (N=3) infants had 1 A&E visit due to injury or ingestion between 12 and 24 months.
- Overall, 1.1% (N=1) infant had 2 or more A&E visits due to injury or ingestion between 12 and 24 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 24 months is 0.06.

NHSCT

- 41 infants reached 24 months by the end of December 2018 and information on A&E visits was recorded for 63.4% (N=26) of these clients.
- No infants recorded any A&E visits due to injury or ingestion between 12 and 24 months.

SEHSCT

- 54 infants reached 24 months by the end of December 2018 and information on A&E visits was recorded for 90.7% (N=49) of these clients.
- Overall, 2% (N=1) infants had 1 A&E visit due to injury or ingestion between 12 and 24 months.
- No infants had recorded 2 or more A&E visits due to injury or ingestion between 12 and 24 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 24 months is 0.02.

SHSCT

- 127 infants reached 24 months by the end of December 2018 and information on A&E visits was recorded for 75.6% (N=96) of these clients.
- Overall, 5.2% (N=5) infants had 1 A&E visit due to injury or ingestion between 12 and 24 months.
- No infants recorded had 2 or more A&E visits due to injury or ingestion between 12 and 24 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 24 months is 0.05.

WHSCT

- 185 infants reached 24 months by the end of December 2018 and information on A&E visits was recorded for 84.9% (N=157) of these clients.
- Overall, 9.6% (N=15) infants had 1 A&E visit due to injury or ingestion between 12 and 24 months.
- Overall, 1.3% (N=2) infant had 2 or more A&E visits due to injury or ingestion between 12 and 24 months.
- Overall, the mean visits to A&E due to injury or ingestion between 12 and 24 months is 0.12.

ASQ

ASQ Infancy

Table 114: ASQ Infancy – All sites

ASQ* Infancy		Communication	Gross Motor	Fine Motor	Problem Solving	Personal Social
Clients ever enrolled		913	913	913	913	913
4 months (I16 visit)						
Active clients at 4 months	N	744	744	744	744	744
Number of infants for active clients at 4 months**	N	748	748	748	748	748
Infants with ASQ data recorded at 4 months	N	622	622	622	622	622
	%	83.2	83.2	83.2	83.2	83.2
Mean ASQ 4 month score	N	56.3	54.7	54.8	56.2	55.9
Seek consultation' cut off scores at 4 months		34.60	38.41	29.62	34.98	33.16
Infants with scores below 'seek consultation' cut off at 4 months	N	10	30	13	10	12
	%	1.6	4.8	2.1	1.6	1.9
10 months (I28 visit)						
Active clients at 10 months	N	667	667	667	667	667
Number of infants for active clients at 10 months*	N	671	671	671	671	671
Infants with ASQ data recorded at 10 months	N	543	543	543	543	543
	%	80.9	80.9	80.9	80.9	80.9
Mean ASQ 10 month score	N	55.7	50.1	56.0	54.7	53.9
Seek consultation' cut off scores at 10 months		22.87	30.07	37.97	32.51	27.25
Infants with scores below 'seek consultation' cut off at 10 months	N	7	86	19	20	9
	%	1.3	15.8	3.5	3.7	1.7

* The Ages and Stages Questionnaire (ASQ) is a general developmental screening tool [Squires, J & Bricker, D. Ages & Stages Questionnaires®, Third Edition (ASQ- 3™). A parent-completed child-monitoring system. Paul H. Brookes Publishing Co., Baltimore (2009)].

**Two clients had twins

All sites

4 Months

- Overall, 748 infants reached 4 months infancy by the end of December 2018 and ASQ data was recorded for 83.2% (N=622) of these clients.
- Communication: 1.6% (N=10) infants had a score below the seek consultation cut-off (34.60). The overall mean for communication is 56.3.
- Gross motor: 4.8% (N=30) infants had a score below the seek consultation cut-off (38.41). The overall mean for gross motor is 54.7.
- Fine motor: 2.1% (N=13) infants had a score below the seek consultation cut-off (29.62). The overall mean for fine motor is 54.8.
- Problem solving: 1.6% (N=10) infants had a score below the seek consultation cut-off (34.98). The overall mean for problem solving is 56.2.
- Personal Social: 1.9% (N=12) infants had a score below the seek consultation cut-off (33.16). The overall mean for personal social is 55.9.

10 Months

- Overall, 671 infants reached 10 months infancy by the end of December 2018 and ASQ data was recorded for 80.9% (N=543) of these clients.
- Communication: 1.3% (N=7) infants had a score below the seek consultation cut-off (22.87). The overall mean for communication is 55.7.
- Gross motor: 15.8% (N=86) infants had a score below the seek consultation cut-off (30.07). The overall mean for gross motor is 50.1.
- Fine motor: 3.5% (N=19) infants had a score below the seek consultation cut-off (37.97). The overall mean for fine motor is 56.0.
- Problem solving: 3.7% (N=20) infants had a score below the seek consultation cut-off (32.51). The overall mean for problem solving is 54.7.
- Personal Social: 1.7% (N=9) infants had a score below the seek consultation cut-off (27.25). The overall mean for personal social is 53.9.

Table 115: ASQ Infancy – BHSCT and all sites

ASQ Infancy		Communication		Gross Motor		Fine Motor		Problem Solving		Personal Social	
		BHSCT	All site total	BHSCT	All site total	BHSCT	All site total	BHSCT	All site total	BHSCT	All site total
Clients ever enrolled		184	913	184	913	184	913	184	913	184	913
4 months (I16 visit)											
Active clients at 4 months	N	153	744	153	744	153	744	153	744	153	744
Number of infants for active clients at 4 months*	N	154	748	154	748	154	748	154	748	154	748
Infants with ASQ data recorded at 4 months	N	135	622	135	622	135	622	135	622	135	622
	%	87.7	83.2	87.7	83.2	87.7	83.2	87.7	83.2	87.7	83.2
Mean ASQ 4 month score	N	57.4	56.3	56.6	54.7	56.7	54.8	57.4	56.2	57.4	55.9
Seek consultation' cut off scores at 4 months		34.60	34.60	38.41	38.41	29.62	29.62	34.98	34.98	33.16	33.16
Infants with scores below 'seek consultation' cut off at 4 months	N	2	10	3	30	0	13	1	10	1	12
	%	1.5	1.6	2.2	4.8	0	2.1	0.7	1.6	0.7	1.9
10 months											
Active clients at 10 months	N	136	667	136	667	136	667	136	667	136	667
Number of infants for active clients at 10 months	N	137	671	137	671	137	671	137	671	137	671
Infants with ASQ data recorded at 10 months	N	121	543	121	543	121	543	121	543	121	543
	%	88.3	80.9	88.3	80.9	88.3	80.9	88.3	80.9	88.3	80.9
Mean ASQ 10 month score	N	55.6	55.7	50.5	50.1	55.9	56.0	54.5	54.7	54.0	53.9
Seek consultation' cut off scores at 10 months		22.87	22.87	30.07	30.07	37.97	37.97	32.51	32.51	27.25	27.25
Infants with scores below 'seek consultation' cut off at 10 months	N	1	7	20	86	4	19	7	20	2	9
	%	0.8	1.3	16.5	15.8	3.3	3.5	5.8	3.7	1.7	1.7

*One client had twins

BHSCT

4 Months

- Overall, 154 infants reached 4 months infancy by the end of December 2018 and ASQ data was recorded for 87.7% (N=135) of these clients.
- Communication: 1.5% (N=2) infants had a score below the seek consultation cut-off (34.60). The overall mean for communication is 57.4.
- Gross motor: 2.2% (N=3) infants had a score below the seek consultation cut-off (38.41). The overall mean for gross motor is 56.6.
- Fine motor: No infants had a score below the seek consultation cut-off (29.62). The overall mean for fine motor is 56.7.
- Problem solving: 0.7% (N=1) infants had a score below the seek consultation cut-off (34.98). The overall mean for problem solving is 57.4.
- Personal Social: 0.7% (N=1) infants had a score below the seek consultation cut-off (33.16). The overall mean for personal social is 57.4.

10 Months

- Overall, 137 infants reached 10 months infancy by the end of December 2018 and ASQ data was recorded for 88.3% (N=121) of these clients.
- Communication: 0.8% (N=1) infants had a score below the seek consultation cut-off (22.87). The overall mean for communication is 55.6.
- Gross motor: 16.5% (N=20) infants had a score below the seek consultation cut-off (30.07). The overall mean for gross motor is 50.5.
- Fine motor: 3.3% (N=4) infants had a score below the seek consultation cut-off (37.97). The overall mean for fine motor is 55.9.
- Problem solving: 5.8% (N=7) infants had a score below the seek consultation cut-off (32.51). The overall mean for problem solving is 54.5.
- Personal Social: 1.7% (N=2) infants had a score below the seek consultation cut-off (27.25). The overall mean for personal social is 54.0.

Table 116 ASQ Infancy- NHSCT and all sites

ASQ Infancy		Communication		Gross Motor		Fine Motor		Problem Solving		Personal Social	
		NHSCT	All site total	NHSCT	All site total	NHSCT	All site total	NHSCT	All site total	NHSCT	All site total
Clients ever enrolled		110	913	110	913	110	913	110	913	110	913
4 months (I16 visit)											
Active clients at 4 months	N	86	744	86	744	86	744	86	744	86	744
Number of infants for active clients at 4 months	N	86	748	86	748	86	748	86	748	86	748
Infants with ASQ data recorded at 4 months	N	54	622	54	622	54	622	54	622	54	622
	%	62.8	83.2	62.8	83.2	62.8	83.2	62.8	83.2	62.8	83.2
Mean ASQ 4 month score	N	54.5	56.3	53.5	54.7	51.4	54.8	54.3	56.2	55.3	55.9
Seek consultation' cut off scores at 4 months		34.60	34.60	38.41	38.41	29.62	29.62	34.98	34.98	33.16	33.16
Infants with scores below 'seek consultation' cut off at 4 months	N	2	10	4	30	3	13	1	10	2	12
	%	3.7	1.6	7.4	4.8	5.6	2.1	1.9	1.6	3.7	1.9
10 months											
Active clients at 10 months	N	68	667	68	667	68	667	68	667	68	667
Number of infants for active clients at 10 months	N	68	671	68	671	68	671	68	671	68	671
Infants with ASQ data recorded at 10 months	N	40	543	40	543	40	543	40	543	40	543
	%	58.8	80.9	58.8	80.9	58.8	80.9	58.8	80.9	58.8	80.9
Mean ASQ 10 month score	N	53.1	55.7	45.4	50.1	52.5	56.0	50.1	54.7	49.4	53.9
Seek consultation' cut off scores at 10 months		22.87	22.87	30.07	30.07	37.97	37.97	32.51	32.51	27.25	27.25
Infants with scores below 'seek consultation' cut off at 10 months	N	2	7	8	86	3	19	2	20	2	9
	%	5.0	1.3	20.0	15.8	7.5	3.5	5.0	3.7	5.0	1.7

NHSCT

4 Months

- Overall, 86 infants reached 4 months infancy by the end of December 2018 and ASQ data was recorded for 62.8% (N=54) of these clients.
- Communication: 3.7% (N=2) infants had a score below the seek consultation cut-off (34.60). The overall mean for communication is 54.5.
- Gross motor: 7.4% (N=4) infants had a score below the seek consultation cut-off (38.41). The overall mean for gross motor is 53.5.
- Fine motor: 5.6% (N=3) infants had a score below the seek consultation cut-off (29.62). The overall mean for fine motor is 51.4.
- Problem solving: 1.9% (N=1) infants had a score below the seek consultation cut-off (34.98). The overall mean for problem solving is 54.3.
- Personal Social: 3.7% (N=2) infants had a score below the seek consultation cut-off (33.16). The overall mean for personal social is 55.3.

10 Months

- Overall, 68 infants reached 10 months infancy by the end of December 2018 and ASQ data was recorded for 58.8% (N=40) of these clients.
- Communication: 5% (N=2) infants had a score below the seek consultation cut-off (22.87). The overall mean for communication is 53.1.
- Gross motor: 20% (N=8) infants had a score below the seek consultation cut-off (30.07). The overall mean for gross motor is 45.4.
- Fine motor: 7.5% (N=3) infants had a score below the seek consultation cut-off (37.97). The overall mean for fine motor is 52.5.
- Problem solving: 5% (N=2) infants had a score below the seek consultation cut-off (32.51). The overall mean for problem solving is 50.1.
- Personal Social: 5% (N=2) infants had a score below the seek consultation cut-off (27.25). The overall mean for personal social is 49.4.

Table 117: ASQ Infancy – SEHSCT and all sites

ASQ Infancy		Communication		Gross Motor		Fine Motor		Problem Solving		Personal Social	
		SEHSCT	All site total	SEHSCT	All site total	SEHSCT	All site total	SEHSCT	All site total	SEHSCT	All site total
Clients ever enrolled		138	913	138	913	138	913	138	913	138	913
4 months (I16 visit)											
Active clients at 4 months	N	92	744	92	744	92	744	92	744	92	744
Number of infants for active clients at 4 months	N	93	748	93	748	93	748	93	748	93	748
Infants with ASQ data recorded at 4 months	N	88	622	88	622	88	622	88	622	88	622
	%	94.6	83.2	94.6	83.2	94.6	83.2	94.6	83.2	94.6	83.2
Mean ASQ 4 month score	N	56.6	56.3	55.5	54.7	54.0	54.8	57.6	56.2	56.1	55.9
Seek consultation' cut off scores at 4 months		34.60	34.60	38.41	38.41	29.62	29.62	34.98	34.98	33.16	33.16
Infants with scores below 'seek consultation' cut off at 4 months	N	0	10	2	30	1	13	0	10	0	12
	%	0	1.6	2.3	4.8	1.1	2.1	0	1.6	0	1.9
10 months											
Active clients at 10 months	N	88	667	88	667	88	667	88	667	88	667
Number of infants for active clients at 10 months	N	89	671	89	671	89	671	89	671	89	671
Infants with ASQ data recorded at 10 months	N	81	543	81	543	81	543	81	543	81	543
	%	91.0	80.9	91.0	80.9	91.0	80.9	91.0	80.9	91.0	80.9
Mean ASQ 10 month score	N	54.8	55.7	51.0	50.1	57.7	56.0	55.0	54.7	54.7	53.9
Seek consultation' cut off scores at 10 months		22.87	22.87	30.07	30.07	37.97	37.97	32.51	32.51	27.25	27.25
Infants with scores below 'seek consultation' cut off at 10 months	N	0	7	11	86	1	19	4	20	1	9
	%	0	1.3	13.6	15.8	1.2	3.5	4.9	3.7	1.2	1.7

SEHSCT

4 Months

- Overall, 93 infants reached 4 months infancy by the end of December 2018 and ASQ data was recorded for 94.6% (N=88) of these clients.
- Communication: No infants had a score below the seek consultation cut-off (34.60). The overall mean for communication is 56.6.
- Gross motor: 2.3% (N=2) infants had a score below the seek consultation cut-off (38.41). The overall mean for gross motor is 55.5.
- Fine motor: 1.1% (N=1) infants had a score below the seek consultation cut-off (29.62). The overall mean for fine motor is 54.0.
- Problem solving: No infants had a score below the seek consultation cut-off (34.98). The overall mean for problem solving is 57.6.
- Personal Social: No infants had a score below the seek consultation cut-off (33.16). The overall mean for personal social is 56.1.

10 Months

- Overall, 89 infants reached 10 months infancy by the end of December 2018 and ASQ data was recorded for 91% (N=81) of these clients.
- Communication: No infants had a score below the seek consultation cut-off (22.87). The overall mean for communication is 54.8.
- Gross motor: 13.6% (N=11) infants had a score below the seek consultation cut-off (30.07). The overall mean for gross motor is 51.0.
- Fine motor: 1.2% (N=1) infants had a score below the seek consultation cut-off (37.97). The overall mean for fine motor is 57.7.
- Problem solving: 4.9% (N=4) infants had a score below the seek consultation cut-off (32.51). The overall mean for problem solving is 55.0.
- Personal Social: 1.2% (N=1) infants had a score below the seek consultation cut-off (27.25). The overall mean for personal social is 54.7.

Table 118: ASQ Infancy – SHSCT and all sites

ASQ Infancy		Communication		Gross Motor		Fine Motor		Problem Solving		Personal Social	
		SHSCT	All site total	SHSCT	All site total	SHSCT	All site total	SHSCT	All site total	SHSCT	All site total
Clients ever enrolled		192	913	192	913	192	913	192	913	192	913
4 months (I16 visit)											
Active clients at 4 months	N	161	744	161	744	161	744	161	744	161	744
Number of infants for active clients at 4 months*	N	163	748	163	748	163	748	163	748	163	748
Infants with ASQ data recorded at 4 months	N	144	622	144	622	144	622	144	622	144	622
	%	88.3	83.2	88.3	83.2	88.3	83.2	88.3	83.2	88.3	83.2
Mean ASQ 4 month score	N	55.9	56.3	53.9	54.7	53.5	54.8	54.8	56.2	55.0	55.9
Seek consultation' cut off scores at 4 months		34.60	34.60	38.41	38.41	29.62	29.62	34.98	34.98	33.16	33.16
Infants with scores below 'seek consultation' cut off at 4 months	N	0	10	8	30	2	13	1	10	1	12
	%	0	1.6	5.6	4.8	1.4	2.1	0.7	1.6	0.7	1.9
10 months											
Active clients at 10 months	N	147	667	147	667	147	667	147	667	147	667
Number of infants for active clients at 10 months	N	149	671	149	671	149	671	149	671	149	671
Infants with ASQ data recorded at 10 months	N	108	543	108	543	108	543	108	543	108	543
	%	72.5	80.9	72.5	80.9	72.5	80.9	72.5	80.9	72.5	80.9
Mean ASQ 10 month score	N	55.0	55.7	47.6	50.1	54.3	56.0	53.1	54.7	52.4	53.9
Seek consultation' cut off scores at 10 months		22.87	22.87	30.07	30.07	37.97	37.97	32.51	32.51	27.25	27.25
Infants with scores below 'seek consultation' cut off at 10 months	N	0	7	21	86	2	19	3	20	0	9
	%	0	1.3	19.4	15.8	1.9	3.5	2.8	3.7	0	1.7

*One client had twins

SHSCT

4 Months

- Overall, 163 infants reached 4 months infancy by the end of December 2018 and ASQ data was recorded for 88.3% (N=144) of these clients.
- Communication: No infants had a score below the seek consultation cut-off (34.60). The overall mean for communication is 55.9.
- Gross motor: 5.6% (N=8) infants had a score below the seek consultation cut-off (38.41). The overall mean for gross motor is 53.9.
- Fine motor: 1.4% (N=2) infants had a score below the seek consultation cut-off (29.62). The overall mean for fine motor is 53.5.
- Problem solving: 0.7% (N=1) infants had a score below the seek consultation cut-off (34.98). The overall mean for problem solving is 54.8.
- Personal Social: 0.7% (N=1) infants had a score below the seek consultation cut-off (33.16). The overall mean for personal social is 55.0.

10 Months

- Overall, 149 infants reached 10 months infancy by the end of December 2018 and ASQ data was recorded for 72.5% (N=108) of these clients.
- Communication: No infants had a score below the seek consultation cut-off (22.87). The overall mean for communication is 55.0.
- Gross motor: 19.4% (N=21) infants had a score below the seek consultation cut-off (30.07). The overall mean for gross motor is 47.6.
- Fine motor: 1.9% (N=2) infants had a score below the seek consultation cut-off (37.97). The overall mean for fine motor is 54.3.
- Problem solving: 2.8% (N=3) infants had a score below the seek consultation cut-off (32.51). The overall mean for problem solving is 53.1.
- Personal Social: No infants had a score below the seek consultation cut-off (27.25). The overall mean for personal social is 52.4.

Table 119: ASQ Infancy – WHSCT and all sites

ASQ Infancy		Communication		Gross Motor		Fine Motor		Problem Solving		Personal Social	
		WHSCT	All site total	WHSCT	All site total	WHSCT	All site total	WHSCT	All site total	WHSCT	All site total
Clients ever enrolled		289	913	289	913	289	913	289	913	289	913
4 months (I16 visit)											
Active clients at 4 months	N	252	744	252	744	252	744	252	744	252	744
Number of infants for active clients at 4 months	N	252	748	252	748	252	748	252	748	252	748
Infants with ASQ data recorded at 4 months	N	201	622	201	622	201	622	201	622	201	622
	%	79.8	83.2	79.8	83.2	79.8	83.2	79.8	83.2	79.8	83.2
Mean ASQ 4 month score	N	56.1	56.3	54.1	54.7	55.5	54.8	56.4	56.2	55.5	55.9
Seek consultation' cut off scores at 4 months		34.60	34.60	38.41	38.41	29.62	29.62	34.98	34.98	33.16	33.16
Infants with scores below 'seek consultation' cut off at 4 months	N	6	10	13	30	7	13	7	10	8	12
	%	3.0	1.6	6.5	4.8	3.5	2.1	3.5	1.6	4.0	1.9
10 months											
Active clients at 10 months	N	228	667	228	667	228	667	228	667	228	667
Number of infants for active clients at 10 months	N	228	671	228	671	228	671	228	671	228	671
Infants with ASQ data recorded at 10 months	N	193	543	193	543	193	543	193	543	193	543
	%	84.6	80.9	84.6	80.9	84.6	80.9	84.6	80.9	84.6	80.9
Mean ASQ 10 month score	N	56.9	55.7	51.7	50.1	56.9	56.0	56.4	54.7	55.3	53.9
Seek consultation' cut off scores at 10 months		22.87	22.87	30.07	30.07	37.97	37.97	32.51	32.51	27.25	27.25
Infants with scores below 'seek consultation' cut off at 10 months	N	4	7	26	86	9	19	4	20	4	9
	%	2.1	1.3	13.5	15.8	4.7	3.5	2.1	3.7	2.1	1.7

WHSCCT

4 Months

- Overall, 252 infants reached 4 months infancy by the end of December 2018 and ASQ data was recorded for 79.8% (N=201) of these clients.
- Communication: 3% (N=6) infants had a score below the seek consultation cut-off (34.60). The overall mean for communication is 56.1.
- Gross motor: 6.5% (N=13) infants had a score below the seek consultation cut-off (38.41). The overall mean for gross motor is 54.1.
- Fine motor: 3.5% (N=7) infants had a score below the seek consultation cut-off (29.62). The overall mean for fine motor is 55.5.
- Problem solving: 3.5% (N=7) infants had a score below the seek consultation cut-off (34.98). The overall mean for problem solving is 56.4.
- Personal Social: 4% (N=8) infants had a score below the seek consultation cut-off (33.16). The overall mean for personal social is 55.5.

10 Months

- Overall, 228 infants reached 10 months infancy by the end of December 2018 and ASQ data was recorded for 84.6% (N=193) of these clients.
- Communication: 2.1% (N=4) infants had a score below the seek consultation cut-off (22.87). The overall mean for communication is 56.9.
- Gross motor: 13.5% (N=26) infants had a score below the seek consultation cut-off (30.07). The overall mean for gross motor is 51.7.
- Fine motor: 4.7% (N=9) infants had a score below the seek consultation cut-off (37.97). The overall mean for fine motor is 56.9.
- Problem solving: 2.1% (N=4) infants had a score below the seek consultation cut-off (32.51). The overall mean for problem solving is 56.4.
- Personal Social: 2.1% (N=4) infants had a score below the seek consultation cut-off (27.25). The overall mean for personal social is 55.3.

ASQ Toddlerhood

Table 120: ASQ Toddlerhood – All sites

ASQ Toddlerhood		Communication	Gross Motor	Fine Motor	Problem Solving	Personal Social
14 months						
Active clients at 14 months	N	578	578	578	578	578
Number of infants for active clients at 14 months*	N	580	580	580	580	580
Infants with ASQ data recorded at 14 months	N	426	426	426	426	426
	%	73.4	73.4	73.4	73.4	73.4
Mean ASQ 14 month score	N	53.5	54.5	54.4	52.6	55.4
Seek consultation' cut off scores at 14 months		17.40	25.8	23.06	22.56	23.18
Infants with scores below 'seek consultation' cut off at 14 months	N	10	21	11	12	7
	%	2.3	5.3	2.5	2.8	1.6
20 months						
Active clients at 20 months	N	508	508	508	508	508
Number of infants for active clients at 20 months*	N	510	510	510	510	510
Infants with ASQ data recorded at 20 months	N	402	402	402	402	402
	%	78.8	78.8	78.8	78.8	78.8
Mean ASQ 20 month score	N	52.5	56.3	55.6	53.4	55.6
Seek consultation' cut off scores at 20 months		20.50	39.89	36.05	28.84	33.36
Infants with scores below 'seek consultation' cut off at 20 months	N	21	13	11	7	11
	%	5.2	3.2	2.7	1.7	2.7

*Two clients had twins

All Sites

14 Months

- Overall, 580 infants reached 14 months toddlerhood by the end of December 2018 and ASQ data was recorded for 73.4% (N=426) of these clients.
- Communication: 2.3% (N=10) infants had a score below the seek consultation cut-off (17.40). The overall mean for communication is 53.5.
- Gross motor: 5.3% (N=21) infants had a score below the seek consultation cut-off (25.8). The overall mean for gross motor is 54.5.
- Fine motor: 2.5% (N=11) infants had a score below the seek consultation cut-off (23.06). The overall mean for fine motor is 54.4.
- Problem solving: 2.8% (N=12) infants had a score below the seek consultation cut-off (22.56). The overall mean for problem solving is 52.6.
- Personal Social: 1.6% (N=7) infants had a score below the seek consultation cut-off (23.18). The overall mean for personal social is 55.4.

20 Months

- Overall, 510 infants reached 20 months toddlerhood by the end of December 2018 and ASQ data was recorded for 78.8% (N=402) of these clients.
- Communication: 5.2% (N=21) infants had a score below the seek consultation cut-off (20.50). The overall mean for communication is 52.5.
- Gross motor: 3.2% (N=13) infants had a score below the seek consultation cut-off (39.89). The overall mean for gross motor is 56.3.
- Fine motor: 2.7% (N=11) infants had a score below the seek consultation cut-off (36.05). The overall mean for fine motor is 55.6.
- Problem solving: 1.7% (N=7) infants had a score below the seek consultation cut-off (28.84). The overall mean for problem solving is 53.4.
- Personal Social: 2.7% (N=11) infants had a score below the seek consultation cut-off (33.36). The overall mean for personal social is 55.6.

Table 121: ASQ Toddlerhood – BHSCT and all sites

ASQ Toddlerhood		Communication		Gross Motor		Fine Motor		Problem Solving		Personal Social	
		BHSCT	All site total	BHSCT	All site total	BHSCT	All site total	BHSCT	All site total	BHSCT	All site total
14 months (T41 visit)											
Active clients at 14 months	N	118	578	118	578	118	578	118	578	118	578
Number of infants for active clients at 14 months*	N	119	580	119	580	119	580	119	580	119	580
Infants with ASQ data recorded at 14 months	N	105	426	105	426	105	426	105	426	105	426
	%	88.2	73.4	88.2	73.4	88.2	73.4	88.2	73.4	88.2	73.4
Mean ASQ 14 month score	N	51.9	53.5	56.5	54.5	52.3	54.4	51.0	52.6	55.5	55.4
Seek consultation' cut off scores at 14 months		17.40	17.40	25.8	25.8	23.06	23.06	22.56	22.56	23.18	23.18
Infants with scores below 'seek consultation' cut off at 14 months	N	5	10	2	21	4	11	4	12	2	7
	%	4.8	2.3	2.0	5.3	3.8	2.5	3.8	2.8	1.9	1.6
20 months (T49 visit)											
Active clients at 20 months	N	102	508	102	508	102	508	102	508	102	508
Number of infants for active clients at 20 months	N	103	510	103	510	103	510	103	510	103	510
Infants with ASQ data recorded at 20 months	N	89	402	89	402	89	402	89	402	89	402
	%	86.4	78.8	86.4	78.8	86.4	78.8	86.4	78.8	86.4	78.8
Mean ASQ 20 month score	N	52.0	52.5	56.1	56.3	55.4	55.6	51.6	53.4	54.7	55.6
Seek consultation' cut off scores at 20 months		20.50	20.50	39.89	39.89	36.05	36.05	28.84	28.84	33.36	33.36
Infants with scores below 'seek consultation' cut off at 20 months	N	8	21	5	13	5	11	2	7	6	11
	%	9.0	5.2	5.6	3.2	5.6	2.7	2.2	1.7	6.7	2.7

*Two clients had twins

BHSCT

14 Months

- Overall, 119 infants reached 14 months toddlerhood by the end of December 2018 and ASQ data was recorded for 88.2% (N=105) of these clients.
- Communication: 4.8% (N=5) infants had a score below the seek consultation cut-off (17.40). The overall mean for communication is 51.9.
- Gross motor: 2% (N=2) infants had a score below the seek consultation cut-off (25.8). The overall mean for gross motor is 56.5.
- Fine motor: 3.8% (N=4) infants had a score below the seek consultation cut-off (23.06). The overall mean for fine motor is 52.3.
- Problem solving: 3.8% (N=4) infants had a score below the seek consultation cut-off (22.56). The overall mean for problem solving is 51.0.
- Personal Social: 1.9% (N=2) infants had a score below the seek consultation cut-off (23.18). The overall mean for personal social is 55.5.

20 Months

- Overall, 103 infants reached 20 months toddlerhood by the end of December 2018 and ASQ data was recorded for 86.4% (N=89) of these clients.
- Communication: 9% (N=8) infants had a score below the seek consultation cut-off (20.50). The overall mean for communication is 52.0.
- Gross motor: 5.6% (N=5) infants had a score below the seek consultation cut-off (39.89). The overall mean for gross motor is 56.1.
- Fine motor: 5.6% (N=5) infants had a score below the seek consultation cut-off (36.05). The overall mean for fine motor is 55.4.
- Problem solving: 2.2% (N=2) infants had a score below the seek consultation cut-off (28.84). The overall mean for problem solving is 51.6.
- Personal Social: 6.7% (N=6) infants had a score below the seek consultation cut-off (33.36). The overall mean for personal social is 54.7.

Table 122: ASQ Toddlerhood – NHSCT and all sites

ASQ Toddlerhood		Communication		Gross Motor		Fine Motor		Problem Solving		Personal Social	
		NHSCT	All site total	NHSCT	All site total	NHSCT	All site total	NHSCT	All site total	NHSCT	All site total
14 months (T41 visit)											
Active clients at 14 months	N	54	578	54	578	54	578	54	578	54	578
Number of infants for active clients at 14 months	N	54	580	54	580	54	580	54	580	54	580
Infants with ASQ data recorded at 14 months	N	34	426	34	426	34	426	34	426	34	426
	%	63.0	73.4	63.0	73.4	63.0	73.4	63.0	73.4	63.0	73.4
Mean ASQ 14 month score	N	54.6	53.5	52.0	54.5	56.4	54.4	51.0	52.6	54.7	55.4
Seek consultation' cut off scores at 14 months		17.40	17.40	25.8	25.8	23.06	23.06	22.56	22.56	23.18	23.18
Infants with scores below 'seek consultation' cut off at 14 months	N	0	10	3	21	0	11	0	12	0	7
	%	0	2.3	9.1	5.3	0	2.5	0	2.8	0	1.6
20 months (T49 visit)											
Active clients at 20 months	N	41	508	41	508	41	508	41	508	41	508
Number of infants for active clients at 20 months	N	41	510	41	510	41	510	41	510	41	510
Infants with ASQ data recorded at 20 months	N	26	402	26	402	26	402	26	402	26	402
	%	63.4	78.8	63.4	78.8	63.4	78.8	63.4	78.8	63.4	78.8
Mean ASQ 20 month score	N	53.1	52.5	57.7	56.3	56.2	55.6	51.5	53.4	55.0	55.6
Seek consultation' cut off scores at 20 months		20.50	20.50	39.89	39.89	36.05	36.05	28.84	28.84	33.36	33.36
Infants with scores below 'seek consultation' cut off at 20 months	N	1	21	0	13	0	11	0	7	1	11
	%	3.8	5.2	0	3.2	0	2.7	0	1.7	3.8	2.7

NHSCT

14 Months

- Overall, 54 infants reached 14 months toddlerhood by the end of December 2018 and ASQ data was recorded for 63.0% (N=34) of these clients.
- Communication: No infants had a score below the seek consultation cut-off (17.40). The overall mean for communication is 54.6.
- Gross motor: 9.1% (N=3) infants had a score below the seek consultation cut-off (25.8). The overall mean for gross motor is 52.0.
- Fine motor: No infants had a score below the seek consultation cut-off (23.06). The overall mean for fine motor is 56.4.
- Problem solving: No infants had a score below the seek consultation cut-off (22.56). The overall mean for problem solving is 51.0.
- Personal Social: No infants had a score below the seek consultation cut-off (23.18). The overall mean for personal social is 54.7.

20 Months

- Overall, 41 infants reached 20 months toddlerhood by the end of December 2018 and ASQ data was recorded for 63.4% (N=26) of these clients.
- Communication: 3.8% (N=1) infants had a score below the seek consultation cut-off (20.50). The overall mean for communication is 53.1.
- Gross motor: No infants had a score below the seek consultation cut-off (39.89). The overall mean for gross motor is 57.7.
- Fine motor: No infants had a score below the seek consultation cut-off (36.05). The overall mean for fine motor is 56.2.
- Problem solving: No infants had a score below the seek consultation cut-off (28.84). The overall mean for problem solving is 51.5.
- Personal Social: 3.8% (N=1) infants had a score below the seek consultation cut-off (33.36). The overall mean for personal social is 55.0.

Table 123: ASQ Toddlerhood – SEHSCT and all sites

ASQ Toddlerhood		Communication		Gross Motor		Fine Motor		Problem Solving		Personal Social	
		SEHSCT	All site total	SEHSCT	All site total	SEHSCT	All site total	SEHSCT	All site total	SEHSCT	All site total
14 months (T41 visit)											
Active clients at 14 months	N	69	578	69	578	69	578	69	578	69	578
Number of infants for active clients at 14 months	N	69	580	69	580	69	580	69	580	69	580
Infants with ASQ data recorded at 14 months	N	63	426	63	426	63	426	63	426	63	426
	%	91.3	73.4	91.3	73.4	91.3	73.4	91.3	73.4	91.3	73.4
Mean ASQ 14 month score	N	53.9	53.5	54.7	54.5	51.7	54.4	51.8	52.6	54.1	55.4
Seek consultation' cut off scores at 14 months		17.40	17.40	25.82	25.8	23.06	23.06	22.56	22.56	23.18	23.18
Infants with scores below 'seek consultation' cut off at 14 months	N	1	10	2	21	1	11	2	12	1	7
	%	1.6	2.3	3.2	5.3	1.6	2.5	3.2	2.8	1.6	1.6
20 months (T49 visit)											
Active clients at 20 months	N	54	508	54	508	54	508	54	508	54	508
Number of infants for active clients at 20 months	N	54	510	54	510	54	510	54	510	54	510
Infants with ASQ data recorded at 20 months	N	49	402	49	402	49	402	49	402	49	402
	%	90.7	78.8	90.7	78.8	90.7	78.8	90.7	78.8	90.7	78.8
Mean ASQ 20 month score	N	50.4	52.5	55.3	56.3	55.5	55.6	52.8	53.4	55.6	55.6
Seek consultation' cut off scores at 20 months		20.50	20.50	39.89	39.89	36.05	36.05	28.84	28.84	33.36	33.36
Infants with scores below 'seek consultation' cut off at 20 months	N	2	21	3	13	1	11	0	7	0	11
	%	4.1	5.2	6.1	3.2	2.0	2.7	0.0	1.7	0	2.7

SEHSCT

14 Months

- Overall, 69 infants reached 14 months toddlerhood by the end of December 2018 and ASQ data was recorded for 91.3% (N=63) of these clients.
- Communication: 1.6% (N=1) infants had a score below the seek consultation cut-off (17.40). The overall mean for communication is 53.9.
- Gross motor: 3.2% (N=2) infants had a score below the seek consultation cut-off (25.82). The overall mean for gross motor is 54.7.
- Fine motor: 1.6% (N=1) infants had a score below the seek consultation cut-off (23.06). The overall mean for fine motor is 51.7.
- Problem solving: 3.2% (N=2) infants had a score below the seek consultation cut-off (22.56). The overall mean for problem solving is 51.8.
- Personal Social: 1.6% (N=1) infants had a score below the seek consultation cut-off (23.18). The overall mean for personal social is 54.1.

20 Months

- Overall, 54 infants reached 20 months toddlerhood by the end of December 2018 and ASQ data was recorded for 90.7% (N=49) of these clients.
- Communication: 4.1% (N=2) infants had a score below the seek consultation cut-off (20.50). The overall mean for communication is 50.4.
- Gross motor: 6.1% (N=3) infants had a score below the seek consultation cut-off (39.89). The overall mean for gross motor is 55.3.
- Fine motor: 2% (N=1) infants had a score below the seek consultation cut-off (36.05). The overall mean for fine motor is 55.5.
- Problem solving: No infants had a score below the seek consultation cut-off (28.84). The overall mean for problem solving is 52.8.
- Personal Social: No infants had a score below the seek consultation cut-off (33.36). The overall mean for personal social is 55.6.

Table 124: ASQ Toddlerhood – SHSCT and all sites

ASQ Toddlerhood		Communication		Gross Motor		Fine Motor		Problem Solving		Personal Social	
		SHSCT	All site total	SHSCT	All site total	SHSCT	All site total	SHSCT	All site total	SHSCT	All site total
14 months (T41 visit)											
Active clients at 14 months	N	133	578	133	578	133	578	133	578	133	578
Number of infants for active clients at 14 months	N	134	580	134	580	134	580	134	580	134	580
Infants with ASQ data recorded at 14 months	N	71	426	71	426	71	426	71	426	71	426
	%	53.0	73.4	53.0	73.4	53.0	73.4	53.0	73.4	53.0	73.4
Mean ASQ 14 month score	N	50.4	53.5	54.2	54.5	51.5	54.4	49.4	52.6	53.2	55.4
Seek consultation' cut off scores at 14 months		17.40	17.40	25.82	25.8	23.06	23.06	22.56	22.56	23.18	23.18
Infants with scores below 'seek consultation' cut off at 14 months	N	3	10	4	21	4	11	5	12	3	7
	%	4.2	2.3	5.7	5.3	5.6	2.5	7.0	2.8	4.2	1.6
20 months (T49 visit)											
Active clients at 20 months	N	126	508	126	508	126	508	126	508	126	508
Number of infants for active clients at 20 months	N	127	510	127	510	127	510	127	510	127	510
Infants with ASQ data recorded at 20 months	N	92	402	92	402	92	402	92	402	92	402
	%	72.4	78.8	72.4	78.8	72.4	78.8	72.4	78.8	72.4	78.8
Mean ASQ 20 month score	N	50.3	52.5	55.2	56.3	53.0	55.6	51.4	53.4	54.3	55.6
Seek consultation' cut off scores at 20 months		20.50	20.50	39.89	39.89	36.05	36.05	28.84	28.84	33.36	33.36
Infants with scores below 'seek consultation' cut off at 20 months	N	5	21	2	13	2	11	3	7	2	11
	%	5.4	5.2	2.2	3.2	2.2	2.7	3.3	1.7	2.2	2.7

*One client had twins

SHSCT

14 Months

- Overall, 134 infants reached 14 months toddlerhood by the end of December 2018 and ASQ data was recorded for 53% (N=71) of these clients.
- Communication: 4.2% (N=3) infants had a score below the seek consultation cut-off (17.40). The overall mean for communication is 50.4.
- Gross motor: 5.7% (N=4) infants had a score below the seek consultation cut-off (25.82). The overall mean for gross motor is 54.2.
- Fine motor: 5.6% (N=4) infants had a score below the seek consultation cut-off (23.06). The overall mean for fine motor is 51.5.
- Problem solving: 7.0% (N=5) infants had a score below the seek consultation cut-off (22.56). The overall mean for problem solving is 49.4.
- Personal Social: 4.2% (N=3) infants had a score below the seek consultation cut-off (23.18). The overall mean for personal social is 53.2.

20 Months

- Overall, 127 infants reached 20 months toddlerhood by the end of December 2018 and ASQ data was recorded for 72.4% (N=92) of these clients.
- Communication: 5.4% (N=5) infants had a score below the seek consultation cut-off (20.50). The overall mean for communication is 50.3.
- Gross motor: 2.2% (N=2) infants had a score below the seek consultation cut-off (39.89). The overall mean for gross motor is 55.2.
- Fine motor: 2.2% (N=2) infants had a score below the seek consultation cut-off (36.05). The overall mean for fine motor is 53.0.
- Problem solving: 3.3% (N=3) infants had a score below the seek consultation cut-off (28.84). The overall mean for problem solving is 51.4.
- Personal Social: 2.2% (N=2) infants had a score below the seek consultation cut-off (33.36). The overall mean for personal social is 54.3.

Table 125: ASQ Toddlerhood – WHSCT and all sites

ASQ Toddlerhood		Communication		Gross Motor		Fine Motor		Problem Solving		Personal Social	
		WHSCT	All site total	WHSCT	All site total	WHSCT	All site total	WHSCT	All site total	WHSCT	All site total
14 months (T41 visit)											
Active clients at 14 months	N	204	578	204	578	204	578	204	578	204	578
Number of infants for active clients at 14 months	N	204	580	204	580	204	580	204	580	204	580
Infants with ASQ data recorded at 14 months	N	153	426	153	426	153	426	153	426	153	426
	%	75.0	73.4	75.0	73.4	75.0	73.4	75.0	73.4	75.0	73.4
Mean ASQ 14 month score	N	55.7	53.5	53.6	54.5	57.7	54.4	55.8	52.6	56.9	55.4
Seek consultation' cut off scores at 14 months		17.40	17.40	25.82	25.8	23.06	23.06	22.56	22.56	23.18	23.18
Infants with scores below 'seek consultation' cut off at 14 months	N	1	10	10	21	2	11	1	12	1	7
	%	0.7	2.3	7.5	5.3	1.3	2.5	0.7	2.8	0.7	1.6
20 months (T49 visit)											
Active clients at 20 months	N	185	508	185	508	185	508	185	508	185	508
Number of infants for active clients at 20 months	N	185	510	185	510	185	510	185	510	185	510
Infants with ASQ data recorded at 20 months	N	146	402	146	402	146	402	146	402	146	402
	%	78.9	78.8	78.9	78.8	78.9	78.8	78.9	78.8	78.9	78.8
Mean ASQ 20 month score	N	54.8	52.5	57.2	56.3	57.4	55.6	56.3	53.4	57.0	55.6
Seek consultation' cut off scores at 20 months		20.50	20.50	39.89	39.89	36.05	36.05	28.84	28.84	33.36	33.36
Infants with scores below 'seek consultation' cut off at 20 months	N	5	21	3	13	3	11	2	7	2	11
	%	3.4	5.2	2.1	3.2	2.1	2.7	1.4	1.7	1.4	2.7

WHST

14 Months

- Overall, 204 infants reached 14 months toddlerhood by the end of December 2018 and ASQ data was recorded for 75% (N=153) of these clients.
- Communication: 0.7% (N=1) infants had a score below the seek consultation cut-off (17.40). The overall mean for communication is 55.7.
- Gross motor: 7.5% (N=10) infants had a score below the seek consultation cut-off (25.82). The overall mean for gross motor is 53.6.
- Fine motor: 1.3% (N=2) infants had a score below the seek consultation cut-off (23.06). The overall mean for fine motor is 57.7.
- Problem solving: 0.7% (N=1) infants had a score below the seek consultation cut-off (22.56). The overall mean for problem solving is 55.8.
- Personal Social: 0.7% (N=1) infants had a score below the seek consultation cut-off (23.18). The overall mean for personal social is 56.9.

20 Months

- Overall, 185 infants reached 20 months toddlerhood by the end of December 2018 and ASQ data was recorded for 78.9% (N=146) of these clients.
- Communication: 3.4% (N=5) infants had a score below the seek consultation cut-off (20.50). The overall mean for communication is 54.8.
- Gross motor: 2.1% (N=3) infants had a score below the seek consultation cut-off (39.89). The overall mean for gross motor is 57.2.
- Fine motor: 2.1% (N=3) infants had a score below the seek consultation cut-off (36.05). The overall mean for fine motor is 57.4.
- Problem solving: 1.4% (N=2) infants had a score below the seek consultation cut-off (28.84). The overall mean for problem solving is 56.3.
- Personal Social: 1.4% (N=2) infants had a score below the seek consultation cut-off (33.36). The overall mean for personal social is 57.0.

ASQ-SE

Table 126: ASQ-SE 6, 12, 18 and 24 months – All sites

ASQ-SE*				
Clients ever enrolled = 913				
	6 months	12 months	18 months	24 months
Active clients (N)	744	667	578	508
Number of infants for active clients (N)	748	671	580	510
Number of infants with ASQ-SE data recorded (N)	619	535	425	408
Percentage of infants with ASQ-SE data recorded (%)	82.8	79.7	73.3	80.0
Mean ASQ-SE score	7.8	7.6	9.8	11.6
Min ASQ-SE score	0	0	0	0
Max ASQ-SE score	360	60	80	150
'Seek consultation' cut off	45	48	50	50
Number of infants with scores above 'seek consultation' cut off (N)	5	3	6	12
Percentage of infants with a score above 'seek consultation' cut off (%)	0.8	0.6	1.4	2.9

* The ASQ:SE-2 is a set of questionnaires about children's social-emotional development [Squires, J., Bricker, D., & Twombly, E. (2002). Ages and Stages Questionnaires: Social-Emotional (ASQ: SE), A parent-completed, child-monitoring system for social-emotional behaviours. Baltimore: Paul H. Brookes Publishing].

All sites

- Overall, 748 infants reached 6 months infancy by the end of December 2018 and ASQ-SE data was recorded for 82.8% (N=619) of these clients. 0.8% (N=5) infants scored above the seek consultation cut-off (45). The overall mean ASQ-SE score is 7.8.
- Overall, 671 infants reached 12 months infancy by the end of December 2018 and ASQ-SE data was recorded for 79.7% (N=535) of these clients. 0.6% (N=3) infants scored above the seek consultation cut-off (48). The overall mean ASQ-SE score is 7.6.
- Overall, 580 infants reached 18 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 73.3% (N=425) of these clients. 1.4% (N=6) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 9.8.
- Overall, 510 infants reached 24 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 80% (N=408) of these clients. 2.9% (N=12) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 11.6.

Table 127: ASQ-SE 6, 12, 18 and 24 months – BHSCT and all sites

ASQ-SE*								
	6 months		12 months		18 months		24 months	
	BHSCT	All site total	BHSCT	All site total	BHSCT	All site total	BHSCT	All site total
Active clients (N)	153	744	136	667	118	578	102	508
Number of infants for active clients (N)	154	748	137	671	119	580	103	510
Number of infants with ASQ-SE data recorded (N)	135	619	120	535	105	425	89	408
Percentage of infants with ASQ-SE data recorded (%)	87.7	82.8	87.6	79.7	88.2	73.3	86.4	80.0
Mean ASQ-SE score	7.7	7.8	8.5	7.6	12.7	9.8	11.7	11.6
Min ASQ-SE score	0	0	0	0	0	0	0	0
Max ASQ-SE score	180	360	50	60	80	80	95	150
'Seek consultation' cut off	45	45	48	48	50	50	50	50
Number of infants with scores above 'seek consultation' cut off (N)	2	5	1	3	3	6	4	12
Percentage of infants with a score above 'seek consultation' cut off (%)	1.5	0.8	0.8	0.6	2.9	1.4	4.5	2.9

* The ASQ:SE-2 is a set of questionnaires about children's social-emotional development [Squires, J., Bricker, D., & Twombly, E. (2002). Ages and Stages Questionnaires: Social-Emotional (ASQ: SE), A parent-completed, child-monitoring system for social-emotional behaviors. Baltimore: Paul H. Brookes Publishing].

BHSCT

- Overall, 154 infants reached 6 months infancy by the end of December 2018 and ASQ-SE data was recorded for 87.7% (N=135) of these clients. 1.5% (N=2) infants scored above the seek consultation cut-off (45). The overall mean ASQ-SE score is 7.7.
- Overall, 137 infants reached 12 months infancy by the end of December 2018 and ASQ-SE data was recorded for 87.6% (N=120) of these clients. 0.8% (N=1) infants scored above the seek consultation cut-off (48). The overall mean ASQ-SE score is 8.5.
- Overall, 119 infants reached 18 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 88.2% (N=105) of these clients. 2.9% (N=3) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 12.7.
- Overall, 103 infants reached 24 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 86.4% (N=89) of these clients. 4.5% (N=4) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 11.7.

Table 128: ASQ-SE 6, 12, 18 and 24 months – NHSCT and all sites

ASQ-SE*								
	6 months		12 months		18 months		24 months	
	NHSCT	All site total	NHSCT	All site total	NHSCT	All site total	NHSCT	All site total
Active clients (N)	86	744	68	667	54	578	41	508
Number of infants for active clients (N)	86	748	68	671	54	580	41	510
Number of infants with ASQ-SE data recorded (N)	53	619	41	535	35	425	26	408
Percentage of infants with ASQ-SE data recorded (%)	61.6	82.8	60.3	79.7	64.8	73.3	63.4	80.0
Mean ASQ-SE score	16.3	7.8	8.7	7.6	9.7	9.8	18.5	11.6
Min ASQ-SE score	0	0	0	0	0	0	0	0
Max ASQ-SE score	360	360	20	60	30	80	150	150
'Seek consultation' cut off	45	45	48	48	50	50	50	50
Number of infants with scores above 'seek consultation' cut off (N)	1	5	0	3	0	6	2	12
Percentage of infants with a score above 'seek consultation' cut off (%)	1.9	0.8	0	0.6	0	1.4	7.7	2.9

* The ASQ:SE-2 is a set of questionnaires about children's social-emotional development [Squires, J., Bricker, D., & Twombly, E. (2002). Ages and Stages Questionnaires: Social-Emotional (ASQ: SE), A parent-completed, child-monitoring system for social-emotional behaviors. Baltimore: Paul H. Brookes Publishing].

NHSCT

- Overall, 86 infants reached 6 months infancy by the end of December 2018 and ASQ-SE data was recorded for 61.6% (N=53) of these clients. 1.9% (N=1) infants scored above the seek consultation cut-off (45). The overall mean ASQ-SE score is 16.3.
- Overall, 68 infants reached 12 months infancy by the end of December 2018 and ASQ-SE data was recorded for 60.3% (N=41) of these clients. No infants scored above the seek consultation cut-off (48). The overall mean ASQ-SE score is 8.7.
- Overall, 54 infants reached 18 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 64.8% (N=35) of these clients. No infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 9.7.
- Overall, 41 infants reached 24 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 63.4% (N=26) of these clients. 7.7% (N=2) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 18.5.

Table 129: ASQ-SE 6, 12, 18 and 24 months – SEHSCT and all sites

ASQ-SE*								
	6 months		12 months		18 months		24 months	
	SEHSCT	All site total	SEHSCT	All site total	SEHSCT	All site total	SEHSCT	All site total
Active clients (N)	92	744	88	667	69	578	54	508
Number of infants for active clients (N)	93	748	89	671	69	580	54	510
Number of infants with ASQ-SE data recorded (N)	88	619	81	535	63	425	49	408
Percentage of infants with ASQ-SE data recorded (%)	94.6	82.8	91.0	79.7	91.3	73.3	90.7	80.0
Mean ASQ-SE score	6.3	7.8	5.9	7.6	8.2	9.8	12.1	11.6
Min ASQ-SE score	0	0	0	0	0	0	0	0
Max ASQ-SE score	60	360	60	60	60	80	140	150
'Seek consultation' cut off	45	45	48	48	50	50	50	50
Number of infants with scores above 'seek consultation' cut off (N)	2	5	1	3	1	6	2	12
Percentage of infants with a score above 'seek consultation' cut off (%)	2.3	0.8	1.2	0.6	1.6	1.4	4.1	2.9

* The ASQ:SE-2 is a set of questionnaires about children's social-emotional development [Squires, J., Bricker, D., & Twombly, E. (2002). *Ages and Stages Questionnaires: Social-Emotional (ASQ: SE)*, A parent-completed, child-monitoring system for social-emotional behaviors. Baltimore: Paul H. Brookes Publishing].

SEHSCT

- Overall, 93 infants reached 6 months infancy by the end of December 2018 and ASQ-SE data was recorded for 94.6% (N=88) of these clients. 2.3% (N=2) infants scored above the seek consultation cut-off (45). The overall mean ASQ-SE score is 6.3.
- Overall, 89 infants reached 12 months infancy by the end of December 2018 and ASQ-SE data was recorded for 91% (N=81) of these clients. 1.2% (N=1) infants scored above the seek consultation cut-off (48). The overall mean ASQ-SE score is 5.9.
- Overall, 69 infants reached 18 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 91.3% (N=63) of these clients. 1.6% (N=1) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 8.2.
- Overall, 54 infants reached 24 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 90.7% (N=49) of these clients. 4.1% (N=2) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 12.1.

Table 130: ASQ-SE 6, 12, 18 and 24 months – SHSCT and all sites

ASQ-SE *								
	6 months		12 months		18 months		24 months	
	SHSCT	All site total	SHSCT	All site total	SHSCT	All site total	SHSCT	All site total
Active clients (N)	161	744	147	667	133	578	126	508
Number of infants for active clients (N)	163	748	149	671	134	580	127	510
Number of infants with ASQ-SE data recorded (N)	144	619	106	535	68	425	95	408
Percentage of infants with ASQ-SE data recorded (%)	88.3	82.8	71.1	79.7	50.7	73.3	74.8	80.0
Mean ASQ-SE score	12.4	7.8	13.0	7.6	15.9	9.8	15.6	11.6
Min ASQ-SE score	0	0	0	0	0	0	0	0
Max ASQ-SE score	45	360	55	60	60	80	60	150
'Seek consultation' cut off	45	45	48	48	50	50	50	50
Number of infants with scores above 'seek consultation' cut off (N)	0	5	1	3	1	6	2	12
Percentage of infants with a score above 'seek consultation' cut off (%)	0	0.8	0.9	0.6	1.5	1.4	2.1	2.9

* The ASQ:SE-2 is a set of questionnaires about children's social-emotional development [Squires, J., Bricker, D., & Twombly, E. (2002). *Ages and Stages Questionnaires: Social-Emotional (ASQ: SE)*, A parent-completed, child-monitoring system for social-emotional behaviors. Baltimore: Paul H. Brookes Publishing].

SHSCT

- Overall, 163 infants reached 6 months infancy by the end of December 2018 and ASQ-SE data was recorded for 88.3% (N=144) of these clients. No infants scored above the seek consultation cut-off (45). The overall mean ASQ-SE score is 12.4.
- Overall, 149 infants reached 12 months infancy by the end of December 2018 and ASQ-SE data was recorded for 71.1% (N=106) of these clients. 0.9% (N=1) infants scored above the seek consultation cut-off (48). The overall mean ASQ-SE score is 13.0.
- Overall, 134 infants reached 18 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 50.7% (N=68) of these clients. 1.5% (N=1) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 15.9.
- Overall, 127 infants reached 24 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 74.8% (N=95) of these clients. 2.1% (N=2) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 15.6.

Table 131: ASQ-SE 6, 12, 18 and 24 months – WHSCT and all sites

ASQ-SE *								
	6 months		12 months		18 months		24 months	
	WHSCT	All site total	WHSCT	All site total	WHSCT	All site total	WHSCT	All site total
Active clients (N)	252	744	228	667	204	578	185	508
Number of infants for active clients (N)	252	748	228	671	204	580	185	510
Number of infants with ASQ-SE data recorded (N)	199	619	187	535	154	425	149	408
Percentage of infants with ASQ-SE data recorded (%)	79.0	82.8	82.0	79.7	75.5	73.3	80.5	80.0
Mean ASQ-SE score	2.4	7.8	4.5	7.6	5.8	9.8	7.5	11.6
Min ASQ-SE score	0	0	0	0	0	0	0	0
Max ASQ-SE score	25	360	35	60	65	80	60	150
'Seek consultation' cut off	45	45	48	48	50	50	50	50
Number of infants with scores above 'seek consultation' cut off (N)	0	5	0	3	1	6	2	12
Percentage of infants with a score above 'seek consultation' cut off (%)	0	0.8	0	0.6	0.6	1.4	1.3	2.9

* The ASQ:SE-2 is a set of questionnaires about children's social-emotional development [Squires, J., Bricker, D., & Twombly, E. (2002). Ages and Stages Questionnaires: Social-Emotional (ASQ: SE), A parent-completed, child-monitoring system for social-emotional behaviors. Baltimore: Paul H. Brookes Publishing].

WHSCT

- Overall, 252 infants reached 6 months infancy by the end of December 2018 and ASQ-SE data was recorded for 79% (N=199) of these clients. No infants scored above the seek consultation cut-off (45). The overall mean ASQ-SE score is 2.4.
- Overall, 228 infants reached 12 months infancy by the end of December 2018 and ASQ-SE data was recorded for 82% (N=187) of these clients. No infants scored above the seek consultation cut-off (48). The overall mean ASQ-SE score is 4.5.
- Overall, 204 infants reached 18 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 75.5% (N=154) of these clients. 0.6% (N=1) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 5.8.
- Overall, 185 infants reached 24 months toddlerhood by the end of December 2018 and ASQ-SE data was recorded for 80.5% (N=149) of these clients. 1.3% (N=2) infants scored above the seek consultation cut-off (50). The overall mean ASQ-SE score is 7.5.

Immunisations

Immunisations 6 Months

Table 132: Immunisations 6 Months – All sites

Site	6 months										
	Total number enrolled	Number of active clients	Number of infants (including twins)	Infants with immunisation status recorded		Infants with written record of up-to-date immunisations		Infants with up-to-date immunisations – written record and mother’s self-report		Infants recorded as not up-to-date with immunisations	
				N	%	N	%	N	%	N	%
BHSCT	184	153	154	138	89.6	45	32.6	133	96.4	5	3.6
NHSCT	110	86	86	56	65.1	21	37.5	51	91.1	5	8.9
SEHSCT	138	92	93	88	94.6	36	40.9	85	96.6	3	3.4
SHSCT	192	161	163	145	89.0	63	43.4	139	95.9	6	4.1
WHSCT	289	252	252	205	81.3	91	44.4	192	93.7	13	6.3
All sites	913	744	748	632	84.5	256	40.5	600	94.9	32	5.1

This table reports on immunisations for infants reaching 6 months (N=748) by 31.12.18.

All sites

- Overall, 748 infants reached 6 months infancy and immunisation status was recorded for 84.5% (N=632) of these clients. 94.9% (N= 600) had up-to-date immunisations (written record or mother’s self-report) and 5.1% (N=32) did not have records reported as up-to-date.

BHSCT

- In the BHSCT, 154 infants reached 6 months infancy and immunisation status was recorded for 89.6% (N=138) of these clients.
- 96.4% (N=133) had up-to-date immunisations (written record or mother’s self-report) and 3.6% (N=5) did not have records reported as up-to-date.

NHSCT

- In the NHSCT, 86 infants reached 6 months infancy and immunisation status was recorded for 65.1% (N=56) of these clients.
- 91.1% (N=51) had up-to-date immunisations (written record or mother's self-report) and 8.9% (N=5) did not have records reported as up-to-date.

SEHSCT

- In the SEHSCT, 93 infants reached 6 months infancy and immunisation status was recorded for 94.6% (N=88) of these clients.
- 96.6% (N=85) had up-to-date immunisations (written record or mother's self-report) and 3.4% (N=3) did not have records reported as up-to-date.

SHSCT

- In the SHSCT, 163 infants reached 6 months infancy and immunisation status was recorded for 89% (N=145) of these clients.
- 95.9% (N=139) had up-to-date immunisations (written record or mother's self-report) and 4.1% (N=6) did not have records reported as up-to-date.

WHSCT

- In the WHSCT, 252 infants reached 6 months infancy and immunisation status was recorded for 81.3% (N=205) of these clients.
- 93.7% (N=192) had up-to-date immunisations (written record or mother's self-report) and 6.3% (N=13) did not have records reported as up-to-date.

Immunisations 12 months

Table 133: Immunisations 12 months – All sites

Site	12 months										
	Total number ever enrolled	Number of active clients	Number of infants (including twins)	Infants with immunisation status recorded		Infants with written record of up-to-date immunisations		Infants with up-to-date immunisations – written record and mother’s self-report		Infants recorded as not up-to-date with immunisations	
				N	%	N	%	N	%	N	%
BHSCT	184	136	137	122	89.1	23	18.9	111	91.0	11	9.0
NHSCT	110	68	68	41	60.3	11	26.8	40	97.6	1	2.4
SEHSCT	138	88	89	79	88.8	26	32.9	73	92.4	6	7.6
SHSCT	192	147	149	110	73.8	43	39.1	101	91.8	9	8.2
WHSCT	289	228	228	196	86.0	68	34.7	187	95.4	9	4.6
All sites	913	667	671	548	81.7	171	31.2	512	93.4	36	6.6

This table reports on immunisations for infants reaching 12 months (N=671) by 31.12.18.

All sites

- Overall, 671 infants reached 12 months infancy and immunisation status was recorded for 81.7% (N=548) of these clients.
- 93.4% (N=512) had up-to-date immunisations (written record or mother’s self-report) and 6.6% (N=36) did not have records recorded as up-to-date.

BHSCT

- In the BHSCT, 137 infants reached 12 months infancy and immunisation status was recorded for 89.1% (N=122) of these clients.
- 91% (N=111) had immunisations (written record or mother’s self-report) recorded as up-to-date and 9% (N=11) did not have records recorded as up-to-date.

NHSCT

- In the NHSCT, 68 infants reached 12 months infancy and immunisation status was recorded for 60.3% (N=41) of these clients.
- 97.6% (N=40) had immunisations (written record or mother’s self-report) recorded as up-to-date and 2.4% (N=1) did not have records recorded as up-to-date.

SEHSCT

- In the SEHSCT, 89 infants reached 12 months infancy and immunisation status was recorded for 88.8% (N=79) of these clients
- 92.4% (N=73) had immunisations (written record or mother's self-report) recorded as up-to-date and 7.6% (N=6) did not have records recorded as up-to-date.

SHSCT

- In the SHSCT, 149 infants reached 12 months infancy and immunisation status was recorded for 73.8% (N=110) of these clients
- 91.8% (N=101) had immunisations (written record or mother's self-report) recorded as up-to-date and 8.2% (N=9) did not have records reported as up-to-date.

WHSCT

- In the WHSCT, 228 infants reached 12 months infancy and immunisation status was recorded for 86% (N=196) of these clients
- 95.4% (N=187) had immunisations (written record or mother's self-report) recorded as up-to-date and 4.6% (N=9) did not have records reported as up-to-date.

Immunisations 18 months

Table 134: Immunisations 18 months – All sites

18 months											
Site	Total number ever enrolled	Number of active clients	Number of infants (including twins)	Infants with immunisation status recorded		Infants with written record of up-to-date immunisations		Infants with up-to-date immunisations – written record and mother's self-report		Infants recorded as not up-to-date with immunisations	
	N	N	N	N	%	N	%	N	%	N	%
BHSCT	184	118	119	105	88.2	25	23.8	101	96.2	4	3.8
NHSCT	110	54	54	35	64.8	13	37.1	34	97.1	1	2.9
SEHSCT	138	69	69	63	91.3	8	12.7	60	95.2	3	4.8
SHSCT	192	133	134	70	52.2	24	34.3	67	95.7	3	4.3
WHSCT	289	204	204	160	78.4	58	36.3	157	98.1	3	1.9
All sites	913	578	580	433	74.7	128	29.6	419	96.8	14	3.2

This table reports on immunisations for infants reaching 18 months (N=580) by 31.12.18.

All sites

- Overall, 580 infants reached 18 months toddlerhood and immunisation status was recorded for 74.7% (N=433) of these clients. 96.8% (N=419) had up-to-date immunisations (written record or mother's self-report) and 3.2% (N=14) did not have records reported as up-to-date.

BHSCT

- In the BHSCT, 119 infants reached 18 months toddlerhood and immunisation status was recorded for 88.2% (N=105) of these clients.
- 96.2% (N=101) had immunisations (written record or mother's self-report) recorded as up-to-date and 3.8% (N=4) did not have records reported as up-to-date.

NHSCT

- In the NHSCT, 54 infants reached 18 months toddlerhood and immunisation status was recorded for 64.8% (N=35) of these clients.
- 97.1% (N=34) had immunisations (written record or mother's self-report) recorded as up-to-date and 2.9% (N=1) did not have records reported as up-to-date.

SEHSCT

- In the SEHSCT, 69 infants reached 18 months toddlerhood and immunisation status was recorded for 91.3% (N=63) of these clients.
- 95.2% (N=60) had immunisations (written record or mother's self-report) recorded as up-to-date and 4.8% (N=3) did not have records reported as up-to-date.

SHSCT

- In the SHSCT, 134 infants reached 18 months toddlerhood and immunisation status was recorded for 52.2% (N=70) of these clients.
- 95.7% (N=67) had immunisations (written record or mother's self-report) recorded as up-to-date and 4.3% (N=3) did not have records reported as up-to-date.

WHSCT

- In the WHSCT, 204 infants reached 18 months toddlerhood and immunisation status was recorded for 78.4% (N=160) of these clients.
- 98.1% (N=157) had immunisations (written record or mother's self-report) recorded as up-to-date and 1.9% (N=3) did not have records reported as up-to-date.

Immunisations 24 months

Table 135: Immunisations 24 months – All sites

24 months											
Site	Total number ever enrolled	Number of active clients	Number of infants (including twins)	Infants with immunisation status recorded		Infants with written record of up-to-date immunisations		Infants with up-to-date immunisations – written record and mother’s self-report		Infants recorded as not up-to-date with immunisations	
	N	N	N	N	%	N	%	N	%	N	%
BHSCT	184	102	103	89	86.4	24	27.0	87	97.8	2	2.2
NHSCT	110	41	41	26	63.4	7	26.9	26	100	0	0
SEHSCT	138	54	54	49	90.7	12	24.5	49	100	0	0
SHSCT	192	126	127	96	75.6	35	36.5	93	96.9	3	3.1
WHSCT	289	185	185	157	84.9	60	38.2	156	99.4	1	0.6
All sites	913	508	510	417	81.8	138	33.1	411	98.6	6	1.4

This table reports on immunisations for infants reaching 24 months (N=510) by 31.12.18.

All sites

- Overall, 510 infants reached 24 months toddlerhood and immunisation status was recorded for 81.8% (N=417) of these clients. 98.6% (N=411) had up-to-date immunisations (written record or mother’s self-report) and 1.4% (N=6) did not have records reported as up-to-date.

BHSCT

- In the BHSCT, 103 infants reached 24 months toddlerhood and immunisation status was recorded for 86.4% (N=89) of these clients.
- 97.8% (N=87) had immunisations (written record or mother’s self-report) recorded as up-to-date and 2.2% (N=2) did not have records reported as up-to-date.

NHSCT

- In the NHSCT, 41 infants reached 24 months toddlerhood and immunisation status was recorded for 63.4% (N=26) of these clients.
- 100% (N=26) had immunisations (written record or mother’s self-report) recorded as up-to-date.

SEHSCT

- In the SEHSCT, 54 infants reached 24 months toddlerhood and immunisation status was recorded for 90.7% (N=49) of these clients.
- 100% (N=49) had immunisations (written record or mother's self-report) recorded as up-to-date.

SHSCT

- In the SHSCT, 127 infants reached 24 months toddlerhood and immunisation status was recorded for 75.6% (N=96) of these clients.
- 96.9% (N=93) had immunisations (written record or mother's self-report) recorded as up-to-date and 3.1% (N=3) did not have records reported as up-to-date.

WHSCT

- In the WHSCT, 185 infants reached 24 months toddlerhood and immunisation status was recorded for 84.9% (N=157) of these clients.
- 99.4% (N=156) had immunisations (written record or mother's self-report) recorded as up-to-date and 0.6% (N=1) did not have records reported as up-to-date.